

PUBLISHED VERSION

Alistair M. Senior, Mathieu Lihoreau, Michael A. Charleston, Jerome Buhl, David Raubenheimer and Stephen J. Simpson

Adaptive collective foraging in groups with conflicting nutritional needs

Royal Society Open Science, 2016; 3(4):150638-1-150638-15

© 2016 The Authors. Published by the Royal Society under the terms of the Creative Commons Attribution License <http://creativecommons.org/licenses/by/4.0/>, which permits unrestricted use, provided the original author and source are credited.

Originally published at:

<http://doi.org/10.1098/rsos.150638>

PERMISSIONS

<http://creativecommons.org/licenses/by/4.0/>



Attribution 4.0 International (CC BY 4.0)

This is a human-readable summary of (and not a substitute for) the [license](#).

[Disclaimer](#)



You are free to:

Share — copy and redistribute the material in any medium or format

Adapt — remix, transform, and build upon the material

for any purpose, even commercially.

The licensor cannot revoke these freedoms as long as you follow the license terms.

Under the following terms:



Attribution — You must give **appropriate credit**, provide a link to the license, and **indicate if changes were made**. You may do so in any reasonable manner, but not in any way that suggests the licensor endorses you or your use.

No additional restrictions — You may not apply legal terms or **technological measures** that legally restrict others from doing anything the license permits.

7 December 2016

<http://hdl.handle.net/2440/101604>



Cite this article: Senior AM, Lihoreau M, Charleston MA, Buhl J, Raubenheimer D, Simpson SJ. 2016 Adaptive collective foraging in groups with conflicting nutritional needs. *R. Soc. open sci.* **3**: 150638. <http://dx.doi.org/10.1098/rsos.150638>

Received: 23 November 2015

Accepted: 15 March 2016

Subject Category:

Biology (whole organism)

Subject Areas:

behaviour/ecology/computational biology

Keywords:

foraging, individual-based model, nutritional geometry, collective decisions, social interactions, sociality

Author for correspondence:

Alistair M. Senior

e-mail: alistair.senior@sydney.edu.au

Adaptive collective foraging in groups with conflicting nutritional needs

Alistair M. Senior^{1,2}, Mathieu Lihoreau⁵, Michael A. Charleston⁶, Jerome Buhl⁷, David Raubenheimer^{1,3,4} and Stephen J. Simpson^{1,3}

¹Charles Perkins Centre, ²School of Mathematics and Statistics, ³School of Life and Environmental Sciences, and ⁴Faculty of Veterinary Science, The University of Sydney, Sydney, New South Wales 2006, Australia

⁵Research Center on Animal Cognition (CRCA), Center for Integrative Biology (CBI), Toulouse University, CNRS, UPS, France

⁶School of Physical Sciences, University of Tasmania, Hobart, Tasmania 7005, Australia

⁷School of Agriculture, Food and Wine, The University of Adelaide, Adelaide, South Australia 5005, Australia

Collective foraging, based on positive feedback and quorum responses, is believed to improve the foraging efficiency of animals. Nutritional models suggest that social information transfer increases the ability of foragers with closely aligned nutritional needs to find nutrients and maintain a balanced diet. However, whether or not collective foraging is adaptive in a heterogeneous group composed of individuals with differing nutritional needs is virtually unexplored. Here we develop an evolutionary agent-based model using concepts of nutritional ecology to address this knowledge gap. Our aim was to evaluate how collective foraging, mediated by social retention on foods, can improve nutrient balancing in individuals with different requirements. The model suggests that in groups where inter-individual nutritional needs are unimodally distributed, high levels of collective foraging yield optimal individual fitness by reducing search times that result from moving between nutritionally imbalanced foods. However, where nutritional needs are highly bimodal (e.g. where the requirements of males and females differ) collective foraging is selected against, leading to group fission. In this case, additional mechanisms such as assortative interactions can coevolve to allow collective foraging by subgroups

Electronic supplementary material is available at <http://dx.doi.org/10.1098/rsos.150638> or via <http://rsos.royalsocietypublishing.org>.

of individuals with aligned requirements. Our findings indicate that collective foraging is an efficient strategy for nutrient regulation in animals inhabiting complex nutritional environments and exhibiting a range of social forms.

1. Introduction

Many species live in heterogeneous environments in which essential resources are patchily distributed. Foraging decisions, which have a large impact on fitness, can be especially complex, leading some animals to exploit social information emanating from their conspecifics in locating and selecting foods [1,2]. In gregarious animals, such as many insects, fish, birds and ungulates, social information transfer may result in collective foraging decisions whereby all (or most) individuals in the group decide to exploit the same food resource from several available alternatives [3–5]. Typically, these collective dynamics are driven by positive feedback and quorum responses, whereby the probability of an individual choosing a resource varies positively and nonlinearly with the number of individuals already exploiting that resource [4,6]. Through these processes, groups often make faster and/or more accurate decisions than isolated animals, a phenomenon known as swarm intelligence [4,6]. To date, research on collective foraging behaviour has largely focused on identifying the mechanisms that underpin collective decision-making. However, little is known about the evolutionary roots of this widespread phenomenon [7].

Insightful data come from studies on insects, which exemplify how relatively simple social interactions, such as social retention on, or attraction to, foods based on the number of conspecifics already exploiting that food, can impact the foraging decisions and efficiency of individuals [6,8–10]. Informal comparisons of experimental data suggest that the fitness benefits of collective foraging depend on a subtle interplay between the strength of social effects in a species and the availability of resources in their environment. For instance, studies on domiciliary cockroaches (*Blattella germanica*) illustrate how the number of conspecifics already feeding on a food can influence the choice of an individual, with larger groups being more likely to attract new recruits [11,12]. It is hypothesized that this simple mechanism provides an individual cockroach with an honest signal about the quantity and/or quality of a food source enabling grouped cockroaches to make more accurate decisions than isolated conspecifics [12]. By contrast, experiments in tent caterpillars (*Malacosoma disstria*) indicate that the strength of social retention may compromise the quality of nutritional decisions, which individuals alone make with some accuracy [13]. These differential findings highlight an interesting potential trade-off between group foraging, which may increase the efficiency of individual decision-making, and the specific nutritional needs of the individual.

Conceptual progressions in nutrition research show that foraging decisions are intrinsically complicated by the fact that individuals must take into account their needs for multiple nutrients, which may be contained in differing amounts and ratios in those foods available [14]. Ultimately studies on the evolution of collective foraging decisions must capture the complex multidimensional nature of nutrition, rather than solely focusing on the acquisition of a single resource (e.g. energy [15]). Recently, Lihoreau *et al.* [16] developed an agent-based model (ABM) derived from nutritional geometry, a state-space modelling framework for conceptualizing the nutritional decisions of animals (figure 1*a* and box 1) [14,17,18] to explore the efficiency of collective foraging in complex multi-nutrient environments. Their model suggests that an optimal level of social retention (termed K_{soc}), whereby foragers are more likely to remain on and eat from heavily occupied food sources, improves the nutritional performance of individuals. This simple form of information transfer is sufficient to enable individuals to efficiently comprise a balanced diet from individually imbalanced, but collectively complementary, foods [18]. While this is an important first step, this approach overlooks inter-individual variation in nutritional requirements, which is likely to be present in most animal groups (cf. [15]). A recent meta-analysis exemplifies the probable ubiquity of such variability in species from different trophic levels and a wide range of taxonomic groups [24]. In part, this variation may be readily predictable based on phenotype, for example if the nutrient requirements of an individual vary with age or sex [14,25]. However, even where groups of individuals appear to be outwardly homogeneous, as for instance in a cohort of same sex individuals, heterogeneity in other traits such as metabolic rate may result in variance in nutritional requirements [24]. Given that the nutritional needs of all individuals in a group may never be perfectly aligned, does collective foraging still improve the foraging efficiency of individuals?




Figure 1. Nutritional geometry models. (a) Nutritional geometry (see box 1) model for two nutrients, e.g. protein on the x -axis and carbohydrate on the y -axis. In this example, the individuals' requirements are given by a single coordinate known as the intake target (IT; grey cross hair). The environment contains two foods. Food A is rich in carbohydrate (10 parts to each part protein) and food B is rich in protein. An individual has been able to move its nutritional state (NS) towards the IT, by eating food A for meal 1 and food B for meal 2 (sequence of black arrows). (b) While a single IT may adequately represent the mean requirements of all individuals in a group (grey cross hair), in reality there is likely to be within-group variance, with each individual having its own IT (for a hypothetical group of 20 individuals given here by black points). (c) In some instances, variance in nutritional requirements may be predictable. For example, if the group is composed of individuals with two phenotypes with distinct nutritional requirements (e.g. male and female), individual ITs (black points) may be bimodally distributed in two subgroups around an overall mean (grey cross hair).

Box 1. Key principles of nutritional geometry.

Nutritional geometry is a state-based modelling approach for studying the nutritional strategies of animals, based on graphic representations of individuals, their nutritional requirements, foods and the interactions thereof in a geometric space [14,17,18]. Over recent years, this conceptual framework has become increasingly used to study how animals regulate their acquisition of multiple nutrients simultaneously, and how this varies across feeding guilds, trophic levels and taxonomic groups. In the most basic nutritional geometry models, two nutrients (such as the macronutrients protein and carbohydrate) are depicted in a two-dimensional Cartesian coordinate system forming a nutrient space (see example in figure 1a). An individual's nutritional state (NS) is denoted by its (x, y) coordinates, and moves through the nutrient space when foods are eaten. Foods are radials projecting through the nutrient space at angles from the origin determined by the ratio of the component nutrients contained (nutritional rails; figure 1a). The nutritional requirements of an individual are given by a single coordinate or a broader region within the nutrient space known as the intake target (IT, figure 1a) [14]. Multiple ITs may maximize different life-history traits in the same individual (e.g. maximize growth, reproduction or longevity [19,20]). However, evolutionary theory and experimental evidence suggest that animals evolve strategies that attempt to reach the IT that maximizes overall evolutionary fitness [14,19,21,22]. While most previous nutritional geometry models have assumed that a single IT would maximize the fitness of all individuals within a group or population (e.g. [23]), it is becoming increasingly clear that inter-individual variation in ITs is abundant (figure 1b) [24]. In the relatively homogeneous groups, this variation may be captured by a unimodal distribution of individual ITs surrounding the group mean IT (figure 1b). In more heterogeneous groups, these ITs may have discrete distributions, separating the group into distinct subgroups, as for example, in the case of sex-specific nutritional needs where the ITs of mixed-sex groups may be bimodally distributed (figure 1c).

Here we explored the costs and benefits of collective foraging for nutrient balancing in groups with increasing levels of inter-individual variance in nutritional requirements. We have developed a nutritional geometry focused ABM in which we assigned each agent an individual value of the group retention parameter K_{soc} . Allowing K_{soc} to mutate across generations, we then used an evolutionary

algorithm to explore how ecological and nutritional aspects of the environment affect optimal levels of group retention. Among those factors explored, we tested how aspects of inter-individual variation in nutrient needs interact with the composition and number of foods available to govern the degree of K_{soc} that optimizes an individual's ability to meet its nutritional requirements.

2. Material and methods

2.1. Agent-based model overview

All models were coded and simulation experiments performed in the ABM programming environment Netlogo 5.1 [26]. Model data were analysed and plotted in the statistical programming environment R v. 3.1.2 [27]. Below, we describe our model using the overview, design and details format as is now widely adopted for ABMs [28–30]. All Netlogo code can be found in the electronic supplementary material, S2 ‘Netlogo Code’.

2.1.1. Purpose

The ABMs have been programmed to evaluate how within-group heterogeneity affects the efficacy of social retention as a mechanism to improve individual foraging in a complex nutritional environment. Previous models suggest that a relatively high level of social retention is optimal when all individuals have equal needs [15,16]. Here, we use an evolutionary algorithm to explore how this optimal level of social retention varies with inter-individual variation in nutrient requirements.

2.1.2. Entities, states, variables and scales

The ABM is made up of individuals and their environment. The environment is a two-dimensional Cartesian coordinate system representing the space available for two nutrients (i.e. [figure 1a](#); implemented as in Senior *et al.* [23]). Each individual's nutritional state (NS; all parameters and variables are summarized in [table 1](#)) is given by its (x, y) coordinates. An individual's fitness (F) is maximized when its NS reaches its intake target (IT; [figure 1a,b](#)). Individuals are given a fixed period to reach their IT before the next generation begins, and generations are non-overlapping. The group has a mean IT of (μ_x, μ_y) and we alter the distribution of individual ITs around this mean. An individual's NS moves as it eats. There are N_{food} foods in the environment, each defined by a nutritional rail: a radial projecting through the nutrient space at an angle corresponding to the nutrient balance (V) of that food ([figure 1a](#)). That is, increasing N_{food} increases the diversity of foods, but not their total abundance. Individuals can only eat one food at a time and thus their NS moves in parallel to the nutritional rail for the food consumed ([figure 1a](#)). After eating an amount of a food (φ), an individual may seek an alternative, based on its own nutritional requirements and the number of other individuals consuming that food. The importance ascribed to these two factors is governed by a social retention parameter, K_{soc} , which evolves. The higher the value of K_{soc} , the lower the probability an individual will leave a popular food. We assume that foods can be patchily distributed in space and that there is a travel time (T) associated with moving between foods; we have explored the effects of different values of this time-cost on the evolution of K_{soc} . The distribution of individual ITs is governed by σ_{IT} and B (B is a fixed global parameter which governs the degree of bimodality in ITs; [table 1](#), [figure 1c](#)). We also explored variation in ability to consume food, φ , although no effects were observed (see electronic supplementary material, S1 and [figure S1](#)). The model runs for 1000 generations with K_{soc} values randomly mutating at each generation, and F governing an individual's representation in the subsequent generation. Note that our aim is not to precisely mimic evolution by natural selection but to explore optimal levels of K_{soc} under different environments and levels of group heterogeneity.

2.1.3. Process overview and scheduling

On each iteration of the ABM, the following three processes occur (detailed below, and in [figure 2a](#)): (i) ‘Find Food’, where those individuals not located on a food find one; (ii) ‘Eat’, where individuals located on a food eat some of it (i.e. move their NS through the environment); and (iii) ‘Leave Food’, where individuals may decide to leave a food on which they are feeding to seek an alternative. A new generation begins after 500 iterations. Data on the evolution of K_{soc} are recorded after 1000 generations.

Table 1. All model parameters and variables, their notation, level of operation and values (s.d. = standard deviation).

variable/parameter	notation	level	description	value
nutritional state	NS	individual	an individual's nutritional state, as tracked by its (x, y) position in the nutrient space	variable (x, y)
fitness	F	individual	an individual's fitness, which is a function of the Euclidean distance between an individual's NS and its IT	variable (equation (2.5))
intake target	IT	individual	an individual's IT is the (x, y) coordinate in the nutrient space that maximizes F	variable (x, y)
group mean IT	(μ_x, μ_y)	global	the mean IT (i.e. requirements) of the group for the nutrients on x - and y -axes	(500, 500)
number of foods	N_{food}	global	the number of food types (i.e. food rails) present in environment	2, 3 or 4
nutrient balance	V	global	the amount of y -axis nutrient present within a food, relative to each part of the x -axis nutrient (e.g. carbohydrate to protein in figure 1a)	0.0625, 0.25, 1, 4 and 16
capacity to eat food	φ	individual	the maximum amount of food that an individual can eat on one time step	2; see [25]
social retention	K_{soc}	individual	the degree to which an individual takes into account the popularity of a food, when evaluating whether to seek an alternative	variable and evolvable
time-cost	T	global	the time-cost associated with finding an alternative food	0–4
standard deviation in IT	σ_{IT}	global	the s.d. in ITs (inter-individual)	0–75
bimodality	B	global	the degree of bimodality in ITs (see equations (2.1) and (2.2))	0–0.5
deviation from the mean IT	ε	individual	the amount that an individual's IT deviates from the group mean	random-normal with mean = 0, s.d. = σ_{IT}
distance to the IT	D	individual	the Euclidean distance between an individual's NS and their IT	variable
ideal food rail	α_{ideal}	individual	the angle in radians of the ideal food rail connecting an individual's NS with their IT	variable
food 'popularity'	P	individual	the proportion of the group located on the same food as a given individual	variable
assortative interactions	A_{int}	individual	a binary variable denoting whether an individual is able to discriminate between and interact with conspecifics on the basis of their nutritional needs (1), or not (0)	0 or 1




Figure 2. The model's implementation of nutritional geometry. (a) Overview of the flow of events in the agent-based model. The cycle of events is repeated 500 times, before the next generation. Generations are non-overlapping and individuals are processed randomly (see details in Material and methods). (b) A schematic of the model's implementation of nutritional geometry [14]. The individual's nutritional state (NS) is given as an open point, with its intake target (IT) given as a grey cross hair. The individual is eating a food with a food rail denoted by a solid black line. A hypothetical ideal food (given by the grey dashed line) would allow the individual to reach the IT. The amount of the actual food to eat (i.e. distance to move) to minimize the Euclidean distance between the individual's NS and the IT is found by multiplying D by the angular difference between the ideal food rail and the angle of the actual food rail (arctangent of $V/1$). All parameters and variables are given in table 1. Adapted from Senior *et al.* [23].

2.2. Design concepts

Basic principles. We are interested in understanding how inter-individual variance in nutrient requirements influences optimal social retention during foraging.

Emergence. We are interested in the emergence of social retention, as given by the evolution of the parameter K_{soc} . We observe how this parameter evolves under varying distributions of nutritional requirements and environments.

Adaptation. An individual cannot adapt, but K_{soc} does evolve across generations.

Objectives. All individuals aim to reach their IT, which is the point in nutritional space that maximizes fitness, before the end of the generation.

Sensing. Individuals can sense the proportion of the group that is feeding on the same food as themselves.

Stochasticity. Many events in the ABM occur via Bernoulli trials and certain individual traits vary by values drawn from random distributions. For example, random variation in ITs is drawn from a random-normal distribution (see below for details).

Collectives. Collections of individuals can form on foods, the size of which will be dependent on the current values of K_{soc} .

Observation. Mean K_{soc} values of the entire group are observed after 1000 generations, where each generation consists of 500 model iterations.

2.3. Details

2.3.1. Initialization

The model is initialized with 100 individuals with a NS of (0,0), and $K_{soc}=0$. Individuals set an individual IT as given by either equation (2.1) or (2.2) with equal probability:

$$IT(x, y) = (\mu_x + \varepsilon + \mu_x B, \mu_y + \varepsilon - \mu_y B) \quad (2.1)$$

and

$$IT(x, y) = (\mu_x + \varepsilon - \mu_x B, \mu_y + \varepsilon + \mu_y B), \quad (2.2)$$

where μ_x and μ_y are the group mean requirements of the nutrients given on the x - and y -axes, B is a fixed global parameter taking values between 0 and 0.5 and governing the degree of bimodality in the ITs of all individuals (figure 1c) and ε is a value drawn from a random-normal distribution with a mean of 0 and standard deviation of σ_{IT} . Thus, as σ_{IT} increases, random variation is added to an individual's IT in both the x - and y -dimensions simultaneously, and as B increases individual ITs separate into two subgroups.

2.3.2. Find food

Those individuals that are not located on a food must find one. We assume a patchy environment, and thus every time an individual leaves a food they must spend T iterations searching. After T iterations without food an individual is randomly assigned a food from one of those available (all foods are assigned with equal probability). T may thus be thought of as the travel time between foods, or more conceptually how patchily foods are distributed. Given individuals have a fixed generation time to reach the IT, the inclusion of T thus constitutes a time-cost associated with moving between foods, which induces the selection pressure on K_{soc} . The order of magnitude of T is only relevant in the context of the amount of time individuals are given to reach the IT. However, the specific value of T will impact the strength of selection on K_{soc} . Previous models constrain T to a single value ($T = 2$ [16]), however, here we relax this constraint and explore a range of T values, including 0, which gives us an estimate of the evolution of K_{soc} in an environment where different foods are continuously distributed.

2.3.3. Eat

Eating constitutes an individual moving its NS through the nutrient space in parallel to the nutritional rail of the food consumed. The distance moved (i.e. amount eaten) is given by:

$$\text{distance moved} = \min\{\varphi, D \cos |\alpha_{ideal} - \arctan V|\}, \quad (2.3)$$

where φ is the maximum amount of food that an individual is capable of eating on one iteration, D is the Euclidean distance between the individual's IT and their NS, α_{ideal} is the angle (in radians) of a hypothetical ideal food rail connecting the individual's NS with their IT and the arctangent of V is the angle of the food rail of the food being consumed (a graphical representation of this model is given in figure 2b). Accordingly, individuals follow the 'closest distance rule of compromise' as defined by Simpson & Raubenheimer [14], whereby when possible individuals consume a food to minimize the Euclidean distance between their NS and IT (also see [16,23]).

2.3.4. Leave food

An individual's decision to leave a food is a probabilistic function of its own nutritional requirements (i.e. whether the food will allow the individual to reach its IT), and the choice of the rest of the population. The balance between these two factors is controlled by an individual's K_{soc} via:

$$\text{probability of leaving} = \max\{0.05, (1 - K_{soc})|\alpha_{ideal} - \arctan V| + K_{soc}e^{-7P}\}, \quad (2.4)$$

where e is the base of the natural logarithm, P is the proportion of the group currently consuming the same food as the individual (or the food's 'popularity') and all other parameters are as above. Accordingly, individuals with high K_{soc} are highly influenced by the behaviour of the rest of the group (P), and individuals with lower K_{soc} values pay more attention to whether the food will meet their own nutritional needs ($|\alpha_{ideal} - \arctan V|$). We also assume that all individuals have an innate minimum probability of leaving a food of 0.05 (representing the likelihood an individual may make an imperfect decision; following [16]).

2.3.5. Next generation

After 500 iterations of the above a new daughter generation begins. The fitness (F) of all individuals from the parental generation is calculated via

$$F = e^{-2D}, \quad (2.5)$$

where all variables are as above. Thus, as individuals near their IT, their F approaches 1. The size of the daughter generation is fixed at 100 individuals. Each individual inherits a value of K_{soc} from a parent of

the previous generation. At the point of inheritance (i.e. between generations), K_{soc} mutates by a value drawn at random from a normal distribution with a mean of 0 and standard deviation of 0.025 (and K_{soc} is bound at 0 and 1). All individuals in the parent group with $F \geq 0.25$ ($F < 0.25$ is not considered fit enough to reproduce) have a probability of being selected as the parent of an individual in the daughter group proportional to F (for details see Senior *et al.* [23]). Individuals within the daughter generation have an initial NS of (0, 0), and define their IT following ‘Initialization’ above. The parent generation is then replaced.

3. Results

We varied the individual heterogeneity variables, σ_{IT} and B (values given in table 1) and measured the group mean level of K_{soc} at the end of 1000 generations under each set of variables. For each parameter set we performed 30 model runs and report here the mean and 0.025–0.975 quantile of the results. We explored those effects in differing nutritional environments (i.e. different N_{food} and V ; table 1), and under differing assumptions about time-costs associated with finding foods (T).

3.1. Time-costs to foraging

First, we examined the efficiency of collective foraging in groups of individuals with the same IT (e.g. figure 1*a*). In an environment with two nutritionally imbalanced but complementary foods and in the absence of a time-cost to moving between foods ($T=0$), low levels of K_{soc} evolved and selection was weak (i.e. highly variable values of evolved K_{soc} ; figure 3*a*). However, with any time-cost to searching for foods ($T \geq 1$), K_{soc} evolved to be high and selection was strong (i.e. a narrow 0.025–0.975 quantile; figure 2*a*). At $T \geq 1$, mean $K_{\text{soc}} = 0.89$ (0.025–0.975 quantile = 0.86–0.92), which is equivalent to the optimal K_{soc} estimated by Lihoreau *et al.* [16] in their systematic exploration of the same parameter space. By way of contrast, with the inclusion of a third food that is nutritionally balanced relative to the group IT (rail passing through the IT), increasing T resulted in only slight increases in mean K_{soc} and in a very high variance (at $T=4$, mean $K_{\text{soc}} = 0.53$ and 0.025–0.975 quantile = 0.20–0.85; figure 3*b*).

To explore the mechanisms underlying the evolution of social retention, we re-ran the model with fixed K_{soc} values (0.3 or 0.9) and $T=4$ for a single generation. In both instances, we recorded the movements of individuals through the nutrient space and the time they spent foraging. In the two-food environment, low K_{soc} individuals moved frequently in order to regulate their intake of nutrients, but as a consequence spent a great deal of time foraging (figure 3*c,e*). By contrast, high K_{soc} individuals moved less frequently and took a wider path through the nutrient space (figure 3*c*). Consequently, high K_{soc} individuals spent roughly half as much time foraging as those with lower K_{soc} (figure 3*e*). In the three-food environment, all individuals spent little time foraging regardless of K_{soc} , because they quickly found the nutritionally balanced food and no longer searched for alternatives (figure 3*d,f*). Ultimately, in the absence of balanced foods containing an optimal nutrient mix for all group members, high levels of social retention were selected for by reducing individual search times for nutritionally complementary resources.

Further explorations of the model suggest that increasing the number of foods ($N_{\text{food}} = 4$) selects for lower K_{soc} . In these conditions, high K_{soc} forces individuals to move more often between foods, thereby precluding groups from forming on a given food, preventing foragers from efficiently tracking their IT. These effects are discussed in detail in the electronic supplementary material, S2 and figure S2.

3.2. Unimodal variation in individual nutrient requirements (σ_{IT})

Next, we examined the efficacy of collective foraging in heterogeneous groups, wherein each individual had its own IT. Here, all ITs were unimodally distributed around the group mean (e.g. figure 1*b*) and there was a high time-cost to moving between foods ($T=4$). In these conditions, no single food was balanced for all individuals. In a two-food environment, increasing variability in individual ITs (σ_{IT}) had no effect on evolved levels of K_{soc} (figure 4*a*). However, the inclusion of a third food, which contained an ideal mix of nutrients for the mean requirements of the group (rail passing through the mean group IT), dramatically altered these effects. Counterintuitively, increasing σ_{IT} selected for increases in K_{soc} (figure 3*b*). For example, in this three-food environment with $\sigma_{\text{IT}}=0$, mean $K_{\text{soc}} = 0.54$ (0.025–0.975 quantile = 0.25–0.81), but with $\sigma_{\text{IT}}=75$, mean $K_{\text{soc}} = 0.84$ (0.025–0.975 quantile = 0.52–0.93; figure 4*b*).

To understand the mechanisms underlying these effects, we re-ran the model and recorded the movements of individuals through the nutrient space with fixed high or low K_{soc} (0.3 or 0.9), $T=4$ and




Figure 3. Time-costs to foraging and the evolution of collective foraging. Mean and 0.025–0.975 quantile of K_{soc} after 1000 generations at differing levels of T based on 30 model runs in (a) a two-food and (b) a three-food environment for homogeneous groups of individuals with a single IT. Embedded in each panel is a geometric depiction of the nutritional environment showing the modelled food rails (V). Traces of the movement of 100 agents through the nutrient space (grey lines), and their intake targets (grey cross hair) where $T = 4$ and $K_{\text{soc}} = 0.3$ (i) and $K_{\text{soc}} = 0.9$ (ii) in (c) a two-food environment and (d) a three-food environment. The mean (± 0.025 –0.975 quantile) proportion of that model run spent foraging (i.e. moving between foods), by those individuals in (e) a two-food environment and (f) a three-food environment. $\sigma_{\text{IT}} = 0$ in all instances (see table 1 for all parameters and variables).

$\sigma_{\text{IT}} = 75$ for a single generation. In the two-food environment, the benefits of group retention were high, despite a large degree of variability in individual ITs. In these conditions, high K_{soc} individuals spent less time foraging than low K_{soc} individuals (figure 4c,e). Thus, the benefits of reducing time spent foraging outweighed any benefits associated with closely tracking an individual IT. In the three-food environment, the presence of a food passing through the mean group IT (and thus being close to the requirements of most individuals) caused variability in ITs to select for group retention. Where K_{soc} is low, high variance in individual ITs caused individuals to leave the food that passed through the mean group IT in order to closely track their own ITs, spending a great deal of time foraging (cf. figure 3d,f, where $\sigma_{\text{IT}} = 0$ and $K_{\text{soc}} = 0.3$, with figure 4d,f, where $\sigma_{\text{IT}} = 75$ and $K_{\text{soc}} = 0.3$). By contrast, high K_{soc} (0.9) drew individuals to the food with an equal ratio of nutrients, which had a nutrient balance that was relatively close to each individual's IT (although not meeting it exactly; figure 4d). Being attracted to this relatively balanced food reduced the amount of time an individual spent foraging, allowing individuals to get closer to their own IT than if they moved between foods frequently (figure 4d,f). Therefore, despite high inter-individual variation in nutritional needs, social retention still enhances the ability of individuals to balance their diet from multiple complementary foods in this case.

3.3. Bimodality in nutrient requirements

As well as unimodal variance in nutritional requirements, we also explored a more discrete form of heterogeneity by generating groups with bimodal distributions of ITs (e.g. figure 1c). Overall, we detected an interaction between the degree of bimodality (B) in ITs and the time-cost associated with locating foods (T). As one would predict, in a two-food environment, increasing B decreased the mean level of




Figure 4. Unimodal variance in individual nutrient requirements and the evolution of collective foraging. Mean and 0.025–0.975 quantile of K_{soc} after 1000 generations at differing levels of σ_{IT} and $T = 4$ based on 30 model runs in (a) a two-food and (b) a three-food environment. Embedded in each panel is a geometric depiction of the nutritional environment showing the modelled food rails (V), and a visualization of variation in individual ITs at certain values (σ_{IT} of 0, 37.5 and 75 are shown). Traces of the movement of 100 agents through the nutrient space (grey lines) and their intake targets (black points), where $T = 4$, $\sigma_{\text{IT}} = 75$ and $K_{\text{soc}} = 0.3$ (i) and $K_{\text{soc}} = 0.9$ (ii) in (c) a two-food environment and (d) a three-food environment. The mean (± 0.025 –0.975 quantile) proportion of that model run spent foraging (i.e. moving between foods), by those individuals in (e) a two-food environment and (f) a three-food environment (see table 1 for all parameters and variables).

evolved K_{soc} . However, decreases in K_{soc} were only observed at higher levels of B (e.g. $B > 0.25$), and with low levels of B mean evolved K_{soc} remained high (figure 4). The degree to which K_{soc} decreased at high B was dependent on T . At low to moderate T (i.e. $T = 1$ –3), high B only slightly reduced the evolved level of K_{soc} (mean $K_{\text{soc}} \approx 0.7$; figure 5). However, where both T and B were high, evolved K_{soc} was low (e.g. $T = 4$ and $B = 0.5$, mean $K_{\text{soc}} \approx 0.3$; figure 5).

The presence of a third food rail that meets the mean group IT (but of none of the individual ITs) produced different results (shown in the electronic supplementary material, figure S3). In general, in highly bimodal groups, collective foraging is not an efficient strategy, especially in environments containing foods with balances of nutrients that meet the mean requirements of the group.

3.4. Assortative interactions

A simple mechanism to overcome the constraints that discrete distributions in ITs place on the evolution of social retention may be *assortative interactions*, whereby group members only forage with those that share their nutritional needs. To explore this, we included a trait that allows individuals to discriminate between conspecifics on the basis of their ITs (A_{int} ; table 1), which may coevolve with K_{soc} . We assume that A_{int} is a binary and heritable trait, where $A_{\text{int}} = 1$ allows individuals to interact only with those that share their needs, and $A_{\text{int}} = 0$ does not (the probability of mutating from one state to the other was 0.01). Given that there may be time-costs associated with the processing of information for evaluating




Figure 5. Bimodality in nutrient requirements in a two-food environment and the evolution of collective foraging. Heat maps representing the effects of covarying B and T on the mean level of K_{soc} after 1000 generations, in a two-food environment (0.025–0.975 quantile given on the right). Shown on the left is a nutritional geometry depiction of model settings, including the nutritional value of food rails (V), the mean intake target (grey cross hair) and individual intake targets (black points), distributed around the group mean with a given B (see table 1 for all parameters and variables).

individuals prior to making a foraging decision, we also assume that individuals with $A_{\text{int}} = 1$ take an additional iteration to locate a food source (i.e. the time-cost of moving between foods = $T + A_{\text{int}}$).

We found that where A_{int} could evolve, the constraints that B had on the evolution of K_{soc} were overcome. For example, in a two-food environment with high T and high B (4 and 0.5, respectively), and where A_{int} and K_{soc} can coevolve, K_{soc} evolved to around 0.9 (figure 6a). This result contrasts with the evolved K_{soc} of 0.2 obtained under equivalent model setting where A_{int} could not evolve (figure 4). We also found selection for A_{int} , despite the associated cost. Where B and T are high, A_{int} evolved to be around 0.9 on average, but was selected against in all other areas of the parameter space (figure 6b). Thus, time-costs associated with foraging as well as distinct bimodality in nutrient requirements may select for the evolution of assortative foraging, whereby groups segregate into subgroups of individuals with similar nutritional needs. See the electronic supplementary material, S3, figures S4 and S5, for equivalent results in three- and four-food environments.

4. Discussion

Many animals, from insects to mammals, make collective decisions that enhance the speed and/or accuracy of individual choices [5,31,32]. While previous studies show how collective foraging enables individuals to select the richest or the largest available food resources in their environment [8–10], little is known about whether and how individuals can also collectively regulate their acquisition of vital nutrients from multiple imbalanced foods [15,16]. Our evolutionary model, based on concepts of nutritional geometry, indicates that collective foraging is often an efficient strategy for nutrient balancing, even in groups of individuals with different needs.

4.1. Collective foraging is adaptive in heterogeneous groups

In cases where nutrient intake must be balanced from complementary foods and there is a time-cost to foraging, our model shows that social retention is an effective mechanism for preventing individuals from over-investing in the time spent foraging. This prediction remained true in a number of instances where the specific requirements of individuals actually varied, only appearing to be suboptimal when the group contained an incredibly high degree of bimodality (i.e. figure 5). However, the presence of a food that met the mean requirements of the group (i.e. a food rail passing through the mean group IT) interacted with the distribution of ITs to determine optimal levels of social retention. When such a food was present and intra-group variation in ITs was low, social retention was only weakly




Figure 6. Bimodality in nutritional requirements and the co-evolution of collective foraging and assortative interactions. Heat maps represent the effects of covarying B and T on the mean (0.025–0.975 quantile given on the right) level of (a) K_{soc} and (b) A_{int} after 1000 generations, when the two traits are allowed to co-evolve in a two-food environment (i.e. equivalent of figure 4). Shown on the left is a nutritional geometry depiction of model settings, including the nutritional value of food rails (V), the mean intake target (grey cross hair) and individual intake targets (black points), distributed around the group mean with a given B (see table 1 for all parameters and variables).

selected for, despite costs associated with foraging. Surprisingly, in this environment, increasing inter-individual variance around a single mean (and therefore increasing potential conflict of interest between individuals) selected for stronger social retention (i.e. figure 4b). Here, collective foraging caused individuals to converge on a food that would meet their needs approximately, rather than over-investing (in terms of foraging time) in attempting to accurately cater to the idiosyncrasies of their own dietary requirements. Ecological scenarios such as those identified by our model may help explain how group retention can evolve to be so strong that it effectively traps individuals on nutritionally imbalanced foods, overriding their own nutritional needs, such as has been observed in the forest-tent caterpillar [13], where experimental conditions do not necessarily match natural nutritional environments.

Given the above, we can make a number of predictions about where organisms might, or might not, be expected to display collective foraging based on simple mechanisms of group retention. Namely, our model states that social retention becomes maladaptive under three circumstances: (i) where there is no cost associated with nutrient balancing (e.g. foods are nutritionally balanced or not patchily distributed); (ii) when the environment becomes increasingly complex (large numbers of different types

of foods patchily distributed), and individuals are too sparsely scattered across patches to effectively aggregate; and (iii) when the group has a highly bimodal distribution of requirements. Accordingly, we may expect organisms that inhabit stable environments falling under one, or more, of the above categories not to display social retention, whereas in the reciprocal environment the trait should evolve. Testing such a prediction would necessitate cross-taxa comparative study. However, in organisms that live in fluctuating environments, we might expect individuals to update and adapt their use of social information in response to environmental change [2,33]. For such species, nutritional geometry provides an experimental framework with which to evaluate the use of social information for nutrient balancing. Foods with known nutritional composition can be created, allowing one to manipulate the composition and distribution of foods in the environment [14]. In addition, if the species does not intrinsically contain individuals with differing nutritional needs it may be possible to experimentally modify the distribution of inter-individual requirements by manipulating individual NS (e.g. feeding individuals with differing foods prior to the trial).

4.2. Assortative foraging and group heterogeneity

In the extreme case where inter-individual variance has a highly discrete (multimodal) distribution, for instance, when groups are composed of distinct subsets of individuals with divergent needs, social retention tended to be selected against. Bimodal distributions of ITs may be common in mixed groups composed of males and females. Such observations have been made, for example, in field crickets (*Teleogryllus commodus*) [25] and fruit flies (*Drosophila melanogaster*) [34,35], where females maximize their reproductive fitness on diets richer in protein than males. Our results suggest that where the differences between sexes in nutritional requirements are large, collective foraging may be selected against. However, the constraints that sex-specificity in nutritional requirements place on the evolution of group foraging are easily overcome if individuals interact in an assortative manner, for instance by preferentially following individuals of the same sex rather than the group as a whole.

Sexual segregation of foraging groups has been observed in taxa ranging from marine and terrestrial mammals to fish [36–38]. Numerous competing hypotheses have been proposed to explain this behaviour [38], and comparative analyses restricted to large herbivores have provided perhaps the best insights. A leading hypothesis, termed the ‘forage selection’, or ‘nutritional needs’ hypothesis, explicitly states that sexual segregation results from differential dietary requirements of the sexes [37]. While support from comparative studies for this hypothesis has been mixed [37,39], data have focused on looking for differences in ‘diet-quality’ (e.g. nitrogen content) between males and females, rather than differences in diet composition *per se* (i.e. required ratios and amounts of nutrients, or ITs). Our model suggests that bimodality in nutritional requirements, in combination with selective pressures that favour collective foraging (e.g. time-costs associated with foraging), can lead to the emergence of such sexual segregation.

4.3. Collective nutrient regulation and the evolution of group complexity

It has long been suggested that nutritional constraints, such as limited access to key nutrients, may promote the evolution of cooperation and division of labour in animal groups [40–42]. However, the lack of a conceptual framework for testing these hypotheses has long hampered such research. Our theoretical exploration of the costs and benefits of social retention across multiple levels of group heterogeneities suggests that nutritional factors such as spatial/temporal distribution of foods and individual nutrient requirements are potential drivers of the evolution of collective foraging. Recently, it has been proposed that levels of intra-group variance in ITs and NSs may become greater with increasing degrees of social complexities, as one moves from forming temporary aggregations to living in permanent and fully eusocial colonies that forage from a central sedentary location [15]. Group living may increase inter-individual variance in requirements because of the simultaneous effects of competition over nutrient acquisition, age structures or differential parental nourishment, all of which are associated with cooperation and division of labour [43–45]. Ultimately, these factors may lead to the evolution of highly integrated social groups containing multiple classes of individuals with discrete ITs, as for instance the different castes of individuals that characterize the colonies of eusocial insects (e.g. ants, bees, wasps, termites) [46]. Although not its primary aim, our model predicts an evolutionary relationship between group heterogeneity (as measured by inter-individual variance in ITs) and social interactions (collective foraging) that is modulated by the nutritional environment (nutrient balance of available foods), partially supporting the above hypothesis.

Our results further suggest that collective foraging is not selected for in groups composed of individuals with highly divergent nutritional needs (i.e. bimodal distribution of ITs). However, this does not necessarily mean that intra-group variance is incompatible with sociality. As we have shown, in such species, additional mechanisms can evolve to mitigate the costs of collective foraging. For instance, eusocial insects have evolved sophisticated collective regulatory behaviour whereby a subset of individuals (the foragers) attempts to collect amounts and balances of nutrients that address the divergent nutritional needs of all colony members. In these advanced societies, foragers are able to gather foods that meet the high protein requirements of larvae and the queens, as well as the other adults that are more reliant on carbohydrates, based on a complex system of nutritional feedback between colony members [47–51].

4.4. Further development and conclusions

Our model, derived from nutritional geometry, provides a theoretical platform for exploring the costs and benefits of collective foraging in complex nutritional environments. A major advantage of this approach is that it generates specific predictions that can be empirically tested using well-established experimental designs from nutritional geometry [18]. Using this framework, it is possible to combine behavioural observations of groups of interacting animals feeding on chemically defined diets and to correlate individual nutrient intakes with measures of fitness traits both at the individual and collective levels [18]. Our model assumes a perhaps simplistic nutritional environment (i.e. foods are not finite and do not shift spatially), as our focus is primarily on heterogeneity in individual needs, in an environment that can be experimentally replicated. Future developments of our multidimensional approach, however, could integrate spatio-temporal distributions of finite foods constituting different compositions of essential nutrients, which ultimately determine the probability of a forager locating resources [52,53]. These factors constitute perhaps the most realistic costs to foraging in complex and ecologically relevant environments. Temporal and spatial variability in food abundance will alter the value of social information, thus affecting the nutritional strategy or strategies that will be optimal, both for an individual and groups. What is more, finite food sources will intrinsically capture aspects of intra-group conflict over a limiting resource, which will further affect the cost–benefit trade-off of group living.

Previous models demonstrate that simple mechanisms of social retention can optimize nutrient balancing in groups with homogeneous nutritional requirements. Here we present the first evolutionary ABM that re-evaluates the efficacy of such mechanisms for groups with heterogeneous nutritional needs. Our model suggests that simple mechanisms of social retention can improve foraging efficiency by reducing foraging times, even in relatively heterogeneous groups. Further, counterintuitively in some nutritional environments, heterogeneity in nutritional needs may select for increased collective decision-making. However, in groups with highly bimodal distributions of nutritional requirements (e.g. where needs of the sexes differ greatly) additional mechanisms of assortative foraging must coevolve alongside mechanisms of group retention, a potential route to the evolution of sexual segregation. Finally, our model may imply an evolutionary relationship between nutritional requirements, the ecological costs associated with foraging and the evolution of sociality, and provides a powerful framework for investigating such relationships.

Authors' contributions. A.M.S., M.L. and S.J.S. conceived the study. A.M.S., M.L., J.B. and M.C. designed the model. All authors contributed to the interpretation of the findings and the preparation of the manuscript.

Competing interests. We declare we have no competing interests.

Funding. This work was supported by the Australian Research Council (ARC). S.J.S. is supported by an ARC Laureate Fellowship. J.B. was supported by ARC Future Fellowship (ARC FT110100082) and ARC Discovery Projects (ARC DP1030101670) programs. M.L. is supported by grants from the IDEX of the Federal University of Toulouse, the Fyssen Foundation and the CNRS. D.R. is an affiliate of the New Zealand Institute for Advanced Study.

References

1. Danchin É, Giraldeau L-A, Valone TJ, Wagner RH. 2004 Public information: from nosy neighbors to cultural evolution. *Science* **305**, 487–491. (doi:10.1126/science.1098254)
2. Grüter C, Leadbeater E. 2014 Insights from insects about adaptive social information use. *Trends Ecol. Evol.* **29**, 177–184. (doi:10.1016/j.tree.2014.01.004)
3. Conradt L, Roper TJ. 2005 Consensus decision making in animals. *Trends Ecol. Evol.* **20**, 449–456. (doi:10.1016/j.tree.2005.05.008)
4. Ward AJW, Sumpter DJT, Couzin ID, Hart PJB, Krause J. 2008 Quorum decision-making facilitates information transfer in fish shoals. *Proc. Natl Acad. Sci. USA* **105**, 6948–6953. (doi:10.1073/pnas.0710344105)
5. Couzin ID. 2009 Collective cognition in animal groups. *Trends Cogn. Sci.* **13**, 36–43. (doi:10.1016/j.tics.2008.10.002)
6. Sumpter DJT, Pratt SC. 2009 Quorum responses and consensus decision making. *Phil. Trans. R. Soc.*

- B* **364**, 743–753. (doi:10.1098/rstb.2008.0204)
7. Gordon DM. 2014 The ecology of collective behavior. *PLoS Biol.* **12**, e1001805. (doi:10.1371/journal.pbio.1001805)
 8. Seeley TD, Camazine S, Sneyd J. 1991 Collective decision making in honey bees: how colonies choose among nectar sources. *Behav. Ecol. Sociobiol.* **28**, 277–290. (doi:10.1007/BF00175101)
 9. Beckers R, Deneubourg JL, Goss S, Pasteels JM. 1990 Collective decision making through food recruitment. *Insectes Soc.* **37**, 258–267. (doi:10.1007/BF02224053)
 10. Deneubourg JL, Goss S. 1989 Collective patterns and decision-making. *Ethol. Ecol. Evol.* **1**, 295–311. (doi:10.1080/08927014.1989.9525500)
 11. Lihoreau M, Deneubourg J-L, Rivault C. 2010 Collective foraging decision in a gregarious insect. *Behav. Ecol. Sociobiol.* **64**, 1577–1587. (doi:10.1007/s00265-010-0971-7)
 12. Lihoreau M, Rivault C. 2011 Local enhancement promotes cockroach feeding aggregations. *PLoS ONE* **6**, e22048. (doi:10.1371/journal.pone.0022048)
 13. Dussutour A, Simpson SJ, Despland E, Colasurdo N. 2007 When the group denies individual nutritional wisdom. *Anim. Behav.* **74**, 931–939. (doi:10.1016/j.anbehav.2006.12.022)
 14. Simpson SJ, Raubenheimer D. 2012 *The nature of nutrition: a unifying framework from animal adaptations to human obesity*. Oxford, UK: Princeton University Press.
 15. Lihoreau M, Buhl J, Charleston MA, Sword GA, Raubenheimer D, Simpson SJ. 2015 Nutritional ecology beyond the individual: a conceptual framework for integrating nutrition and social interactions. *Ecol. Lett.* **18**, 273–286. (doi:10.1111/ele.12406)
 16. Lihoreau M, Buhl J, Charleston MA, Sword GA, Raubenheimer D, Simpson SJ. 2014 Modelling nutrition across organizational levels: from individuals to superorganisms. *J. Insect Physiol.* **69**, 2–11. (doi:10.1016/j.jinsphys.2014.03.004)
 17. Raubenheimer D, Simpson SJ. 1993 The geometry of compensatory feeding in the locust. *Anim. Behav.* **45**, 953–964. (doi:10.1006/anbe.1993.1114)
 18. Simpson SJ, Clissold F, Lihoreau M, Ponton F, Wilder SM, Raubenheimer D. 2015 Recent advances in the integrative nutrition of arthropods. *Annu. Rev. Entomol.* **60**, 293–311. (doi:10.1146/annurev-ento-010814-020917)
 19. Lee KP, Simpson SJ, Clissold FJ, Brooks R, Ballard JWO, Taylor PW, Soran N, Raubenheimer D. 2008 Lifespan and reproduction in *Drosophila*: new insights from nutritional geometry. *Proc. Natl Acad. Sci. USA* **105**, 2498–2503. (doi:10.1073/pnas.0710787105)
 20. Solon-Biet SM *et al.* 2015 Macronutrient balance, reproductive function and lifespan in aging mice. *Proc. Natl Acad. Sci. USA* **112**, 3481–3486. (doi:10.1073/pnas.1422041112)
 21. Simpson SJ, Sibly RM, Lee KP, Behmer ST, Raubenheimer D. 2004 Optimal foraging when regulating intake of multiple nutrients. *Anim. Behav.* **68**, 1299–1311. (doi:10.1016/j.anbehav.2004.03.003)
 22. Jensen K, Mayntz D, Toft S, Clissold FJ, Hunt J, Raubenheimer D, Simpson SJ. 2012 Optimal foraging for specific nutrients in predatory beetles. *Proc. R. Soc. B* **279**, 2212–2218. (doi:10.1098/rspb.2011.2410)
 23. Senior AM, Charleston MA, Lihoreau M, Buhl J, Raubenheimer D, Simpson SJ. 2015 Evolving nutritional strategies in the presence of competition: a geometric agent-based model. *PLoS Comput. Biol.* **11**, e1004111. (doi:10.1371/journal.pcbi.1004111)
 24. Senior AM, Nakagawa S, Lihoreau M, Simpson SJ, Raubenheimer D. 2015 An overlooked consequence of dietary mixing: a varied diet reduces inter-individual variance in fitness. *Am. Nat.* **186**, 649–659. (doi:10.1086/683182)
 25. Maklakov AA, Simpson SJ, Zajitschek F, Hall MD, Dessmann J, Clissold F, Raubenheimer D, Bonduriansky R, Brooks RC. 2008 Sex-specific fitness effects of nutrient intake on reproduction and lifespan. *Curr. Biol.* **18**, 1062–1066. (doi:10.1016/j.cub.2008.06.059)
 26. Wilensky U. 1999 NetLogo. Center for Connected Learning and Computer-Based Modeling, Northwestern University Evanston, IL. See <http://ccl.northwestern.edu/netlogo/>.
 27. R Development Core Team. 2015 *R: a language and environment for statistical computing*. Version 3.2.1. Vienna, Austria: R Foundation for Statistical Computing.
 28. Grimm V, *et al.* 2006 A standard protocol for describing individual-based and agent-based models. *Ecol. Model.* **198**, 115–126. (doi:10.1016/j.ecolmodel.2006.04.023)
 29. Grimm V, Berger U, DeAngelis DL, Polhill JG, Giske J, Railsback SF. 2010 The ODD protocol: a review and first update. *Ecol. Model.* **221**, 2760–2768. (doi:10.1016/j.ecolmodel.2010.08.019)
 30. Railsback SF, Grimm V. 2012 *Agent-based and individual-based modeling: a practical introduction*. Princeton, NJ: Princeton University Press.
 31. Krause J, Ruxton GD, Krause S. 2010 Swarm intelligence in animals and humans. *Trends Ecol. Evol.* **25**, 28–34. (doi:10.1016/j.tree.2009.06.016)
 32. Sumpter DJT. 2010 *Collective animal behaviour*. Princeton, NJ: Princeton University Press.
 33. Rieucau G, Giraldeau L-A. 2011 Exploring the costs and benefits of social information use: an appraisal of current experimental evidence. *Phil. Trans. R. Soc. B* **366**, 949–957. (doi:10.1098/rstb.2010.0325)
 34. Reddiex AJ, Gosden TP, Bonduriansky R, Chenoweth SF. 2013 Sex-specific fitness consequences of nutrient intake and the evolvability of diet preference. *Am. Nat.* **182**, 91–102. (doi:10.1086/670649)
 35. Lee KP, Kim J-S, Min K-J. 2013 Sexual dimorphism in nutrient intake and life span is mediated by mating in *Drosophila melanogaster*. *Anim. Behav.* **86**, 987–992. (doi:10.1016/j.anbehav.2013.08.018)
 36. Leung ES, Chilvers BL, Nakagawa S, Moore AB, Robertson BC. 2012 Sexual segregation in juvenile New Zealand sea lion foraging ranges: implications for intraspecific competition, population dynamics and conservation. *PLoS ONE* **7**, e45389. (doi:10.1371/journal.pone.0045389)
 37. Main MB. 2008 Reconciling competing ecological explanations for sexual segregation in ungulates. *Ecology* **89**, 693–704. (doi:10.1890/07-0645.1)
 38. Wearmouth VJ, Sims DW. 2008 Sexual segregation in marine fish, reptiles, birds and mammals: behaviour patterns, mechanisms and conservation implications. In *Advances in marine biology* (ed. WS David), pp. 107–170. New York, NY: Academic Press.
 39. Mysterud A. 2000 The relationship between ecological segregation and sexual body size dimorphism in large herbivores. *Oecologia* **124**, 40–54. (doi:10.1007/s004420050023)
 40. Hunt JH, Kensing BJ, Kossuth JA, Henshaw MT, Norberg K, Wolschin F, Amdam GV. 2007 A diapause pathway underlies the gyne phenotype in *Polistes* wasps, revealing an evolutionary route to caste-containing insect societies. *Proc. Natl Acad. Sci. USA* **104**, 14 020–14 025. (doi:10.1073/pnas.0705660104)
 41. Amdam GV, Page RE. 2010 The developmental genetics and physiology of honeybee societies. *Anim. Behav.* **79**, 973–980. (doi:10.1016/j.anbehav.2010.02.007)
 42. West-Eberhard MJ. 2003 *Developmental plasticity and evolution*. Oxford, UK: Oxford University Press.
 43. Eggert A-K, Otte T, Müller JK. 2008 Starving the competition: a proximate cause of reproductive skew in burying beetles (*Nicrophorus vespilloides*). *Proc. R. Soc. B* **275**, 2521–2528. (doi:10.1098/rspb.2008.0661)
 44. Salomon M, Mayntz D, Lubin Y. 2008 Colony nutrition skews reproduction in a social spider. *Behav. Ecol.* **19**, 605–611. (doi:10.1093/beheco/arm008)
 45. Schwander T, Lo N, Beekman M, Oldroyd BP, Keller L. 2010 Nature versus nurture in social insect caste differentiation. *Trends Ecol. Evol.* **25**, 275–282. (doi:10.1016/j.tree.2009.12.001)
 46. Hölldobler B, Wilson EO. 2008 *The superorganism: the beauty, elegance and strangeness of insect societies*, 1st edn. New York, NY: W. H. Norton & Company.
 47. Dussutour A, Simpson SJ. 2009 Communal nutrition in ants. *Curr. Biol.* **19**, 740–744. (doi:10.1016/j.cub.2009.03.015)
 48. Cook SC, Eubanks MD, Gold RE, Behmer ST. 2010 Colony-level macronutrient regulation in ants: mechanisms, hoarding and associated costs. *Anim. Behav.* **79**, 429–437. (doi:10.1016/j.anbehav.2009.11.022)
 49. Altaye SZ, Pirk CWW, Crewe RM, Nicolson SW. 2010 Convergence of carbohydrate-biased intake targets in caged worker honeybees fed different protein sources. *J. Exp. Biol.* **213**, 3311–3318. (doi:10.1242/jeb.046953)
 50. Stabler D, Paoli PP, Nicolson SW, Wright GA. 2015 Nutrient balancing of the adult worker bumblebee (*Bombus terrestris*) depends on its dietary source of essential amino acids. *J. Exp. Biol.* **218**, 793–802. (doi:10.1242/jeb.114249)
 51. Dussutour A, Simpson SJ. 2012 Ant workers die young and colonies collapse when fed a high-protein diet. *Proc. R. Soc. B* **279**, 2402–2408. (doi:10.1098/rspb.2012.0051)
 52. Reynolds AM, Sword GA, Simpson SJ, Reynolds DR. 2009 Predator percolation, insect outbreaks, and phase polyphenism. *Curr. Biol.* **19**, 20–24. (doi:10.1016/j.cub.2008.10.070)
 53. Bhattacharya K, Vicsek T. 2014 Collective foraging in heterogeneous landscapes. *J. R. Soc. Interface* **11**, 20140674. (doi:10.1098/rsif.2014.0674)