

PUBLISHED VERSION

M.K. Hyde, R.U. Newton, D.A Galvão, WA, R.A. Gardiner, S Occhipinti, A. Lowe, G.A. Wittert, SA, S.K. Chambers

Men's help-seeking in the first year after diagnosis of localised prostate cancer

European Journal of Cancer Care, 2017; 26(2):e12497-1-e12497-12


© 2016 The Authors. European Journal of Cancer Care Published by John Wiley & Sons Ltd. This is an open access article under the terms of the Creative Commons Attribution-NonCommercial License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes.

Originally published at:

<http://doi.org/10.1111/ecc.12497>

PERMISSIONS

<http://creativecommons.org/licenses/by-nc/4.0/>



Attribution-NonCommercial 4.0 International (CC BY-NC 4.0)



This is a human-readable summary of (and not a substitute for) the [license](#). [Disclaimer](#).

You are free to:

- Share** — copy and redistribute the material in any medium or format
- Adapt** — remix, transform, and build upon the material

The licensor cannot revoke these freedoms as long as you follow the license terms.

Under the following terms:

-  **Attribution** — You must give [appropriate credit](#), provide a link to the license, and [indicate if changes were made](#). You may do so in any reasonable manner, but not in any way that suggests the licensor endorses you or your use.
-  **NonCommercial** — You may not use the material for [commercial purposes](#).

No additional restrictions — You may not apply legal terms or [technological measures](#) that legally restrict others from doing anything the license permits.

10 August 2017

<http://hdl.handle.net/2440/105094>

Men's help-seeking in the first year after diagnosis of localised prostate cancer

M.K. HYDE, PHD, SENIOR RESEARCH FELLOW, *Menzies Health Institute Queensland, Griffith University, Gold Coast, Qld, and Cancer Council Queensland, Fortitude Valley, Qld*, R.U. NEWTON, PHD, PROFESSOR, *Exercise Medicine Research Institute, Edith Cowan University, Perth, WA, and University of Queensland Centre for Clinical Research, University of Queensland, Brisbane, Qld*, D.A. GALVÃO, PHD, PROFESSOR, *Exercise Medicine Research Institute, Edith Cowan University, Perth, WA*, R.A. GARDINER, MD, MBBS, FRACS, FRCS, PROFESSOR AND CONSULTING UROLOGIST, *Exercise Medicine Research Institute, Edith Cowan University, Perth, WA, University of Queensland Centre for Clinical Research, University of Queensland, Brisbane, Qld, and Department of Urology, Royal Brisbane and Women's Hospital, Brisbane, Qld*, S. OCCHIPINTI, PHD, SENIOR LECTURER, *Menzies Health Institute Queensland, Griffith University, Gold Coast, Qld*, A. LOWE, PHD, CHIEF EXECUTIVE OFFICER, *Menzies Health Institute Queensland, Griffith University, Gold Coast, Qld, and Prostate Cancer Foundation of Australia, St Leonards, NSW*, G.A. WITTERT, MD, FRACP, FRCP, PROFESSOR, *Freemasons Foundation Centre for Men's Health, School of Medicine, University of Adelaide, Adelaide, SA*, & S.K. CHAMBERS, RN, PHD, PROFESSOR, *Menzies Health Institute Queensland, Griffith University, Gold Coast, Qld, Cancer Council Queensland, Fortitude Valley, Qld, Exercise Medicine Research Institute, Edith Cowan University, Perth, WA, University of Queensland Centre for Clinical Research, University of Queensland, Brisbane, Qld, and Prostate Cancer Foundation of Australia, St Leonards, NSW, Australia*

HYDE M.K., NEWTON R.U., GALVÃO D.A., GARDINER R.A., OCCHIPINTI S., LOWE A., WITTERT G.A. & CHAMBERS S.K. (2016) *European Journal of Cancer Care* **26**, e12497, doi: 10.1111/ecc.12497

Men's help-seeking in the first year after diagnosis of localised prostate cancer

This study describes sources of support utilised by men with localised prostate cancer in the first year after diagnosis and examines characteristics associated with help-seeking for men with unmet needs. A cross-sectional survey of 331 patients from a population-based sample who were in the first year after diagnosis ($M = 9.6$, $SD = 1.9$) was conducted to assess sources of support, unmet supportive care needs, domain-specific quality of life and psychological distress. Overall, 82% of men reported unmet supportive care needs. The top five needs were sexuality (58%); prostate cancer-specific (57%); psychological (47%); physical and daily living (41%); and health system and information (31%). Professional support was most often sought from doctors (51%). Across most domains, men who were older ($P_s \leq 0.03$), less well educated ($P_s \leq 0.04$) and more depressed ($P_s \leq 0.05$) were less likely to seek help for unmet needs. Greater sexual help-seeking was related to better sexual function ($P = 0.03$), higher education ($P \leq 0.03$) and less depression ($P = 0.05$). Unmet supportive care needs are highly prevalent after localised prostate cancer diagnosis with older age, lower education and higher depression apparent barriers to help-seeking. Interventions that link across medicine, nursing and community based peer support may be an accessible approach to meeting these needs. Clinical Trial Registry: Trial Registration: ACTRN12611000392965.

Keywords: prostate cancer, supportive care needs, help-seeking, survivorship.

Correspondence address: Melissa Hyde, Menzies Health Institute Queensland, Griffith University, Gold Coast Campus, QLD 4222, Australia (e-mail: melissa.hyde@griffith.edu.au).

Accepted 10 March 2016

DOI: 10.1111/ecc.12497

European Journal of Cancer Care, 2017, **26**, e12497, DOI: 10.1111/ecc.12497

INTRODUCTION

Globally, over 1 million new cases of prostate cancer were diagnosed in 2012 with 68% of these occurring in more developed countries (Ferlay *et al.* 2014). Prostate cancer incidence is highest in Australia/New Zealand, North America, Western and Northern Europe (age

standardised rates per 100 000 range from 85.0 to 111.6), with incidence expected to increase globally to 1.7 million cases in 2030 (Center *et al.* 2012; Ferlay *et al.* 2014). Survival rates for prostate cancer have increased in most countries in the past two decades (Allemani *et al.* 2015) such that in the UK, North America and Australia/New Zealand approximately 90% of men now survive their prostate cancer at least 5 years and >80% survive for 10 years or more (Australian Institute of Health and Welfare 2013; Cancer Research UK 2014; American Cancer Society 2015). Although many more men are surviving prostate cancer, they are living with high and persistent symptom burdens often not addressed in follow-up care (Smith *et al.* 2000; Bernat *et al.* 2015; Carlsson *et al.* 2015; Gavin *et al.* 2015). In a recent UK study, approximately one-third of men up to 2 years post-prostate cancer diagnosis reported concerns with sexual, urinary and bowel function (Watson *et al.* 2015) and there is evidence to suggest that these effects can persist for a decade or more (Bernat *et al.* 2015; Carlsson *et al.* 2015). As well, men who were younger or received multi-modal treatment including androgen deprivation therapy (ADT) experienced worse effects (Smith *et al.* 2007; Carlsson *et al.* 2015). In addition to substantial symptom burden, prostate cancer survivors' supportive care needs are frequently not met (Steginga *et al.* 2001; Lintz *et al.* 2003; Smith *et al.* 2007; Chambers *et al.* 2015a,b; Watson *et al.* 2015). Up to one half of men report ongoing unmet sexuality, psychological, and health system and information needs after prostate cancer treatment (Steginga *et al.* 2001; Lintz *et al.* 2003; Smith *et al.* 2007; Bernat *et al.* 2015; Watson *et al.* 2015).

Problematically, although prior studies suggest that men access health care services (Holden *et al.* 2005, 2006) at similar frequencies to women (Hourani *et al.* 2016), they often do not actively seek help (or receive treatment) for the full range of their concerns (Addis & Mahalik 2003; Shabsigh *et al.* 2004; Galdas *et al.* 2005; Steginga *et al.* 2008; Forsythe *et al.* 2013; Hyde *et al.* 2016; Yousaf *et al.* 2015) unless prompted to do so by a partner or a direct enquiry from a health professional (Holden *et al.* 2006). In particular, men with prostate cancer are less likely to discuss their social and emotional concerns with health care providers compared to women with breast cancer (Forsythe *et al.* 2013). Reasons for men's hesitancy to seek help may include under-reporting of emotional, physical or sexual concerns (Kunkel *et al.* 2000; Holden *et al.* 2006; Bernat *et al.* 2015; Yousaf *et al.* 2015); somatisation of mental health symptoms (Kockler & Heun 2002; Fiske *et al.*

2009); self-reliance and a desire to maintain a sense of normalcy (Gray *et al.* 2000; Yousaf *et al.* 2015); concerns about burdening others (Gray *et al.* 2000); discomfort or embarrassment (Yousaf *et al.* 2015); being less well educated or unaware (Dunn *et al.* 1999); and preference for self-management (Klafke *et al.* 2014). However, although recent studies have identified the symptom burden and supportive care needs of prostate cancer survivors and the factors that contribute to these (Smith *et al.* 2000; McDowell *et al.* 2010, 2011; Bernat *et al.* 2015; Gavin *et al.* 2015; Watson *et al.* 2015), patterns of help-seeking in this population to address unmet needs are less well described (Neese *et al.* 2003; Hyde *et al.* 2016). Rutten *et al.* (2005) propose that cancer patient's unmet needs and sources of support should be examined with specific reference to cancer survivorship stage (e.g., diagnosis, treatment, post-treatment) in order to be responsive to context. For prostate cancer survivors, unmet supportive care needs are highest close to the time of treatment (Smith *et al.* 2007; Harrison *et al.* 2009). Accordingly, we describe men's patterns of help-seeking in the first year after prostate cancer treatment, their unmet supportive care needs and from this examine factors related to help-seeking in the context of unmet need.

METHODS

Recruitment

Men who were diagnosed with localised prostate cancer in the state of Queensland, Australia were recruited between September 2011 and November 2012 via the Queensland Cancer Registry as part of a randomised controlled trial that is ongoing (Chambers *et al.* 2011; Galvão *et al.* 2015). Men were eligible for the trial if they had undergone/were undergoing prostate cancer treatment; could read and speak English; had no prior history of head injury, dementia or current psychiatric illness; had no concurrent cancer; and received clearance to participate from their physician. This study reports cross-sectional baseline data from a sub-group of men drawn from the larger population-based cohort who were in the first year after diagnosis and had received treatment. The study was approved by the Griffith University Human Research Ethics Committee and human research ethics committees of hospitals across Queensland. All participants provided written informed consent.

Measures

Baseline assessment occurred via computer-assisted telephone interview using previously validated and reliable

self-report measures (Chambers *et al.* 2011; Galvão *et al.* 2015).

Help-seeking and sources accessed

Help-seeking for prostate cancer-related concerns since diagnosis was assessed using a self-report yes/no measure. Type of help sought was measured with a prompted list on which men indicated all resources (e.g. Internet, brochures/books) and sources of support (e.g. doctor, nurse/other health professional, family/friends, support group) they had accessed for their concerns since diagnosis (McDowell *et al.*, 2011; Hyde *et al.* 2016). An open-ended response option was provided for men to indicate if they had used a resource or source of support that was not listed.

Unmet supportive care needs

The Supportive Care Needs Survey Short Form-34 (SCNS-SF34) measured men's need for help across psychological, health systems and information, patient care and support, physical and daily living, and sexuality domains (Boyes *et al.* 2009). A previously validated eight-item prostate cancer-specific module was also included to assess urinary, bowel, hormone and masculinity-related sexuality (e.g. feeling like you've lost part of your manhood) needs (Steginga *et al.* 2001). Items were rated 1 (no need/not applicable), 2 (need was satisfied), 3 (low need), 4 (moderate need) or 5 (high need).

Disease-specific quality of life

The domain summary scores from the Expanded UCLA Prostate Index Composite (EPIC) was applied to measure disease-specific quality of life for urinary ($\alpha = 0.85$), bowel ($\alpha = 0.87$), sexual ($\alpha = 0.89$) and hormonal ($\alpha = 0.69$) function (Litwin *et al.* 1998; Wei *et al.* 2000). Scores for each domain were transformed to a 0–100 scale with higher scores indicating better functioning.

Distress

The Brief Symptom Inventory-18 (BSI-18) including subscales of anxiety ($\alpha = 0.73$), depression ($\alpha = 0.86$) and somatisation ($\alpha = 0.67$) and a Global Severity Index (GSI) ($\alpha = 0.88$) assessed psychological distress (Derogatis & Lopez 2000). Men reported the degree of distress experienced for each symptom in the week prior to assessment, scored 0 (not at all) to 4 (extremely), with higher scores indicating greater distress. Raw

scores were converted to standardised *t*-scores to identify men with clinically significant symptoms (Chambers *et al.* 2014). Consistent with previous studies of cancer patients, caseness was identified using a cut-off *t*-score ≥ 57 on the BSI-18 GSI or on at least two of the BSI-18 subscales (Zabora *et al.* 2001; Chambers *et al.* 2014).

Statistical analysis

Descriptive statistics were calculated for socio-demographic and treatment characteristics, and to describe psychological distress, disease-specific quality of life, unmet supportive care needs, help-seeking and sources of help accessed. For men who expressed some level of need (scored ≥ 3) overall and in each SCNS-SF34 domain (except patient care and support because too few men reported an unmet need on this domain), logistic regression was used to examine associations between help-seeking and the following variables: age, education, months since diagnosis, hormone treatment; BSI-18 somatic, depression, anxiety subscales; and EPIC urinary, bowel and sexual domain summary scores. The hormone summary score was not included in the analysis due to the small number of men who received ADT. Logistic regressions were also conducted to examine associations between the variables specified above and supportive care needs (except patient care and support).

RESULTS

Recruitment and participant characteristics

Initially, 1899 patients were identified as potentially eligible for the trial and of these, 1770 doctors were contacted for permission to recruit their patients of whom 88.4% ($n = 1564$) gave consent to do so. Six hundred and seventy-nine of 1501 patients contacted agreed to participate, of which 463 met eligibility criteria, gave consent and completed baseline assessment (Galvão *et al.* 2015). Within this group, 331 patients were in the first year after diagnosis and had received treatment for localised prostate cancer. Analyses in this study are based on data from this sub-group of men.

Mean age of men was 64.5 years (median = 65.0; SD = 7.6). Most were in a relationship (86%), educated at university/college (25%) or trade/technical (38%) level, and just over half had an income \geq AUD \$60 000. Men were within the first 12 months of diagnosis (M = 9.6 months; median = 10.0; SD = 1.9). Men had received treatment approximately 6 months prior to the study (M = 6.4 months; median = 6.8; SD = 2.5), and

Table 1. Patient socio-demographic and treatment characteristics and EPIC domain summary scores ($N = 331$)

Characteristics	Mean (SD)/range or %
Age (years)	64.5 (7.6)/44–89
Education (highest level completed)	
University or college degree	25.4
Trade/technical certificate/diploma	38.4
Senior high school	10.0
Junior high school	19.9
Primary school	6.0
Did not complete primary school	0.3
Marital status	
Married	81.6
Defacto	3.9
Divorced/separated	4.5
Widowed	1.2
Single	8.8
Gross household income (AUD)	
< \$20 000	11.2
\$20 000 to \$39 999	23.0
\$40 000 to \$59 999	14.2
\$60 000 to \$79 999	12.7
\$80 000+	35.6
Don't know/unwilling to answer	3.3
Months since diagnosis	9.6 (1.9)/2.9–12.5
Months since treatment	6.4 (2.5)/0.1–11.3
Treatment received	
Radical prostatectomy	68.9
EBRT with ADT	13.0
EBRT without ADT	2.7
Brachytherapy with ADT	2.1
Brachytherapy without ADT	5.4
EBRT & Brachytherapy with ADT	1.5
ADT only	2.7
Other	3.6
EPIC domain summary scores	
Urinary	85.3 (15.6)
Bowel	93.1 (11.0)
Hormonal	80.6 (13.9)
Sexual	36.0 (22.6)

ADT, androgen deprivation therapy; AUD, Australian dollars; EBRT, external beam radiation therapy; EPIC, expanded prostate cancer incidence composite.

most (68%) were treated with radical prostatectomy (Table 1).

Disease-specific quality of life

Table 1 reports EPIC domain summary scores and of these men reported few difficulties in the areas of urinary and bowel function and hormonal effects. By contrast, sexual function scores were lower. Specifically, 41% of men considered their sexual function to be a moderate/big problem in the month prior to baseline assessment; with 14.2% and 4.5% reporting urinary and bowel dysfunction, respectively, to be a moderate/big problem in the same time frame. Sixty-four men received ADT and of these approximately 10% identified hot flushes,

depression and body weight as a moderate/big problem. Lack of energy was a moderate/big problem for 22% of these men.

Psychological distress

The standardised mean BSI GSI was 45.5 (SD = 8.5, range 36–72), with mean subscale scores of 47.4 for somatisation (SD = 6.8, range 42–81), 46.3 for depression (SD = 7.2, range 42–77) and 45.2 for anxiety (SD = 7.5, range 39–72). Forty-two men (12.7%) were identified as reaching high distress using the cut-off specified (t -score ≥ 57).

Unmet supportive care needs

Overall, 82% of men had some (low, moderate or high) level of need across any supportive care need domains. Of these, over half had concerns about sexuality (58%) and prostate cancer-specific needs (57%) that were not addressed. Approximately, half reported psychological (47%) and physical and daily living (41%) needs, and less than one-third had health system and information (31%) or patient care and support (17%) needs. Moderate–high need was expressed most frequently on sexuality (40%) and prostate cancer-specific (34%) domains (Table 2). The top 10 items for which men reported moderate to high unmet needs are described in Table 2. Of these, sexuality needs caused the most concern or discomfort with approximately one-third of men reporting they needed help with changes in sexual feelings and relationships and their sense of masculinity (feeling like a man). Physical needs, particularly not being able to do things as before, lack of energy/tiredness, and urinary incontinence were moderate–high concerns for $\geq 10\%$ of men. Men also expressed moderate–high need for help with their own (10%) or close others' (12%) psychological well-being, and their feelings of uncertainty about the future (9%). A moderate–high information need regarding being informed about things to do to get well was reported by 9% of men. Table 3 shows that men who experienced increased anxiety or somatic symptoms and were treated with ADT (vs. not treated) were more likely to express unmet supportive care needs across a range of need areas. Better outcomes on sexual, urinary and bowel domains and older age were associated with less unmet needs (Table 3). Education and months since diagnosis were not related to unmet needs.

Help-seeking and sources of support accessed

Overall, 94% of men reported that they had accessed resources and/or support for their prostate cancer-related

Table 2. Supportive care needs domains and items (SCNS SF-34) (*N* = 331)

Supportive care needs*†	Some need %	No need or need satisfied %	Low need %	Moderate-high need %
Physical and daily living need	40.5	59.5	18.1	22.4
Not being able to do the things you used to do	24.8	75.2	10.3	14.5‡
Lack of energy/tiredness	24.8	75.2	14.8	10.0‡
Feeling unwell a lot of the time	8.7	91.3	3.0	5.7
Work around the home	8.5	91.5	4.9	3.6
Pain	6.6	93.4	3.0	3.6
Psychological need	47.1	52.9	23.0	24.1
Concerns about the worries of those close to you	23.0	77.0	11.2	11.8‡
Fears about the cancer spreading	21.5	78.5	12.7	8.8
Uncertainty about the future	21.1	78.9	12.1	9.0‡
Feeling down or depressed	17.5	82.5	7.3	10.2‡
Anxiety	15.7	84.3	7.3	8.4
Feelings of sadness	15.4	84.6	7.8	7.6
Worry that the results of treatment are beyond your control	14.2	85.8	7.3	6.9
Learning to feel in control of your situation	12.4	87.6	6.9	5.5
Feelings about death or dying	9.1	90.9	5.8	3.3
Keeping a positive outlook	8.2	91.8	3.9	4.3
Health system and information need	30.8	69.2	10.3	20.5
Being informed about things you can do to help yourself get well	14.5	85.5	5.5	9.0‡
Having one member of staff with whom you can talk to about all aspects of your condition, treatment and follow-up	13.0	87.0	4.5	8.5
Having access to professional counselling (e.g. psychologist, social worker, counsellor, nurse specialist) if you, family or friends need it	10.9	89.1	4.5	6.4
Being given information (written, diagrams, drawings) about aspects of managing your illness and side effects at home	9.4	90.6	4.2	5.2
Being given explanations of those tests for which you would like explanations	9.1	90.9	3.6	5.5
Being adequately informed about cancer which is under control or diminishing	8.8	91.2	2.4	6.4
Being adequately informed about the benefits and side effects of treatments before you choose to have them	8.5	91.5	2.4	6.1
Being informed about your test results as soon as feasible	7.3	92.7	2.1	5.2
Being given written information about the important aspects of your care	5.8	94.2	1.6	4.2
Being treated like a person not just another case	5.7	94.3	1.8	3.9
Being treated in a hospital or clinic that is as physically pleasant as possible	3.6	96.4	2.1	1.5
Patient care and support need	16.7	83.3	6.7	10.0
Reassurance by medical staff that the way you feel is normal	8.5	91.5	4.3	4.2
More choice about which cancer specialists you see	6.4	93.6	1.8	4.6
More choice about which hospital you attend	6.1	93.9	2.1	4.0
Hospital staff attending promptly to your physical needs	5.4	94.6	2.7	2.7
Hospital staff acknowledging, and showing sensitivity to, your feelings and emotional needs	2.4	97.6	0.6	1.8
Sexuality need	58.0	42.0	17.8	40.2
Changes in sexual feelings	47.7	52.3	18.7	29.0‡
Changes in sexual relationships	41.4	58.6	13.0	28.4‡
To be given information about sexual relationships	19.9	80.1	9.0	10.9‡
Prostate cancer-specific need	57.4	42.6	23.0	34.4
Feeling like you've lost part of your manhood	38.1	61.9	16.0	22.1‡
Urinary incontinence	26.0	74.0	12.7	13.3‡
Problems with your bowel habits	11.5	88.5	4.8	6.7
Hot flushes§	6.9	93.1	3.1	3.8
Difficulties in passing urine	6.6	93.4	3.6	3.0

Table 2. *Continued*

Supportive care needs*†	Some need %	No need or need satisfied %	Low need %	Moderate–high need %
Feeling that what you say is not taken seriously by others	9.4	90.6	4.5	4.9
Feeling as if you are going through a change of life like women do‡	3.4	96.6	1.1	2.3
Feeling like you have lost the ability to be aggressive	2.4	97.6	1.2	1.2

*Level of need for help in the last month.

†Need was scored 1–2 = no need/need satisfied, 3 = some need, 4–5 = moderate–high need.

‡Top 10 moderate–high supportive care need.

§*n* = 262 (69 men receiving androgen deprivation therapy were excluded).

concerns. Excluding resources accessed, 79.5% reported seeking support while 20.5% did not access any form of support. Breaking this down further to examine seeking support from a health professional (doctor, nurse, counselling, or cancer helpline), 61% reported seeking support while 39% did not seek any support from health professional(s) for prostate cancer-related concerns. The most common resources accessed were brochures or books from the doctor (78%) and the Internet (53%). Sources of support most frequently accessed by men since their prostate cancer diagnosis were a doctor (51%) and family or friends (48%). Local nurse counsellors, cancer helpline, and counselling services and libraries were rarely accessed. While prostate cancer support group use was not high, it was substantially more than other types of counselling services (Table 4).

Associations between help-seeking and unmet supportive care needs

Associations between socio-demographic and treatment characteristics, disease-specific quality of life, psychological distress, and help-seeking for men who had some level of need overall and on each supportive care needs domain were examined and results of the logistic regressions are displayed in Table 5. Men who expressed *some level of need overall* (*n* = 272) were less likely to seek help if they were older (OR 0.94, CI 0.89–0.99), limited to high school (OR 0.25, CI 0.09–0.74) or primary school (OR 0.07, CI 0.02–0.28) level education and had increased depression symptoms (OR 0.83, CI 0.70–0.98). Being older (OR 0.88, CI 0.80–0.98) and those limited to primary school level education (OR 0.14, CI 0.02–0.90) were associated with a lower likelihood of seeking help for *physical and daily living needs*. Men who had *psychological or health system and information needs* were less likely to seek help if they had not progressed beyond primary school level education (OR_{Psychological} 0.09, CI 0.02–0.51; OR_{HealthSystInfo} 0.06, CI 0.01–0.64) and increased depression symptoms

(OR_{Psychological} 0.71, CI 0.57–0.87; OR_{HealthSystInfo} 0.77, CI 0.60–1.00). As well, there was a trend for older men with unmet health system and information needs to seek help less often (0.91, CI 0.83–1.00, *P* = 0.057). Men with unmet *sexuality needs* were less likely to seek help if they were educated at high school (OR 0.24, CI 0.07–0.90) or primary school (OR 0.10, CI 0.02–0.57) level, more time had passed since diagnosis (OR 0.72, CI 0.53–0.97), they experienced more depression symptoms (OR 0.81, CI 0.66–1.00), and had better bowel function (OR 0.93, CI 0.87–1.00, *P* = 0.056), with help-seeking also associated with better sexual function (OR 1.03, CI 1.00–1.06). Being older (OR 0.92, CI 0.86–0.99), educated not further than primary school level (OR 0.08, CI 0.02–0.42), and increased depression (OR 0.83, CI 0.68–1.00) were associated with less help-seeking for men with *prostate cancer-specific needs*.

DISCUSSION

The present study confirms previous research over the past decade showing a high prevalence of unmet supportive care needs in men with localised prostate cancer (Steginga *et al.* 2001; Lintz *et al.* 2003; Smith *et al.* 2007; Watson *et al.* 2015). Hence, despite the development of clinical practice guidelines for men with prostate cancer (National Health and Medical Research Council 2003; American Urological Association 2007; Australian Cancer Network Management of Metastatic Prostate Cancer Working Party 2010; Parker *et al.* 2015) and generic guidelines for psychosocial care in oncology (National Comprehensive Cancer Network 2002; National Breast Cancer Centre and National Cancer Control Initiative 2003; Holland *et al.* 2011), the pattern of need is unremitting and sexuality needs in particular appear intransigent. These findings have implications moving forward given the recent focus on prostate cancer survivorship guidelines (Skolarus *et al.* 2014; Resnick *et al.* 2015) and the observation that knowing what might assist men and actually

Table 3. Factors associated with unmet supportive care needs in men with prostate cancer (N = 331)

Supportive care needs	Any need		Physical and daily living need		Psychological need		Health system and info need		Sexuality need		Prostate cancer-specific need*	
	OR	(CI 95%)	OR	(CI 95%)	OR	(CI 95%)	OR	(CI 95%)	OR	(CI 95%)	OR	(CI 95%)
Age (years)	0.96	(0.91–1.01)	0.14	0.98 (0.94–1.02)	0.24	0.92 (0.89–0.96)	< 0.001	0.95 (0.91–0.99)	0.02	0.92 (0.88–0.96)	< 0.001	0.97 (0.93–1.01)
Education												
University/college	Ref.		Ref.		Ref.		Ref.		Ref.		Ref.	
Trade/technical	0.90	(0.42–1.96)	0.79	1.22 (0.64–2.34)	0.54	0.67 (0.36–1.27)	0.22	1.67 (0.83–3.36)	0.15	0.97 (0.51–1.86)	0.92	0.62 (0.32–1.22)
High school	1.38	(0.59–3.22)	0.46	1.30 (0.66–2.59)	0.45	1.20 (0.62–2.31)	0.59	1.52 (0.72–3.21)	0.27	1.08 (0.55–2.13)	0.83	1.41 (0.70–2.84)
Primary school	1.28	(0.40–4.17)	0.69	1.06 (0.33–3.40)	0.92	1.98 (0.62–6.29)	0.25	1.28 (0.39–4.17)	0.69	1.27 (0.38–4.27)	0.70	2.27 (0.59–8.69)
Months since dx	1.09	(0.92–1.30)	0.32	0.95 (0.83–1.09)	0.50	0.98 (0.86–1.12)	0.80	1.07 (0.93–1.24)	0.35	1.13 (0.98–1.30)	0.08	1.06 (0.92–1.23)
ADT												
Absence	Ref.		Ref.		Ref.		Ref.		Ref.		Ref.	
Presence	0.53	(0.21–1.35)	0.18	2.29 (1.13–4.65)	0.02	1.52 (0.75–3.11)	0.25	2.15 (1.05–4.39)	0.04	0.50 (0.25–1.01)	0.06	0.39 (0.18–0.86)
BSI-18												
Somatization	1.28	(0.96–1.71)	0.10	1.24 (1.05–1.48)	0.01	1.29 (1.08–1.54)	0.005	1.09 (0.93–1.28)	0.27	0.89 (0.76–1.04)	0.14	1.08 (0.89–1.30)
Depression	0.93	(0.72–1.20)	0.56	1.02 (0.88–1.19)	0.80	1.09 (0.93–1.28)	0.30	0.96 (0.83–1.11)	0.59	1.00 (0.84–1.18)	0.97	1.01 (0.84–1.21)
Anxiety	1.44 (1.00–2.08)		0.05	1.17 (0.97–1.40)	0.10	1.08 (0.89–1.30)	0.44	1.20 (1.01–1.43)	0.04	1.18 (0.95–1.45)	0.13	1.24 (0.98–1.56)
EPIC												
Urinary	0.98	(0.95–1.01)	0.17	0.98 (0.96–1.00)	0.02	1.00 (0.98–1.02)	0.68	0.97 (0.96–0.99)	0.006	1.00 (0.98–1.02)	0.81	0.95 (0.93–0.98)
Bowel	0.96	(0.90–1.03)	0.30	0.95 (0.92–0.99)	0.01	0.98 (0.94–1.01)	0.19	0.99 (0.96–1.02)	0.56	0.97 (0.93–1.00)	0.07	0.95 (0.91–1.00)
Sexual	0.97 (0.96–0.99)	< 0.001	< 0.001	1.00 (0.99–1.02)	0.61	0.99 (0.97–1.00)	0.01	0.98 (0.97–1.00)	0.007	0.96 (0.95–0.98)	< 0.001	< 0.001

Ref., Reference; ADT, androgen deprivation therapy; dx, diagnosis.

*The two items that could be confounded with side effects from androgen deprivation therapy were excluded from this scale for analysis (experiencing hot flushes, and going through a change of life like women do). Bolding in the table denotes a statistically significant result.

Table 4. Resources and sources of support accessed since diagnosis ($N = 331$)

Help accessed	%
Resources	
Brochures or books from doctor	77.6
Internet	52.6
Brochures or books from family/friends	7.3
Library	2.7
Sources of support	
Doctor	50.8
Family/friends	48.0
Nurse/Other health professional	17.8
Prostate cancer support group	7.9
Cancer Helpline	4.5
Counselling service	2.1
Local nurse counsellors	0.0

moving this into practice are different questions (Deane 2014).

In this regard, although men may access health care services and do so more frequently as they age, this may not equate to help-seeking for reproductive health concerns such as erectile dysfunction (Holden *et al.* 2005, 2006). In a population-based study of close to 6000 Australian men, 21% reported moderate to severe erectile dysfunction and of these 30% had spoken to a health professional about their condition, with this proportion decreasing by age (17% of men aged ≥ 70 years had discussed their concerns) (Holden *et al.* 2005, 2006). Prostate cancer survivors are also less likely to raise their concerns with health care providers (Forsythe *et al.* 2013), and in particular discuss sexual concerns much less frequently in follow-up care than concerns about urinary or bowel function (Watson *et al.* 2015). This creates a challenge for health professionals in terms of how to best identify men with unmet needs and provide accessible support. As well it indicates a need for health care professionals to initiate discussion with men about sexuality irrespective of whether sexual concerns are expressed (Holden *et al.* 2006; Forsythe *et al.* 2013). The top three sources of support accessed by men in the present study were the medical professional, nurse and to a lesser extent prostate cancer support groups. These three points of support may present as a triad that if well linked and resourced may have the greatest potential for making a difference in men's unmet needs after prostate cancer diagnosis and treatment. Family and friends as a preferred source of support may also help to promote men's awareness and encourage help-seeking from available health and supportive care services (Holden *et al.* 2006; Smith *et al.* 2006). As well, the high utilisation of the Internet points to this as a key resource that could be leveraged to improve access to self-management tools as well as professional support.

Men who were older, less well educated and more depressed were less likely to seek help despite having unmet needs for support. For age, this may relate to masculine values around stoicism and self-reliance that may be more strongly held by older men (Chambers *et al.* 2015a,b) and suggests the importance of continuing to understand the unique challenges faced by specific sub-groups of men. Level of education likely reflects health literacy that is especially problematic given low health literacy is also related to poorer health outcomes (DeWalt *et al.* 2004) and poorer use of health services (Berkman *et al.* 2011). For example, in a population-based study of Australian men, lower levels of functional health literacy were associated with obstructive sleep apnoea and comorbidities including sedentary lifestyle, depression and cardiovascular disease (Li *et al.* 2014). The relationship between higher depression and not seeking help speaks to the need for regular assessment of psychosocial needs (Forsythe *et al.* 2013) including systematic distress screening, rather than waiting for distressed men to self-present (Chambers *et al.* 2014). Finally, help-seeking was associated with better sexual function and this may reflect that men who sought help had better outcomes. Alternatively, it may suggest that men were more likely to seek support for sexuality needs if they had better sexual function. This latter interpretation is consistent with prior research suggesting severity of erectile dysfunction deters help-seeking such that sexuality interventions may require tailoring to sexual function (Hyde *et al.* 2016).

Strengths of this study include a large cohort of men drawn from a population-based cancer registry; the use of valid and reliable measures; and addressing the knowledge-gap regarding associations between supportive care needs and help-seeking. Limitations include the cross-sectional design which precludes inferences about causality, and retrospective self-report assessment of help-seeking. Longitudinal assessment of supportive care needs for men with localised and advanced prostate cancer and their patterns of help-seeking are a key focus for future research.

CONCLUSION

The diagnosis and treatment of prostate cancer is followed by well-described supportive care needs that are frequently unmet and this was apparent in the current study, particularly for sexuality needs. Few men accessed the range of health professional and community support options available and men with greatest need were least likely to seek support. Older age, lower education, and depression are risk factors for men not seeking help. A new approach to

Table 5. Factors associated with help-seeking for unmet supportive care needs in men with prostate cancer

Supportive care needs Factors	Any need (n = 272)		Physical and daily living need (n = 134)		Psychological need (n = 156)		Health system and info need (n = 102)		Sexuality need (n = 192)		Prostate cancer-specific need* (n = 190)	
	OR [CI 95%]	P	OR [CI 95%]	P	OR [CI 95%]	P	OR [CI 95%]	P	OR [CI 95%]	P	OR [CI 95%]	P
Age (years)	0.94 (0.89–0.99)	0.03	0.88 (0.80–0.98)	0.02	0.92 (0.85–1.01)	0.10	0.91 (0.83–1.00)	0.06	0.95 (0.88–1.03)	0.20	0.92 (0.86–0.99)	0.03
Education	Ref.		Ref.		Ref.		Ref.		Ref.		Ref.	
University/college	0.38 (0.13–1.14)	0.09	0.58 (0.11–2.93)	0.51	0.66 (0.15–2.89)	0.58	0.43 (0.07–2.69)	0.37	0.69 (0.17–2.80)	0.61	0.31 (0.08–1.27)	0.10
Trade/technical	0.25 (0.09–0.74)	0.01	0.67 (0.12–3.74)	0.65	0.68 (0.16–2.84)	0.59	0.28 (0.04–1.73)	0.17	0.24 (0.07–0.90)	0.03	0.32 (0.08–1.28)	0.11
High school	0.07 (0.02–0.28)	< 0.001	0.14 (0.02–0.90)	0.04	0.09 (0.02–0.51)	0.007	0.06 (0.01–0.64)	0.02	0.10 (0.02–0.57)	0.009	0.08 (0.02–0.42)	0.003
Primary school	0.92 (0.76–1.10)	0.36	0.99 (0.74–1.32)	0.95	0.98 (0.75–1.28)	0.88	0.74 (0.53–1.05)	0.09	0.72 (0.53–0.97)	0.03	0.96 (0.77–1.21)	0.75
Months since dx												
ADT	Ref.		Ref.		Ref.		Ref.		Ref.		Ref.	
Absence	0.82 (0.34–2.01)	0.67	0.85 (0.22–3.19)	0.80	0.49 (0.14–1.76)	0.27	0.98 (0.21–4.65)	0.98	0.42 (0.12–1.50)	0.18	0.92 (0.29–2.90)	0.88
Presence												
BSI-18												
Somatization	1.11 (0.90–1.36)	0.32	1.23 (0.89–1.70)	0.20	1.08 (0.83–1.42)	0.56	1.35 (0.92–1.97)	0.12	1.04 (0.75–1.44)	0.81	1.17 (0.89–1.54)	0.25
Depression	0.83 (0.70–0.98)	0.03	0.93 (0.74–1.17)	0.51	0.71 (0.57–0.87)	0.001	0.77 (0.60–1.00)	0.05	0.81 (0.66–1.00)	0.05	0.83 (0.68–1.00)	0.05
Anxiety	1.09 (0.86–1.40)	0.47	0.88 (0.64–1.22)	0.45	1.28 (0.91–1.80)	0.15	1.23 (0.86–1.78)	0.26	1.10 (0.81–1.50)	0.54	1.07 (0.81–1.41)	0.65
EPIC												
Urinary	1.00 (0.97–1.02)	0.82	1.02 (0.98–1.05)	0.35	1.00 (0.96–1.03)	0.81	1.01 (0.98–1.05)	0.47	0.99 (0.96–1.03)	0.72	1.00 (0.97–1.03)	0.93
Bowel	0.97 (0.93–1.02)	0.20	0.99 (0.94–1.04)	0.65	0.98 (0.92–1.03)	0.39	0.99 (0.94–1.06)	0.93	0.93 (0.87–1.00)	0.06	1.00 (0.95–1.04)	0.79
Sexual	1.00 (0.98–1.02)	0.94	1.00 (0.97–1.04)	0.93	0.98 (0.96–1.01)	0.26	1.00 (0.96–1.04)	0.97	1.03 (1.00–1.06)	0.03	1.01 (0.98–1.03)	0.53

Ref., Reference; ADT, androgen deprivation therapy; dx, diagnosis.

*The two items that could be confounded with side effects from androgen deprivation therapy were excluded from this scale for analysis (experiencing hot flushes, and going through a change of life like women do). Bolding in the table denotes a statistically significant result.

supportive care for men with prostate cancer seems warranted that links across medicine, nursing and community-based peer support. Future research is needed to establish the optimal mode for intervening to reduce the morbidity associated with this disease.

ACKNOWLEDGEMENTS

This project was funded by Cancer Australia and *beyond-blue* (ID APP1008320). SKC had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis. The sponsors did not participate in the design or conduct of the study; collection, management, analysis and interpretation of the data; or in the preparation, review or approval of the manuscript. DAG is funded by a Movember New Directions Development Award from Prostate Cancer

Foundation of Australia and Cancer Council Western Australia Research Fellowship. SKC is an Australian Research Council Future Fellow; and AG by a Cancer Institute NSW grant. We gratefully acknowledge the support of the Urological Society of Australia and New Zealand; of Mr Bill McHugh, Mr Spence Broughton and Mr Peter Dornan as consumer advisors; and of Ms Brigid Hanley and Ms Sylvia Burns as prostate cancer nurse advisors in the undertaking of this research. We also thank and acknowledge Ms Anna Stiller for research assistance.

CONFLICT OF INTEREST

The authors have no conflict of interest, including relevant financial interests, activities, relationships and affiliations to declare.

REFERENCES

- Addis M.E. & Mahalik J.R. (2003) Men, masculinity, and the contexts of help seeking. *American Psychologist* **58**, 5–14.
- Allemani C., Weir H., Carreira H., Harewood R., Spika D., Wang X., Bannon F., Ahn J., Johnson C., Bonaventure A., Marcos-Gragera R., Stiller C., Azevedo e Silva G., Chen W., Ogunbiyi O., Rachet B., Soeberg M., You H., Matsuda T., Bielska-Lasota M., Storm H., Tucker T. & Coleman M. (2015) Global surveillance of cancer survival 1995–2009: analysis of individual data for 25 676 887 patients from 279 population-based registries in 67 countries (CONCORD-2). *The Lancet* **385**, 977–1010.
- American Cancer Society (2015) Survival rates for prostate cancer. Available at: <http://www.cancer.org/cancer/prostatecancer/detailedguide/prostate-cancer-survival-rates> (accessed 6/11/2015).
- American Urological Association (2007) Guideline for the management of clinically localized prostate cancer: 2007 update. *The Journal of Urology* **177**, 2106–2131.
- Australian Cancer Network Management of Metastatic Prostate Cancer Working Party (2010) *Clinical Practice Guidelines for the Management of Locally Advanced and Metastatic Prostate Cancer*. Cancer Council Australia and Australian Cancer Network, Sydney, NSW, Australia.
- Australian Institute of Health and Welfare (AIHW) (2013) Prostate cancer in Australia. Cancer series no. 79. Available at: <http://www.aihw.gov.au/WorkArea/DownloadAsset.aspx?id=60129545133> (accessed 3/11/2015).
- Berkman N.D., Sheridan S.L., Donahue K.E., Halpern D.J. & Crotty K. (2011) Low health literacy and health outcomes: an updated systematic review. *Annals of Internal Medicine* **155**, 97–107.
- Bernat J., Wittman D., Hawley S., Hamstra D., Helfand A., Haggstrom D., Darwish-Yassine M. & Skolarus T. (2015) Symptom burden and information needs in prostate cancer survivors: a case for tailored long-term survivorship care. *BJU International* doi:10.1111/bju.13329.
- Boyes A., Girgis A. & Lecathelinais C. (2009) Brief assessment of adult cancer patients' perceived needs: development and validation of the 34-item Supportive Care Needs Survey (SCNS-SF34). *Journal of Evaluation in Clinical Practice* **14**, 602–606.
- Cancer Research UK (2014) Prostate cancer statistics: prostate cancer survival. Available at: <http://www.cancerresearchuk.org/health-professional/cancer-statistics/statistics-by-cancer-type/prostate-cancer#heading-Two> (accessed 3/11/2015).
- Carlsson S., Drevin L., Loeb S., Widmark A., Lissbrant I., Robinson D., Johansson E., Stattin P. & Fransson P. (2015) Population-based study of long-term functional outcomes after prostate cancer treatment. *BJU International* doi:10.1111/bju.13179.
- Center M., Jemal A., Lortet-Tieulent J., Ward E., Ferlay J., Brawley O. & Bray F. (2012) International variation in prostate cancer incidence and mortality rates. *European Urology* **61**, 1079–1092.
- Chambers S.K., Newton R., Girgis A., Nielsen L., Lepore S., Mihalopoulos C., Gardiner R., Galvão D. & Occhipinti S. (2011) Living with prostate cancer: randomised controlled trial of a multimodal supportive care intervention for men with prostate cancer. *BMC Cancer* **11**, 317.
- Chambers S.K., Zajdlewicz L., Youlden D.R., Holland J.C. & Dunn J. (2014) The validity of the distress thermometer in prostate cancer populations. *Psycho-Oncology* **23**, 195–203.
- Chambers S.K., Hyde M.K., Oliffe J.L., Zajdlewicz L., Lowe A., Wootten A. & Dunn J. (2015a) Measuring masculinity in the context of chronic disease. *Psychology of Men and Masculinity*, doi:10.1037/men0000018.
- Chambers S.K., Scuffham P.A., Baade P.D., Lowe A.P., Dunn J., Galvão D.A., Gordon L., Smith D.P., Sandoe D.F., Wootten A.C. & Spry N.A. (2015b) Advancing prostate cancer survivorship research in Australia. *Cancer Forum* **39**, 204–209.
- Deane L.A. (2014) Recommendations for prostate cancer survivorship care. *Journal of Men's Health* **11**, 108.
- Derogatis L.R. & Lopez M.C. (2000) *Brief Symptom Inventory 18: Administration, Scoring and Procedures Manual*. National Computer Systems, Minneapolis, MN, USA.
- DeWalt D.A., Berkman N.D., Sheridan S., Lohr K.N. & Pignone M.P. (2004) Literacy and health outcomes. *Journal of General Internal Medicine* **19**, 1228–1239.
- Dunn J., Steginga S.K., Occhipinti S., McCaffrey J. & Collins D.M. (1999) Men's preferences for sources of

- information about and support for cancer. *Journal of Cancer Education* **14**, 238–242.
- Ferlay J., Soerjomataram I., Ervik M., Dikshit R., Eser S., Mathers C., Rebelo M., Parkin D. M., Forman D. & Bray F. (2014) GLOBOCAN 2012 v1.1, Cancer Incidence and Mortality Worldwide: IARC CancerBase No. 11 [Internet]. International Agency for Research on Cancer, Lyon, France. Available at: <http://globocan.iarc.fr> (accessed 14/11/2015).
- Fiske A., Wetherell J.L. & Gatz M. (2009) Depression in older adults. *Annual Review of Clinical Psychology* **5**, 363–389.
- Forsythe L., Kent E., Weaver K., Buchanan N., Hawkins N., Rodriguez J., Ryerson A. & Rowland J. (2013) Receipt of psychosocial care among cancer survivors in the United States. *Journal of Clinical Oncology* **31**, 1961–1969.
- Galdas P.M., Cheater F. & Marshall P. (2005) Men and health help-seeking behaviour: literature review. *Journal of Advanced Nursing* **49**, 616–623.
- Galvão D., Newton R., Gardiner R., Girgis A., Lepore S., Stiller A., Mihalopolous C., Occhipinti S. & Chambers S. (2015) Compliance to exercise-oncology guidelines in prostate cancer survivors and associations with psychological distress, unmet supportive care needs, and quality of life. *Psycho-Oncology* **24**, 1241–1249.
- Gavin A.T., Drummond F.J., Donnelly C., O'Leary E., Sharp L. & Kinnear H.R. (2015) Patient reported “ever had” and “current” long term physical symptoms following prostate cancer treatments. *BJU International* **116**, 397–406.
- Gray R., Fitch M., Phillips C., Labrecque M. & Fergus K. (2000) Managing the impact of illness: the experiences of men with prostate cancer and their spouses. *Journal of Health Psychology* **5**, 531–548.
- Harrison J., Young J., Price M., Butow P. & Solomon M. (2009) What are the unmet supportive care needs of people with cancer? A systematic review. *Supportive Care in Cancer* **17**, 1117–1128.
- Holden C., McLachlan R., Pitts M., Cumming R., Wittert G., Agius P., Handelsman D. & de Kretser D. (2005) Men in Australia Telephone Survey (MATEs): a national survey of the reproductive health and concerns of middle-aged and older Australian men. *The Lancet* **366**, 218–224.
- Holden C.A.E.A., Jolley D., McLachlan R., Pitts M., Cumming R., Wittert G., Handelsman D. & De Kretser D. (2006) Men in Australia Telephone Survey (MATEs): predictors of men's help-seeking behaviour for reproductive health disorders. *Medical Journal of Australia* **185**, 418–422.
- Holland J., Watson M. & Dunn J. (2011) The IPOS New International Standard of Quality Cancer Care: integrating the psychosocial domain into routine care. *Psycho-Oncology* **20**, 677–680.
- Hourani L., Williams J., Bray R., Wilk J. & Hoge C. (2016) Gender differences in posttraumatic stress disorder and help seeking in the U.S. army. *Journal of Women's Health* **25**, 22–31. doi: 10.1089/jwh.2014.5078
- Hyde M.K., Zajdlewicz L., Wootten A., Nelson C., Lowe A., Dunn J. & Chambers S.K. (2016) Medical help-seeking for sexual concerns in prostate cancer survivors. *Sexual Medicine* **4**, e7–e17. doi: 10.1016/j.exsm.2015.12.004
- Klafke N., Elliott J., Olver I. & Wittert G. (2014) Australian men with cancer practice complementary therapies (CTs) as a coping strategy. *Psycho-Oncology* **23**, 1236–1242.
- Kockler M. & Heun R. (2002) Gender differences of depressive symptoms in depressed and nondepressed elderly persons. *International Journal of Geriatric Psychiatry* **17**, 65–72.
- Kunkel E., Bakker J., Myers R., Oyesanmi O. & Gomella L. (2000) Biopsychosocial aspects of prostate cancer. *Psychosomatics* **41**, 85–94.
- Li J.J., Appleton S.L., Wittert G.A., Vakulin A., McEvoy R.D., Antic N.A. & Adams R.J. (2014) The relationship between functional health literacy and obstructive sleep apnea and its related risk factors and comorbidities in a population cohort of men. *Sleep* **37**, 571–578.
- Lintz K., Moynihan C., Steginga S., Norman A., Eeles R., Huddart R., Dearnaley D. & Watson M. (2003) Prostate cancer patients' support and psychological care needs: survey from a non-surgical oncology clinic. *Psycho-Oncology* **12**, 769–783.
- Litwin M.S., Hays R.D., Fink A., Ganz P.A., Leake B. & Brook R.H. (1998) The UCLA Prostate Cancer Index: development, reliability, and validity of a health-related quality of life measure. *Medical Care* **36**, 1002–1012.
- McDowell M., Occhipinti S., Ferguson M. & Chambers S. (2011) Prospective predictors of psychosocial support service use after cancer. *Psycho-Oncology* **20**, 788–791.
- McDowell M.E., Occhipinti S., Ferguson M., Dunn J. & Chambers S.K. (2010) Predictors of change in unmet supportive care needs in cancer. *Psycho-Oncology* **19**, 508–516.
- National Breast Cancer Centre and National Cancer Control Initiative (2003) *Clinical Practice Guidelines for the Psychosocial Care of Adults with Cancer*. National Breast Cancer Centre, Camperdown, NSW, Australia
- National Comprehensive Cancer Network (2002) *Practice Guidelines in Oncology-v.1.2002: distress management* (Rep. No. Version 1). National Comprehensive Network, Fort Washington, PA.
- National Health and Medical Research Council (2003) *Clinical Practice Guidelines: Evidence-based Information and Recommendations for the Management of Localised Prostate Cancer*. Commonwealth of Australia, NHMRC, Canberra, Australia.
- Neese L., Schover L., Klein E., Zippe C. & Kupelian P. (2003) Finding help for sexual problems after prostate cancer treatment: a phone survey of men's and women's perspectives. *Psycho-Oncology* **12**, 463–473.
- Parker C., Gillissen S., Heidenreich A. & Horwich A. & on behalf of the ESMO Guidelines Committee (2015) Cancer of the prostate: ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up. *Annals of Oncology* **26** (Suppl. 5), v69–v77.
- Resnick M.J., Lacchetti C., Bergman J., Hauke R.J., Hoffman K.E., Kungel T.M., Morgans A.K. & Penson D.F. (2015) Prostate cancer survivorship care guideline: American Society of Clinical Oncology clinical practice guideline endorsement. *Journal of Clinical Oncology* **33**, 1078–1085.
- Rutten L., Arora N., Bakos A., Aziz N. & Rowland J. (2005) Information needs and sources of information among cancer patients: a systematic review of research (1980–2003). *Patient Education and Counseling* **57**, 250–261.
- Shabsigh R., Perelman M., Laumann E. & Lockhart D. (2004) Drivers and barriers to seeking treatment for erectile dysfunction: a comparison of six countries. *BJU International* **94**, 1055–1065.
- Skolarus T.A., Wolf A., Erb N.L., Brooks D.D., Rivers B.M., Underwood W., Salner A.L., Zelefsky M.J., Aragon-Ching J.B., Slovin S.F. & Wittmann D.A. (2014) American Cancer Society prostate cancer survivorship care guidelines. *CA: A Cancer Journal for Clinicians* **64**, 225–249.
- Smith D., Carvalhal G., Schneider K., Krygiel J., Yan Y. & Catalona W. (2000) Quality-of-life outcomes for men with prostate carcinoma detected by screening. *Cancer* **88**, 1454–1463.
- Smith J.A., Braunack-Mayer A. & Wittert G.A. (2006) What do we know about

- men's help-seeking and health service use? *Medical Journal of Australia* **184**, 81–83.
- Smith D., Supramaniam R., King M., Ward J., Berry M. & Armstrong B. (2007) Age, health, and education determine supportive care needs of men younger than 70 years with prostate cancer. *Journal of Clinical Oncology* **25**, 2560–2566.
- Steginga S.K., Occhipinti S., Dunn J., Gardiner R., Heathcote P. & Yaxley J. (2001) The supportive care needs of men with prostate cancer (2000). *Psycho-Oncology* **10**, 66–75.
- Steginga S.K., Campbell A., Ferguson M., Beeden A., Walls M., Cairns W. & Dunn J. (2008) Socio-demographic, psychosocial and attitudinal predictors of help seeking after cancer diagnosis. *Psycho-Oncology* **17**, 997–1005.
- Watson E., Shinkins B., Frith E., Neal D., Hamdy F., Walter F., Weller D., Wilkinson C., Faithfull S., Wolstenholme J., Sooriakumaran P., Kastner C., Campbell C., Neal R., Butcher H., Matthews M., Perera R. & Rose P. (2015) Symptoms, unmet needs, psychological well-being and health status in survivors of prostate cancer: implications for redesigning follow-up. *BJU International*. doi:10.1111/bju.13122.
- Wei J.T., Dunn R.L., Litwin M.S., Sandler H.M. & Sanda M.G. (2000) Development and validation of the expanded prostate cancer index composite (EPIC) for comprehensive assessment of health-related quality of life in men with prostate cancer. *Urology* **56**, 899–905.
- Yousaf O., Grunfeld E.A. & Hunter M.S. (2015) A systematic review of the factors associated with delays in medical and psychological help-seeking among men. *Health Psychology Review* **9**, 264–276.
- Zabora J., Brintzenhofesoc K., Jacobsen P., Curbow B., Piantadosi S., Hooker C., Owens A. & Derogatis L. (2001) A new psychosocial screening instrument for use with cancer patients. *Psychosomatics* **42**, 241–246.