



Activity of Bacteriophages in Removing Biofilms of *Pseudomonas aeruginosa* Isolates from Chronic Rhinosinusitis Patients

Stephanie A. Fong¹, Amanda Drilling¹, Sandra Morales², Marjolein E. Cornet³, Bradford A. Woodworth⁴, Wytske J. Fokkens³, Alkis J. Psaltis¹, Sarah Vreugde¹ and Peter-John Wormald^{1*}

¹ Department of Surgery—Otolaryngology, Head and Neck Surgery, The Queen Elizabeth Hospital, Woodville South, SA, Australia, ² AmpliPhi Australia, Brookvale, NSW, Australia, ³ Department of Otorhinolaryngology, Academic Medical Center, University of Amsterdam, Amsterdam, Netherlands, ⁴ Department of Otolaryngology-Head and Neck Surgery, University of Alabama at Birmingham, Birmingham, AL, United States

OPEN ACCESS

Edited by:

Dongsheng Zhou,
Beijing Institute of Microbiology and
Epidemiology, China

Reviewed by:

Eduard Torrents,
Institute for Bioengineering of
Catalonia, Spain
Wei Zhang,
Nanjing Agricultural University, China
Feng Han,
Ocean University of China, China

*Correspondence:

Peter-John Wormald
peterj.wormald@adelaide.edu.au

Received: 06 July 2017

Accepted: 08 September 2017

Published: 22 September 2017

Citation:

Fong SA, Drilling A, Morales S, Cornet ME, Woodworth BA, Fokkens WJ, Psaltis AJ, Vreugde S and Wormald P-J (2017) Activity of Bacteriophages in Removing Biofilms of *Pseudomonas aeruginosa* Isolates from Chronic Rhinosinusitis Patients. *Front. Cell. Infect. Microbiol.* 7:418. doi: 10.3389/fcimb.2017.00418

Introduction: *Pseudomonas aeruginosa* infections are prevalent amongst chronic rhinosinusitis (CRS) sufferers. Many *P. aeruginosa* strains form biofilms, leading to treatment failure. Lytic bacteriophages (phages) are viruses that infect, replicate within, and lyse bacteria, causing bacterial death.

Aim: To assess the activity of a phage cocktail in eradicating biofilms of *ex vivo* *P. aeruginosa* isolates from CRS patients.

Methods : *P. aeruginosa* isolates from CRS patients with and without cystic fibrosis (CF) across three continents were multi-locus sequence typed and tested for antibiotic resistance. Biofilms grown *in vitro* were treated with a cocktail of four phages (CT-PA). Biofilm biomass was measured after 24 and 48 h, using a crystal violet assay. Phage titrations were performed to confirm replication of the phages. A linear mixed effects model was applied to assess the effects of treatment, time, CF status, and multidrug resistance on the biomass of the biofilm.

Results: The isolates included 44 strain types. CT-PA treatment significantly reduced biofilm biomass at both 24 and 48 h post-treatment ($p < 0.0001$), regardless of CF status or antibiotic resistance. Biomass was decreased by a median of 76% at 48 h. Decrease in biofilm was accompanied by a rise in phage titres for all except one strain.

Conclusion: A single dose of phages is able to significantly reduce biofilms formed *in vitro* by a range of *P. aeruginosa* isolates from CRS patients. This represents an exciting potential and novel targeted treatment for *P. aeruginosa* biofilm infections and multidrug resistant bacteria.

Keywords: bacteriophage, *Pseudomonas aeruginosa*, chronic rhinosinusitis, cystic fibrosis, biofilm, multidrug resistant

INTRODUCTION

Chronic rhinosinusitis (CRS) is an inflammatory condition of the nose and paranasal sinuses, persisting for 12 weeks or longer. Bacterial biofilms have been implicated in recalcitrant CRS and increase the tolerance of bacteria to antibiotics through numerous mechanisms. These include metabolic heterogeneity of the bacteria within the biofilm, enzymatic deactivation, anionic charges due to extracellular DNA within the biofilm matrix, and changes in gene expression (de la Fuente-Nú-éz et al., 2013). Biofilms have been found on the sinonasal mucosa of up to 54% of CRS sufferers, compared to 8% of control patients (Chen et al., 2012). Furthermore, multiple studies have noted a higher prevalence of biofilms in patients who are undergoing revision surgery (Psaltis et al., 2007; Chen et al., 2012). In particular, the presence of biofilm-forming *Pseudomonas aeruginosa* strains has been associated with poor resolution of symptoms and signs of CRS following endoscopic sinus surgery (Bendouah et al., 2006).

P. aeruginosa has been identified in the sinuses of 9% of CRS patients, and is associated with poorer quality of life measured by disease severity scores, such as the Visual Analogue Scale (VAS) and Sinonasal Outcome Test-22 (SNOT-22) (Cleland et al., 2013, 2016). *P. aeruginosa* sinus infections also commonly afflict patients with cystic fibrosis (CF), with the species being identified in sinus cultures of up to 49% of CF patients with CRS (Rasmussen et al., 2012). Moreover, *P. aeruginosa* is intrinsically resistant to many classes of antibiotics, and acquired antibiotic resistance is increasing worldwide. New therapeutic strategies are therefore required to combat these difficult to treat bacterial infections in the context of CRS.

Bacteriophages are viruses that infect bacteria. Lytic bacteriophages are able to lyse their host bacterium, after replicating themselves within the host bacterium. Lysis of the host not only kills the bacterium, but also releases the progeny copies of the phage for re-infection of other bacteria (Guttman et al., 2004). Bacteriophages are species specific, and so can be used to target pathogenic bacteria, without disturbing non-harmful commensal bacteria (Hanlon, 2007). Bacteriophages are also able to penetrate bacterial biofilms (Vilas Boas et al., 2016).

These characteristics make bacteriophages an attractive non-antibiotic therapy for treating bacterial biofilms in CRS. This study aims to assess the activity of a cocktail (mixture) of four *P. aeruginosa* bacteriophages (CT-PA) in removing *ex vivo* biofilms formed by PA isolates from CRS patients, both with and without CF. The prevalence of antibiotic resistance in these clinical isolates was also assessed.

METHODS

Bacterial Strains and Growth Conditions

This study was approved by the Human Research Ethics Committee of The Queen Elizabeth Hospital, Adelaide, South Australia.

P. aeruginosa strains were isolated by an independent pathology laboratory (Adelaide Pathology Partners, Adelaide,

South Australia) from endoscopically-guided sinus swabs, from patients who met the European Position Paper on Rhinosinusitis and Nasal Polyps (EPOS) 2012 criteria for chronic rhinosinusitis (Fokkens et al., 2012). Clinical *P. aeruginosa* isolates of patients with CF were kindly donated by the Department of Otorhinolaryngology, Academic Medical Centre (Amsterdam, Netherlands) and *P. aeruginosa* sinus isolates from CRS patients with and without CF were kindly donated by the Department of Otolaryngology-Head and Neck Surgery, University of Alabama at Birmingham (Birmingham, AL). *P. aeruginosa* isolates were stored in 25% glycerol in nutrient broth at -80°C . *P. aeruginosa* laboratory reference strain ATCC 15692 (PAO1) was obtained from American Type Culture Collection (Manassas, VA, USA) as a control for phage sensitivity and biofilm assays. Isolates were plated from frozen glycerol stocks onto 1.5% nutrient agar, and broth cultures were grown in nutrient broth. Agar plates and broth cultures were incubated at 37°C .

Bacteriophage Cocktail

Stocks of 4 anti-*P. aeruginosa* bacteriophages (Pa 193, Pa 204, Pa 222, Pa 223), as well as heat-inactivated stocks, were supplied by AmpliPhi Biosciences (Brookvale, New South Wales, Australia). Pa 193 and Pa 204 are of the Myoviridae family, and Pa 222 and Pa 223 are of the Podoviridae family. All 4 phages have been characterized as strictly lytic by genome sequencing (unpublished data). Prior to each assay, the stock suspension of each bacteriophage was titrated against a selected *P. aeruginosa* bacterial strain using the soft agar overlay small drop assay, as described below. Equal concentrations of each bacteriophage were combined to form the bacteriophage cocktail (CT-PA).

Multi-Locus Sequence Typing

The protocol for multi-locus sequence typing (MLST) of *P. aeruginosa* isolates has been described by Curran et al (Curran et al., 2004). For each strain, a single colony was used to inoculate 5 mL of nutrient broth, and the culture was grown overnight on a shaker at 37°C . DNA extraction from a 1 mL aliquot of the overnight culture was performed using the DNeasy Blood and Tissue DNA extraction kit (Qiagen, Hilden, Germany), following the protocol for Gram negative bacteria recommended by the manufacturer. PCR of the 7 MLST loci was performed using the Taq PCR kit (New England Biolabs, Ipswich MA, USA) using the protocol and primers described by Curran et al (GeneWorks, Adelaide, South Australia). PCR products

TABLE 1 | Characteristics of *P. aeruginosa* clinical isolates studied.

		CF isolates	Non-CF isolates	Total
Country of origin	Australia	2	19	21
	Netherlands	19	0	19
	USA	3	4	7
Total		24	23	47

Phylogenetic and Burst Analysis

Phylogenetic analysis of the clinical isolates and laboratory reference strain PAO1 were conducted using MEGA version 6.06 (Tamura et al., 2013). MLST sequence alleles for each of the 7 loci were aligned using the MUSCLE programme in MEGA (Edgar, 2004). The concatenated alignments were used to generate dendrograms using the Neighbor-Joining method

with bootstrapping analysis (2,000 replicates) (Felsenstein, 1985). Evolutionary distances were computed using the Maximum Composite Likelihood method, with ambiguous positions removed for each sequence pair (Tamura et al., 2004). Outgroup sequences were identified through BLAST searching for sequences of sufficient homology from other *Pseudomonas* species. eBURST version 3 (eburst.mlst.net) was used to identify clonal complexes and BURST groups (Feil et al., 2004). Bootstrapping analysis with 2000 replicates was used for all BURST analyses.

Minimum Inhibitory Concentration (MIC) Assays

Resistance to commonly used antibiotics was determined using broth microdilution minimum inhibitory concentration (MIC) assays, as described by Wiegand et al. (2008). Antibiotics tested were: gentamicin, ciprofloxacin, ceftazidime, piperacillin, and amikacin, obtained from Sigma-Aldrich (Castle Hill, NSW, Australia). Isolates were designated as being sensitive, resistant, or having intermediate sensitivity to the antibiotics based on Clinical and Laboratory Standards Institute (CLSI) cut-offs.

Sensitivity to CT-PA Phages and Enumeration of Phage

The ability of the 4 phages to lyse each bacterial isolate was tested using the spot test described by Mazzocco et al, with a drop size of 5 μ L (Mazzocco et al., 2009). All plaque assays were performed in duplicate. Enumeration of phage in stocks using the small drop plaque assay system was performed prior to each assay, with a concentration of 10^8 PFU/mL used for isolate sensitivity assays. A selected reference bacterial strain was used for titration of each of the 4 different phage stocks. The small drop plaque assay was also used to assess the phage concentration in the liquid contents of the biofilm assay wells, at 48 h after treatments were applied.

Biofilm Assay

The microtitre dish biofilm formation assay as described by O'Toole was used to assess the ability of CT-PA to eradicate *P. aeruginosa* biofilms *in vitro* (O'Toole, 2011). A 1.0 McFarland unit suspension in 0.45% saline of the isolate was diluted into 10 times the volume of nutrient broth (Oxoid, Hants, UK), and gently mixed by inversion. 150 μ L/well of the resulting suspension was plated into a clear polystyrene 96-well plate (Greiner Bio-One, Kremsmünster, Austria). Wells adjacent to the edge of the plate were filled with 180 μ L sterile PBS as a sterility control. The plate was then incubated for 48 h on a gyratory mixer at 37°C. After 48 h, the liquid contents of the bacterial wells were gently aspirated, followed by washing twice with sterile PBS to remove any remaining planktonic cells. Treatments included each of the 4 phages and the phage cocktail in nutrient broth at concentrations of 10^7 and 10^8 PFU/mL, as well as equivalent volumes of heat-inactivated stocks of the 4 phages, with PBS as negative control and 2.5% cetylpyridinium chloride (Sigma-Aldrich, St Louis, MO, USA) as a positive control. 180 μ L of each treatment was plated in quadruplicate and biofilms were assessed at 24 and 48 h after treatment. Forty-eight hours treatment plates had 50 μ L/well nutrient broth replenished at 24 h. At the designated time point (either 24 or 48 h after treatment),




FIGURE 2 | Spot test assay. Phage activity on the spot test assay was graded as: **(A)** Full activity—a completely clear (non-turbid) spot with no bacterial growth observed within the spot, **(B)** Partial activity—a turbid spot easily seen with the naked eye, or a clear spot containing isolated bacterial colonies, or **(C)** No activity—no easily discernible spot seen with the naked eye.

TABLE 2 | Host range of phages and phage cocktail in CRS clinical isolates.

<i>P. aeruginosa</i> isolate	Phage treatment				
	Pa 193	Pa 204	Pa 222	Pa 223	CT-PA
PAO1 (reference strain)	++	++	++	0	++
Aus 1	0	0	++	++	++
Aus 2	Unable to test				
Aus 3	++	++	+	++	++
Aus 4	++	0	++	++	++
Aus 5	++	++	++	++	++
Aus 6	+	0	+	+	+
Aus 7	+	0	++	++	++
Aus 8	0	0	++	++	++
Aus 9	0	0	+	+	+
Aus 10	++	++	0	0	++
Aus 11	0	0	0	0	0
Aus 12	++	++	0	0	++
Aus 13	0	0	0	0	0
Aus 14	++	++	+	0	++
Aus 15	++	++	+	+	++
Aus 16	++	++	++	++	++
Aus 17	++	++	0	0	++
Aus 18	+	+	+	+	++
Aus 19	++	++	+	+	++
Aus 20	++	++	++	++	++
Aus 21	0	0	0	0	0
Ned 1	+	0	0	++	++
Ned 2	++	++	++	++	++
Ned 3	0	0	++	++	++
Ned 4	0	0	0	0	0
Ned 5	++	++	++	++	++
Ned 6	+	++	+	+	++
Ned 7	+	+	++	++	++
Ned 8	++	0	0	++	++
Ned 9	++	++	0	0	++
Ned 10	++	++	+	++	++
Ned 11	++	++	++	++	++
Ned 12	+	0	+	0	+
Ned 13	++	0	+	+	++
Ned 14	0	0	+	+	+
Ned 15	++	++	++	+	++
Ned 16	+	0	+	0	+
Ned 17	Unable to test				
Ned 18	0	0	++	+	++
Ned 19	+	0	0	0	+
USA 1-1	0	0	0	0	0
USA 1-2	0	0	+	++	++
USA 2	++	++	++	++	++
USA 3	++	++	++	++	++
USA 4-2	++	++	++	++	++
USA 5	++	++	+	+	++
USA 6	+	+	++	+	++
Total CI sensitive (%)	33 (73)	24 (53)	33 (73)	32 (71)	40 (89)
Total CI resistant (%)	12 (27)	21 (47)	12 (27)	13 (29)	5 (11)

++, Full activity spots on small drop plaque assay.

+, Partial activity spots on small drop plaque assay.

0, No activity on small drop plaque assay.

CI, clinical isolates.

The bold values are the number (count) of isolates, as opposed to the percentage of isolates shown in parentheses.

liquid contents of the wells were transferred into a fresh plate for post-treatment phage titration. The biofilm plates were gently washed twice with sterile PBS, and then stained with 190 μ L/well 0.5% crystal violet (Sigma, St Louis, MO, USA) for 30 min. The stained plates were rinsed by two rounds of gentle immersion into distilled water, and left to dry overnight. The crystal violet stain was eluted by application of 200 μ L/well 30% acetic acid (Chem-Supply, Adelaide, South Australia) and the plate was incubated at room temperature for 30 min. Absorbance at 595 nm was measured for each well using the Fluostar Optima microplate reader (BMG Labtech, Ortenberg, Germany), with 200 μ L 30% acetic acid in unstained wells used as blanks.

Statistics

Linear mixed effects models were applied to assess the effects of group, time point, CF status and multidrug resistance on absorbance (A595) data from the microtitre plate biofilm formation assay. A595 values were log transformed prior to analysis due to violations of the distributional assumptions of linear regression. Pairwise, *post-hoc* comparisons for group were assessed at $p < 0.010$ due to the large number of comparisons made. The data were analyzed using SAS v9.4 (SAS Institute Inc., Cary, NC, USA).

RESULTS


Multiple MLST Sequence Types Identified Amongst CRS Clinical Isolates

In total, 47 *P. aeruginosa* isolates were collected from the upper and lower airways from 44 patients suffering from CRS and/or CF in 3 continents. These included 19 isolates from the upper and lower respiratory tracts of patients with CF (Amsterdam, Netherlands), 7 sinus isolates from CRS patients with and without CF (Birmingham, AL, USA), and 21 sinus isolates from CRS patients with and without CF (Adelaide, Australia). Isolate characteristics are displayed in **Table 1**. MLST revealed 44 distinct sequence types. Isolate MLST profiles and burst groups are detailed in Supplementary Table 1. One clinical isolate from a CF patient in the Netherlands was of the same sequence type as PAO1. Two Australian non-CF isolates shared an identical sequence type (348), and another sequence type (274) was shared

TABLE 3 | Antibiotic resistance of clinical isolates.

	No. of resistant isolates (%)		
	CF isolates	Non-CF isolates	All isolates
Gentamicin	9 (43)	1 (5)	10 (25)
Amikacin	10 (48)	1 (5)	11 (28)
Ciprofloxacin	13 (62)	2 (11)	15 (38)
Ceftazidime	12 (57)	1 (5)	13 (33)
Piperacillin	3 (14)	0 (0)	3 (8)
Multidrug resistant	11 (52)	0 (0)	11 (28)
Total no. of isolates tested	21	19	40

The bold values are the number (count) of isolates, as opposed to the percentage of isolates shown in parentheses.




by one isolate from each of Australia, the United States, and the Netherlands.

BURST analysis identified only one clonal complex, using the most stringent definition of sequence types sharing six or more alleles in common. The clonal complex consisted of two single locus variants isolated from the same CF patient. The phylogeny inferred by the Neighbor-Joining method is displayed in **Figure 1**.

CT-PA Host Range in CRS Clinical Isolates

40/45 isolates (89%) were lysed by CT-PA, with partial or full activity spots seen on the small drop plaque assay. Examples of full and partial activity spots are shown in **Figure 2**. Two isolates could not be tested; one due to interference from plaques throughout the bacterial lawn including areas where the phage had not been applied (thought to be due to induction of a prophage), and the other due to poor growth in culture. When



tested individually, 73, 53, 73, and 71% of isolates were sensitive to Pa 193, Pa 204, Pa 222, and Pa 223 respectively, as displayed in **Table 2**. 39/45 isolates (87%) were susceptible to 2 or more of the 4 phages.

Multidrug Resistance in CF Clinical Isolates

Isolates that displayed sensitivity to CT-PA were tested for antibiotic resistance, using CLSI cut-offs for intermediate resistance and resistance. The number of isolates displaying resistance or intermediate resistance to the five antibiotics tested is shown in **Table 3**. Eleven isolates, all from CF patients, were multidrug resistant according to the definition proposed by Magiorakos et al. (2012).

Biofilm Reduction by CT-PA

All 40 isolates that displayed sensitivity to CT-PA on the spot test assay, as well as reference strain ATCC 15692, were tested using the microtitre dish biofilm assay. Absorbance readings from the crystal violet biofilm assays of all isolates at 24 and 48 h after treatment are displayed in **Figure 3**. Statistically significant reductions in biofilm biomass compared to the negative treatment control (nutrient broth) were seen with both 10^7 and 10^8 PFU/mL of CT-PA, Pa 222 and Pa 223 at both 24 and 48 h ($p < 0.001$). Significant reductions were also seen with 10^7 PFU/mL Pa 193 and 10^8 PFU/mL Pa 204 at 48 h ($p < 0.01$).

For all isolates tested, the median biofilm biomass reduction was 70% and 64% after 48 h treatment with the lower and higher concentrations of CT-PA, respectively. 34 (85%) and 32 (80%) of the clinical isolates, respectively showed a reduction in biomass compared to the negative treatment control at 48 h, as displayed in **Figure 4**. The median biomass reductions at the 24-h time

point were similar for both concentrations of CT-PA at 67%. The maximum biomass reductions were 98.6 and 99.6% at 24 h, and 99.4 and 98.8% at 48 h, for the lower and higher concentrations, respectively.

Analysis of two-way interactions showed no statistically significant interactions between CF status or multidrug resistance and treatment effect.

Post-treatment Phage Titres

For all except three of the 40 isolates tested, titres of two or more of the four phages had increased after 48 h treatment in the biofilm assay. This indicates successful phage infection and replication, and good complementation between the four phages comprising the CT-PA cocktail. Of the remaining three isolates, two showed an increase in titres of one phage, and one did not show an increase in any phage titres. Post-treatment phage titres for PAO1 and the 40 clinical isolates are displayed in **Table 4**.

DISCUSSION

CT-PA bacteriophage cocktail displayed suitable anti-biofilm activity *in vitro*. It had a broad host range in the 45 isolates tested, with 89% of isolates susceptible. The use of a cocktail as opposed to individual phages increased the host range significantly, with only 53–73% of isolates being susceptible to each of the four phages individually. This is consistent with previous observations that the use of a cocktail of phages rather than individual phages improves activity by expanding the host range and by preventing the development of bacteriophage-insensitive mutant bacteria (Chan and Abedon, 2012; Hall et al., 2012; Drilling et al., 2014b).

The lytic effect of CT-PA on planktonic bacteria translated well to a reduction in biofilm, consistent with previous reports showing efficacy of bacteriophage to reduce biofilm *in vitro* and *in vivo* (Drilling et al., 2014a,b). Moreover, the increase in phage titres following treatment for almost all isolates implies successful phage infection, replication, and lysis of host bacteria, given the strictly lytic nature of all phages comprising the CT-PA phage cocktail.

Our results suggested slightly more biofilm removal by 10^7 PFU/mL compared to 10^8 PFU/mL CT-PA after 48 h treatment, despite similar anti-biofilm activity of the two treatment concentrations at 24 h. Studies of other *P. aeruginosa* phages have demonstrated that a higher concentration of phage treatment does not always result in greater magnitude of biofilm removal or bacterial killing (Knezevic et al., 2011; Worley-Morse et al., 2014). A non-linear relationship may be expected for some phage combinations over time due to the self-replicating nature of phage therapy (Abedon, 2011).

The isolates in our study sample consisted of 44 MLST sequence types from both the sinuses of non-CF CRS sufferers, and the upper and lower respiratory tracts of CF patients. BURST and phylogenetic analysis was consistent with the predominantly non-clonal population structure that has been described by *P. aeruginosa* strain typing studies (Kiewitz and Tümmler, 2000; Curran et al., 2004). Clonal groups (identical ST) and clonal complexes were uncommon within our collection of isolates. The clonal and BURST groups identified contain isolates from

TABLE 4 | Post-treatment phage titres.

Strain	Post-treatment phage titres (PFU/mL)							
	10 ⁸ PFU/mL treatment				10 ⁷ PFU/mL treatment			
	Pa 193	Pa 204	Pa 222	Pa 223	Pa 193	Pa 204	Pa 222	Pa 223
PAO1	2.0E+10	3.0E+09	4.0E+10	1.0E+08	1.0E+10	2.0E+09	2.0E+10	1.0E+09
Aus 1	1.1E+08	5.7E+07	4.9E+07	1.0E+02	4.3E+06	4.0E+06	1.8E+08	2.0E+02
Aus 3	1.8E+10	9.2E+09	4.3E+09	3.9E+09	9.0E+09	2.9E+09	8.5E+08	6.5E+02
Aus 4	3.0E+08	9.0E+08	3.0E+08	1.0E+07	7.0E+08	4.0E+09	5.0E+06	3.0E+05
Aus 5	8.0E+10	2.4E+09	6.4E+08	2.0E+03	5.0E+09	1.3E+10	1.3E+09	1.2E+09
Aus 6	7.0E+07	2.0E+09	3.2E+09	2.1E+05	7.8E+06	7.6E+10	8.3E+09	2.1E+06
Aus 7	2.2E+09	1.0E+08	5.0E+08	4.0E+08	9.0E+09	1.0E+07	2.0E+08	1.0E+09
Aus 8	2.0E+03	8.0E+08	2.0E+03	1.0E+07	3.0E+09	2.0E+09	1.0E+08	5.0E+05
Aus 9	2.0E+07	4.8E+03	2.4E+09	1.3E+03	4.1E+04	6.0E+02	7.8E+08	2.1E+03
Aus 10	5.0E+07	1.0E+11	9.3E+08	1.2E+06	3.0E+06	7.0E+09	2.0E+09	8.0E+05
Aus 12	2.3E+09	2.6E+10	5.1E+09	3.8E+03	7.3E+08	8.2E+10	4.5E+09	2.7E+04
Aus 14	2.7E+08	5.3E+10	9.0E+07	4.8E+06	1.7E+08	2.4E+09	2.0E+06	3.9E+05
Aus 15	1.2E+11	3.0E+10	1.2E+10	5.1E+10	2.0E+11	3.0E+11	4.0E+10	2.0E+05
Aus 16	2.0E+10	1.0E+11	3.0E+09	2.0E+08	1.0E+10	1.0E+10	2.0E+07	2.0E+07
Aus 17	1.9E+10	5.2E+11	1.9E+10	9.5E+06	5.9E+10	1.5E+11	1.5E+10	5.2E+06
Aus 18	9.0E+08	1.0E+08	2.0E+10	2.5E+08	3.0E+07	2.0E+07	7.0E+10	1.0E+08
Aus 19	5.0E+08	2.0E+03	3.0E+10	2.0E+03	2.0E+03	2.0E+03	1.0E+09	2.0E+03
Aus 20	1.0E+10	4.0E+10	9.0E+07	1.0E+10	2.0E+10	2.0E+03	1.7E+07	5.0E+05
Ned 1	4.0E+07	5.3E+06	7.0E+05	5.9E+03	2.2E+06	7.1E+06	9.5E+03	6.4E+03
Ned 2	7.2E+09	1.1E+10	5.7E+07	1.5E+08	8.0E+09	2.0E+08	2.1E+07	8.0E+07
Ned 3	8.0E+07	1.9E+07	3.6E+08	1.5E+08	9.8E+05	2.0E+06	3.5E+07	5.7E+08
Ned 5	2.5E+10	2.1E+11	4.0E+07	7.7E+07	1.8E+10	1.1E+11	4.0E+07	9.0E+05
Ned 6	8.6E+09	2.4E+10	7.1E+08	1.8E+09	5.1E+09	1.7E+10	1.2E+08	5.2E+08
Ned 7	6.8E+10	1.5E+09	7.4E+07	8.4E+07	5.5E+10	1.6E+09	1.2E+07	2.6E+08
Ned 8	4.3E+10	2.2E+12	2.2E+09	2.1E+06	7.1E+10	2.4E+05	3.7E+13	1.5E+11
Ned 9	2.2E+10	2.3E+10	1.2E+07	5.5E+10	4.4E+10	1.0E+11	2.3E+06	1.6E+04
Ned 10	1.7E+08	1.1E+10	5.5E+06	7.0E+04	7.0E+07	1.7E+10	3.1E+05	2.0E+06
Ned 11	1.9E+09	3.9E+09	1.2E+09	8.5E+08	9.0E+09	6.9E+09	1.5E+09	2.4E+07
Ned 12	2.0E+08	1.0E+08	5.0E+09	1.7E+05	2.5E+06	1.0E+08	8.0E+09	4.0E+05
Ned 13	4.0E+09	3.2E+07	4.6E+08	1.0E+03	5.4E+09	4.6E+09	6.2E+08	0.0E+00
Ned 14	1.1E+09	5.3E+09	1.5E+08	1.8E+07	1.5E+09	4.5E+09	6.9E+07	4.3E+07
Ned 15	2.4E+11	1.9E+10	5.3E+09	4.9E+06	6.3E+10	1.5E+10	7.3E+09	5.6E+06
Ned 16	2.0E+11	3.4E+08	1.7E+08	5.1E+07	6.8E+10	3.0E+08	4.8E+08	8.0E+06
Ned 18	3.2E+07	1.4E+07	3.7E+09	1.8E+06	2.3E+06	1.4E+06	5.4E+09	2.7E+06
Ned 19	4.2E+08	1.2E+09	7.0E+07	2.2E+07	2.3E+06	6.2E+09	0.0E+00	6.8E+05
USA 1-2	3.0E+07	1.3E+07	3.0E+09	1.0E+06	6.0E+05	1.0E+07	3.0E+03	2.0E+06
USA 2	1.0E+09	2.0E+10	1.0E+09	1.0E+05	2.0E+10	3.0E+10	1.0E+11	6.0E+04
USA 3	1.0E+08	1.0E+12	1.0E+10	1.0E+01	1.0E+07	2.0E+06	8.0E+09	8.0E+03
USA 4-2	1.4E+09	9.0E+09	1.6E+09	1.0E+08	1.0E+10	2.0E+10	2.0E+09	1.0E+07
USA 5	1.0E+09	9.0E+09	2.0E+10	1.0E+05	1.0E+11	1.0E+09	5.0E+10	3.0E+04
USA 6	2.6E+10	2.0E+11	1.2E+10	1.2E+06	3.8E+11	4.0E+11	8.0E+09	5.0E+08

both CF and non-CF CRS patients, suggesting that *P. aeruginosa* sinus infections in both groups may initially be caused by similar strains. This is consistent with the observation made by Cramer et al that the dominant clones amongst CF patients are also common in the environment and other human disease habitats (Cramer et al., 2012).

However, frequent antibiotic resistance was displayed by CF isolates, with 52% of the CF strains tested meeting the criteria for multidrug-resistance. Rates of antibiotic resistance in non-CF CRS isolates were much lower, and comparable to those observed in similar studies of CRS patients and in data from national antimicrobial surveillance programs (Kingdom

et al., 2004; Sasikala and Sundararaj, 2012; Sader et al., 2014, 2017; AURA, 2016). Higher rates of antibiotic resistance in *P. aeruginosa* isolates from CF patients compared to non-CF patients have been well-documented (Doring et al., 2000; Henwood et al., 2001). Although some of the isolates in our study were from the lower respiratory tract of CF patients, a high degree of similarity in the adaptive mutations and gene expression of CF sinus and lung *P. aeruginosa* isolates has been found (Ciofu et al., 2013). Furthermore, the paranasal sinuses have been found to be a reservoir of recurrent pulmonary infections in CF patients, with identical strains often identified in the upper and lower respiratory tracts of patients (Roby et al., 2008; Johansen et al., 2012; Aanaes, 2013; Aanaes et al., 2013). Thus, the selective pressures in the upper respiratory tract of CF patients, including antibiotics administered for management of bacterial sinus infections, have potential implications for the development of chronic lung infections (Hansen et al., 2012). The progression to chronic *P. aeruginosa* lung infection in CF has been shown to lead to deterioration in lung function and decreased life expectancy (Mastella et al., 1999; Li et al., 2005).

The proportion of clinical isolates that were resistant to lysis by the CT-PA phage cocktail was small (11%), and did not include any multidrug resistant isolates. Importantly, the anti-biofilm activity of CT-PA was not affected by multidrug resistance, or whether the isolate was from a CF patient. The CF isolates tested were largely comprised of isolates with a small colony or mucoid phenotype. These phenotypes are associated with chronic infection, and display increased tolerance to various antibiotics (Nichols et al., 1988; Govan and Deretic, 1996; Haussler et al., 1999; Guss et al., 2009; Hansen et al., 2012). These findings suggest that phage therapy may have potential in treating CF patients with chronic *P. aeruginosa* infection, for whom clinicians are currently faced with a dilemma of how to treat respiratory infections without creating further antibiotic resistance. We have previously demonstrated the ability of high-volume nasal irrigations to penetrate into the paranasal sinuses following functional endoscopic sinus surgery (FESS). (Grobler et al., 2008) The phage cocktail could be administered via nasal irrigations in order to treat recurrent sinus infections following FESS.

In vivo animal studies have demonstrated enhanced bacterial clearance and prolonged survival in phage-treated animals (in *Galleria mellonella*, canine otitis, murine lung, corneal and peritoneal *P. aeruginosa* infection models) (Heo et al., 2009; Debarbieux et al., 2010; Hawkins et al., 2010; Alemayehu et al., 2012; Fukuda et al., 2012; Beeton et al., 2015). A limited number of early-stage clinical trials of phage therapy for treatment of *P. aeruginosa* infections have been undertaken. A double-blind randomized controlled trial of a topically-applied phage cocktail for antibiotic-resistant *P. aeruginosa* chronic otitis showed statistically significant improvement from baseline in both symptom and clinical severity scores in the phage treatment group (Wright et al., 2009). No such improvement was observed in the placebo group. Conversely, two clinical trials of phage cocktails targeting multiple bacterial pathogens (*P. aeruginosa*, *S. aureus*, and *E. coli*) for treatment of burns and chronic leg ulcers, respectively showed no significant differences between phage and control therapies (Rhoads et al., 2009; Rose et al., 2014).

The aforementioned clinical trials all reported excellent safety profiles. A study of long-term safety of a *S. aureus* phage cocktail applied topically to sheep sinuses found no local or systemic safety concerns (Drilling et al., 2017). Although not widespread, clinical use of phage therapy has occurred through phage therapy and research institutes, such as the Eliava Institute in Tbilisi, Georgia, and the Institute of Immunology and Experimental Therapy in Wroclaw, Poland (Sulakvelidze et al., 2001; Kutateladze, 2015). An analysis of 153 patients treated for various infections with phage therapy at the latter institute revealed no major safety concerns (Międzybrodzki et al., 2012).

Whilst further evaluation and optimisation of phage cocktail treatment of *P. aeruginosa* infections in CRS and CF are required, the safety profile of phage therapy and its activity against antibiotic-resistant isolates make it an attractive potential therapeutic option. Further pre-clinical and clinical studies will be of great benefit in determining whether phage therapy can fulfill the potential demonstrated in this study.

AUTHOR CONTRIBUTIONS

SF contributed to study design, execution of experiments, data analysis, and writing of manuscript. AD contributed to study design, execution of experiments, data analysis, and review of manuscript. SM and SV contributed to study design, and review of manuscript. MC assisted with collection of clinical samples, and data collection. BW and WF contributed to collection of clinical samples, and review of manuscript. AP and PW contributed to study design, collection of clinical samples, and review of manuscript.

FUNDING

This work was supported in part by The Hospital Research Foundation, Woodville, South Australia, and Australian Government Research Training Program scholarships to SF. This work was supported in part by National Institutes of Health (NIH)/National Heart, Lung, and Blood Institute (1 R01 HL133006-02) and National Institute of Diabetes and Digestive and Kidney Diseases (5P30DK072482-04, CF Research Center Pilot Award) to BW.

ACKNOWLEDGMENTS

We wish to acknowledge the Department of Otolaryngology-Head and Neck Surgery at University of Alabama at Birmingham, and the Department of Otorhinolaryngology of the Academic Medical Centre, Amsterdam, for their assistance in the collection and donation of *P. aeruginosa* clinical isolates. We thank Dr. Stuart Howell for his assistance with statistical analyses.

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <http://journal.frontiersin.org/article/10.3389/fcimb.2017.00418/full#supplementary-material>

REFERENCES

- AURA (2016). *AURA 2016: First Australian Report on Antimicrobial Use and Resistance in Human Health*. Sydney, NSW: Australian Commission on Safety and Quality in Health Care (ACSQHC)
- Aanaes, K. (2013). Bacterial sinusitis can be a focus for initial lung colonisation and chronic lung infection in patients with cystic fibrosis. *J. Cyst. Fibros.* 12(Suppl. 2), S1–S20. doi: 10.1016/S1569-1993(13)00150-1
- Aanaes, K., von Buchwald, C., Hjuler, T., Skov, M., Alanin, M., and Johansen, H. K. (2013). The effect of sinus surgery with intensive follow-up on pathogenic sinus bacteria in patients with cystic fibrosis. *Am. J. Rhinol. Allergy* 27, e1–e4. doi: 10.2500/ajra.2013.27.3829
- Abedon, S. (2011). “Phage therapy pharmacology: calculating phage dosing,” in *Advances in Applied Microbiology*, Vol. 77, eds I. Allen, S. S. Laskin, and M. G. Geoffrey (San Diego, CA: Academic Press), 1–40.
- Alemayehu, D., Casey, P. G., McAuliffe, O., Guinane, C. M., Martin, J. G., Shanahan, F., et al. (2012). Bacteriophages phiMR299-2 and phiNH-4 can eliminate *Pseudomonas aeruginosa* in the murine lung and on cystic fibrosis lung airway cells. *MBio* 3, e00029–e00012. doi: 10.1128/mBio.00029-12
- Beeton, M. L., Alves, D. R., Enright, M. C., and Jenkins, A. T. A. (2015). Assessing phage therapy against *Pseudomonas aeruginosa* using a *Galleria mellonella* infection model. *Int. J. Antimicrob. Agents* 46, 196–200. doi: 10.1016/j.ijantimicag.2015.04.005
- Bendouah, Z., Barbeau, J., Hamad, W. A., and Desrosiers, M. (2006). Biofilm formation by *Staphylococcus aureus* and *Pseudomonas aeruginosa* is associated with an unfavorable evolution after surgery for chronic sinusitis and nasal polyposis. *Otolaryngol. Head Neck Surg.* 134, 991–996. doi: 10.1016/j.otohns.2006.03.001
- Chan, B. K., and Abedon, S. T. (2012). “Phage therapy pharmacology: phage cocktails,” in *Advances in Applied Microbiology*, Vol. 78, eds I. Allen, S. S. Laskin, and M. G. Geoffrey (San Diego, CA: Academic Press), 1–23.
- Chen, H.-H., Liu, X., Ni, C., Lu, Y.-P., Xiong, G.-Y., Lu, Y.-Y., et al. (2012). Bacterial biofilms in chronic rhinosinusitis and their relationship with inflammation severity. *Auris Nasus Larynx* 39, 169–174. doi: 10.1016/j.anl.2011.04.014
- Ciofu, O., Johansen, H. K., Aanaes, K., Wassermann, T., Alhede, M., von Buchwald, C., et al. (2013). *P. aeruginosa* in the paranasal sinuses and transplanted lungs have similar adaptive mutations as isolates from chronically infected CF lungs. *J. Cyst. Fibros.* 12, 729–736. doi: 10.1016/j.jcf.2013.02.004
- Cleland, E. J., Bassiouni, A., Vreugde, S., and Wormald, P. J. (2016). The bacterial microbiome in chronic rhinosinusitis: richness, diversity, postoperative changes, and patient outcomes. *Am. J. Rhinol. Allergy* 30, 37–43. doi: 10.2500/ajra.2016.30.4261
- Cleland, E. J., Bassiouni, A., and Wormald, P. J. (2013). The bacteriology of chronic rhinosinusitis and the pre-eminence of *Staphylococcus aureus* in revision patients. *Int. Forum Allergy Rhinol.* 3, 642–646. doi: 10.1002/alr.21159
- Cramer, N., Wiehlmann, L., Ciofu, O., Tamm, S., Høiby, N., and Tümmler, B. (2012). Molecular epidemiology of chronic *Pseudomonas aeruginosa* airway infections in cystic fibrosis. *PLoS ONE* 7:e50731. doi: 10.1371/journal.pone.0050731
- Curran, B., Jonas, D., Crundmann, H., Pitt, T., and Dowson, C. (2004). Development of a multilocus sequence typing scheme for the opportunistic pathogen *Pseudomonas aeruginosa*. *J. Clin. Microbiol.* 42, 5644–5649. doi: 10.1128/JCM.42.12.5644-5649.2004
- Debarbieux, L., Leduc, D., Maura, D., Morello, E., Criscuolo, A., Grossi, O., et al. (2010). Bacteriophages can treat and prevent *Pseudomonas aeruginosa* lung infections. *J. Infect. Dis.* 201, 1096–1104. doi: 10.1086/651135
- de la Fuente-Núñez, C., Reffuveille, F., Fernández, L., and Hancock, R. E. W. (2013). Bacterial biofilm development as a multicellular adaptation: antibiotic resistance and new therapeutic strategies. *Curr. Opin. Microbiol.* 16, 580–589. doi: 10.1016/j.mib.2013.06.013
- Doring, G., Conway, S. P., Heijerman, H. G., Hodson, M. E., Høiby, N., Smyth, A., et al. (2000). Antibiotic therapy against *Pseudomonas aeruginosa* in cystic fibrosis: a European consensus. *Eur. Respir. J.* 16, 749–767. doi: 10.1034/j.1399-3003.2000.16d30.x
- Drilling, A. J., Ooi, M. L., Miljkovic, D., James, C., Speck, P., Vreugde, S., et al. (2017). Long-term safety of topical Bacteriophage application to the frontal sinus region. *Front. Cell. Infect. Microbiol.* 7:49. doi: 10.3389/fcimb.2017.00049
- Drilling, A., Morales, S., Boase, S., Jervis-Bardy, J., James, C., Jardeleza, C., et al. (2014a). Safety and efficacy of topical bacteriophage and ethylenediaminetetraacetic acid treatment of *Staphylococcus aureus* infection in a sheep model of sinusitis. *Int. Forum Allergy Rhinol.* 4, 176–186. doi: 10.1002/alr.21270
- Drilling, A., Morales, S., Jardeleza, C., Vreugde, S., Speck, P., and Wormald, P.-J. (2014b). Bacteriophage reduces biofilm of *Staphylococcus aureus ex vivo* isolates from chronic rhinosinusitis patients. *Am. J. Rhinol. Allergy* 28, 3–11. doi: 10.2500/ajra.2014.28.4001
- Edgar, R. C. (2004). MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Res.* 32, 1792–1797. doi: 10.1093/nar/gkh340
- Feil, E. J., Li, B. C., Aanensen, D. M., Hanage, W. P., and Spratt, B. G. (2004). eBURST: inferring patterns of evolutionary descent among clusters of related bacterial genotypes from multilocus sequence typing data. *J. Bacteriol.* 186, 1518–1530. doi: 10.1128/JB.186.5.1518-1530.2004
- Felsenstein, J. (1985). Confidence limits on phylogenies: an approach using the bootstrap. *Evolution* 39, 783–791. doi: 10.1111/j.1558-5646.1985.tb00420.x
- Fokkens, W. J., Lund, V. J., Mullol, J., Bachert, C., Alobid, I., Baroody, F., et al. (2012). EPOS 2012: European position paper on rhinosinusitis and nasal polyps 2012. A summary for otorhinolaryngologists. *Rhinology* 50, 1–12. doi: 10.4193/Rhino50E2
- Fukuda, K., Ishida, W., Uchiyama, J., Rashed, M., Kato, S., Morita, T., et al. (2012). *Pseudomonas aeruginosa* keratitis in mice: effects of topical bacteriophage KPP12 administration. *PLoS ONE* 7:e47742. doi: 10.1371/journal.pone.0047742
- Govan, J. R., and Deretic, V. (1996). Microbial pathogenesis in cystic fibrosis: mucoid *Pseudomonas aeruginosa* and *Burkholderia cepacia*. *Microbiol. Rev.* 60, 539–574.
- Grobler, A., Weitzel, E. K., Buele, A., Jardeleza, C., Cheong, Y. C., Field, J., et al. (2008). Pre- and postoperative sinus penetration of nasal irrigation. *Laryngoscope* 118, 2078–2081. doi: 10.1097/MLG.0b013e31818208c1
- Guss, J., Abuzeid, W. M., Doghramji, L., Edelstein, P. H., and Chiu, A. G. (2009). Fluoroquinolone-resistant *Pseudomonas aeruginosa* in chronic rhinosinusitis. *ORL. J. Otorhinolaryngol. Relat. Spec.* 71, 263–267. doi: 10.1159/000242428
- Guttman, B., Raya, R., and Kutter, E. (2004). “Basic phage biology,” in *Bacteriophages: Biology and Applications*, eds E. Kutter and A. Sulakvelidze (Boca Raton, FL: CRC Press), 30–63.
- Hall, A. R., Vos, D., Friman, V. P., Pirnay, J. P., and Buckling, A. (2012). Effects of sequential and simultaneous application of bacteriophages on populations of *Pseudomonas aeruginosa in vitro* and in waxmoth larvae. *Appl. Environ. Microbiol.* 78, 5646–5652. doi: 10.1128/AEM.00757-12.
- Hanlon, G. W. (2007). Bacteriophages: an appraisal of their role in the treatment of bacterial infections. *Int. J. Antimicrob. Agents* 30, 118–128. doi: 10.1016/j.ijantimicag.2007.04.006
- Hansen, S. K., Rau, M. H., Johansen, H. K., Ciofu, O., Jelsbak, L., Yang, L., et al. (2012). Evolution and diversification of *Pseudomonas aeruginosa* in the paranasal sinuses of cystic fibrosis children have implications for chronic lung infection. *ISME J.* 6, 31–45. doi: 10.1038/ismej.2011.83
- Hausler, S., Tümmler, B., Weissbrodt, H., Rohde, M., and Steinmetz, I. (1999). Small-colony variants of *Pseudomonas aeruginosa* in cystic fibrosis. *Clin. Infect. Dis.* 29, 621–625. doi: 10.1086/598644
- Hawkins, C., Harper, D., Burch, D., Anggard, E., and Soothill, J. (2010). Topical treatment of *Pseudomonas aeruginosa* otitis of dogs with a bacteriophage mixture: a before/after clinical trial. *Vet. Microbiol.* 146, 309–313. doi: 10.1016/j.vetmic.2010.05.014
- Henwood, C. J., Livermore, D. M., James, D., and Warner, M. (2001). *Pseudomonas* Study Group t. Antimicrobial susceptibility of *Pseudomonas aeruginosa*: results of a UK survey and evaluation of the British Society for Antimicrobial Chemotherapy disc susceptibility test. *J. Antimicrob. Chemother.* 47, 789–799. doi: 10.1093/jac/47.6.789
- Heo, Y. J., Lee, Y. R., Jung, H. H., Lee, J., Ko, G., and Cho, Y. H. (2009). Antibacterial efficacy of phages against *Pseudomonas aeruginosa* infections in mice and *Drosophila melanogaster*. *Antimicrob. Agents Chemother.* 53, 2469. doi: 10.1128/AAC.01646-08
- Johansen, H. K., Aanaes, K., Pressler, T., Nielsen, K. G., Fisker, J., Skov, M., et al. (2012). Colonisation and infection of the paranasal sinuses in cystic fibrosis patients is accompanied by a reduced PMN response. *J. Cyst. Fibros.* 11, 525–531. doi: 10.1016/j.jcf.2012.04.011

- Kiewitz, C., and Tümler, B. (2000). Sequence diversity of *Pseudomonas aeruginosa*: impact on population structure and Genome evolution. *J. Bacteriol.* 182, 3125–3135. doi: 10.1128/JB.182.11.3125-3135.2000
- Kingdom, T. T., and Swain, R. E. (2004). The microbiology and antimicrobial resistance patterns in chronic rhinosinusitis. *Am. J. Otolaryngol.* 25, 323–328. doi: 10.1016/j.amjoto.2004.03.003
- Knezevic, P., Obrecht, D., Curcin, S., Petrusic, M., Aleksic, V., Kostanjsek, R., et al. (2011). Phages of *Pseudomonas aeruginosa*: response to environmental factors and *in vitro* ability to inhibit bacterial growth and biofilm formation. *J. Appl. Microbiol.* 111, 245–254. doi: 10.1111/j.1365-2672.2011.05043.x
- Kutateladze, M. (2015). Experience of the Eliava Institute in bacteriophage therapy. *Virol. Sin.* 30, 80–81. doi: 10.1007/s12250-014-3557-0
- Li, Z., Kosorok, M. R., Farrell, P. M., Laxova, A., West, S. E. H., Green, C. G., et al. (2005). Longitudinal development of mucoid *Pseudomonas aeruginosa* infection and lung disease progression in children with cystic fibrosis. *JAMA* 293, 581–588. doi: 10.1001/jama.293.5.581
- Magiorakos, A. P., Srinivasan, A., Carey, R. B., Carmeli, Y., Falagas, M. E., Giske, C. G., et al. (2012). Multidrug-resistant, extensively drug-resistant and pandrug-resistant bacteria: an international expert proposal for interim standard definitions for acquired resistance. *Clin. Microbiol. Infect.* 18, 268–281. doi: 10.1111/j.1469-0691.2011.03570.x
- Mastella, G., Hodson, M. E., Koch, C., Navarro, J., Rainisio, M., Harms, H. K., et al. (1999). Factors affecting velocity of lung function deterioration in ERCF cystic fibrosis patients over 2–4 years follow-up. *Neth. J. Med.* 54:S22. doi: 10.1016/S0300-2977(99)90050-5
- Mazzocco, A., Waddell, T. E., Lingohr, E., and Johnson, R. P. (2009). “Enumeration of Bacteriophages Using the Small Drop Plaque Assay System,” in *Bacteriophages: Methods and Protocols*, Vol. 1, *Isolation, Characterization, and Interactions*, eds M. R. J. Clokie and A. M. Kropinski (Totowa, NJ: Humana Press), 81–85.
- Międzybrodzki, R., Borysowski, J., Weber-Dąbrowska, B., Fortuna, W., Letkiewicz, S., Szufnarowski, K., et al. (2012). “Clinical aspects of phage therapy,” in *Advances in Virus Research*, Vol. 83, eds Małgorzata and S. Waclaw (San Diego, CA: Academic Press), 73–121.
- Nichols, W. W., Dorrington, S. M., Slack, M. P., and Walmsley, H. L. (1988). Inhibition of tobramycin diffusion by binding to alginate. *Antimicrob. Agents Chemother.* 32, 518–523. doi: 10.1128/AAC.32.4.518
- O’Toole, G. A. (2011). Microtiter dish biofilm formation assay. *J. Vis. Exp.* e2437. doi: 10.3791/2437
- Psaltis, A. J., Ha, K. R., Beule, A. G., Tan, L. W., and Wormald, P. J. (2007). Confocal scanning laser microscopy evidence of biofilms in patients with chronic rhinosinusitis. *Laryngoscope* 117, 1302–1306. doi: 10.1097/MLG.0b013e31806009b0
- Rasmussen, J., Aanaes, K., Norling, R., Nielsen, K. G., Johansen, H. K., and von Buchwald, C. (2012). CT of the paranasal sinuses is not a valid indicator for sinus surgery in CF patients. *J. Cyst. Fibros.* 11, 93–99. doi: 10.1016/j.jcf.2011.09.009
- Rhoads, D. D., Wolcott, R. D., Kuskowski, M. A., Wolcott, B. M., Ward, L. S., and Sulakvelidze, A. (2009). Bacteriophage therapy of venous leg ulcers in humans: results of a phase I safety trial. *J. Wound Care* 18, 237–243. doi: 10.12968/jowc.2009.18.6.42801
- Roby, B. B., McNamara, J., Finkelstein, M., and Sidman, J. (2008). Sinus surgery in cystic fibrosis patients: comparison of sinus and lower airway cultures. *Int. J. Pediatr. Otorhinolaryngol.* 72, 1365–1369. doi: 10.1016/j.ijporl.2008.05.011
- Rose, T., Verbeke, G., de Vos, D., Merabishvili, M., Vanechoutte, M., Lavigne, R., et al. (2014). Experimental phage therapy of burn wound infection: Difficult first steps. *Int. J. Burns Trauma* 4, 66–73.
- Sader, H. S., Farrell, D. J., Flamm, R. K., and Jones, R. N. (2014). Antimicrobial susceptibility of Gram-negative organisms isolated from patients hospitalised with pneumonia in US and European hospitals: results from the SENTRY Antimicrobial Surveillance Program, 2009–2012. *Int. J. Antimicrob. Agents* 43, 328–334. doi: 10.1016/j.ijantimicag.2014.01.007
- Sader, H. S., Huband, M. D., Castanheira, M., and Flamm, R. K. (2017). *Pseudomonas aeruginosa* antimicrobial susceptibility results from four years (2012 to 2015) of the international network for optimal resistance monitoring program in the United States. *Antimicrob. Agents Chemother.* 61:e02252-16. doi: 10.1128/AAC.02252-16
- Sasikala, S., and Sundararaj, T. (2012). Characterization of *Pseudomonas* strains from sinusitis patients. *BMC Infect. Dis.* 12:P64. doi: 10.1186/1471-2334-12-S1-P64
- Sulakvelidze, A., Alavidze, Z., and Morris, J. G. (2001). Bacteriophage Therapy. *Antimicrob. Agents Chemother.* 45, 649–659. doi: 10.1128/AAC.45.3.649-659.2001
- Tamura, K., Nei, M., and Kumar, S. (2004). Prospects for inferring very large phylogenies by using the neighbor-joining method. *Proc. Natl. Acad. Sci. U.S.A.* 101, 11030–11035. doi: 10.1073/pnas.0404206101
- Tamura, K., Stecher, G., Peterson, D., Filipski, A., and Kumar, S. (2013). MEGA6: molecular evolutionary genetics analysis version 6.0. *Mol. Biol. Evol.* 30, 2725–2729. doi: 10.1093/molbev/mst197
- Vilas Boas, D., Almeida, C., Sillankorva, S., Nicolau, A., Azeredo, J., and Azevedo, N. F. (2016). Discrimination of bacteriophage infected cells using locked nucleic acid fluorescent *in situ* hybridization (LNA-FISH). *Biofouling* 32, 179–190. doi: 10.1080/08927014.2015.1131821
- Wiegand, I., Hilpert, K., and Hancock, R. E. W. (2008). Agar and broth dilution methods to determine the minimal inhibitory concentration (MIC) of antimicrobial substances. *Nat. Protocols* 3, 163–175. doi: 10.1038/nprot.2007.521
- Worley-Morse, T. O., Zhang, L., and Gunsch, C. K. (2014). The long-term effects of phage concentration on the inhibition of planktonic bacterial cultures. *Env. Sci. Process. Impacts* 16, 81–87. doi: 10.1039/C3EM00427A
- Wright, A., Hawkins, C. H., Anggard, E. E., and Harper, D. R. (2009). A controlled clinical trial of a therapeutic bacteriophage preparation in chronic otitis due to antibiotic-resistant *Pseudomonas aeruginosa*; a preliminary report of efficacy. *Clin. Otolaryngol.* 34, 349–357. doi: 10.1111/j.1749-4486.2009.01973.x

Conflict of Interest Statement: SM is an employee of AmpliPhi Australia, a biotechnology company focused on the development of bacteriophage products.

The other authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest

Copyright © 2017 Fong, Drilling, Morales, Cornet, Woodworth, Fokkens, Psaltis, Vreugde and Wormald. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.