

Azhar Iqbal, Derek Abbott

**A game theoretical perspective on the quantum probabilities associated with a GHZ state**

Quantum Information Processing, 2018; 17(11):313-1-313-13

© Springer Science+Business Media, LLC, part of Springer Nature 2018

*This is a post-peer-review, pre-copyedit version of an article published in **Quantum Information Processing**. The final authenticated version is available online at:*

<http://dx.doi.org/10.1007/s11128-018-2083-7>

**PERMISSIONS**

<https://www.springer.com/gp/open-access/publication-policies/self-archiving-policy>

**Self-archiving for articles in subscription-based journals**

Springer journals' [policy on preprint sharing](#).

By signing the Copyright Transfer Statement you still retain substantial rights, such as self-archiving:

*Author(s) are permitted to self-archive a pre-print and an author's **accepted manuscript** version of their Article.*

.....

*b. An Author's Accepted Manuscript (AAM) is the version accepted for publication in a journal following peer review but prior to copyediting and typesetting that can be made available under the following conditions:*

*(i) Author(s) retain the right to make an AAM of their Article available on their own personal, self-maintained website immediately on acceptance,*

*(ii) Author(s) retain the right to make an AAM of their Article available for public release on any of the following 12 months after first publication ("Embargo Period"): their employer's internal website; their institutional and/or funder repositories. AAMs may also be deposited in such repositories immediately on acceptance, provided that they are not made publicly available until after the Embargo Period.*

*An acknowledgement in the following form should be included, together with a link to the published version on the publisher's website: "This is a post-peer-review, pre-copyedit version of an article published in [insert journal title]. The final authenticated version is available online at: [http://dx.doi.org/\[insert DOI\]](http://dx.doi.org/[insert DOI])".*

When publishing an article in a subscription journal, without open access, authors sign the Copyright Transfer Statement (CTS) which also details Springer's self-archiving policy.

See Springer Nature [terms of reuse](#) for archived author accepted manuscripts (AAMs) of subscription articles.

**19 June 2020**

# A game theoretical perspective on the quantum probabilities associated with a GHZ state

Azhar Iqbal and Derek Abbott

School of Electrical & Electronic Engineering, the University of Adelaide,  
South Australia 5005, Australia.

## Abstract

In the standard approach to quantum games, players' moves are local unitary transformations on an entangled state that is subsequently measured. Players' payoffs are then obtained as expected values of the entries in the payoff matrix of the classical game on a set of quantum probabilities obtained from the quantum measurement. In this paper, we approach quantum games from a diametrically opposite perspective. We consider a classical three-player symmetric game along with a known expression for a set of quantum probabilities relevant to a tripartite Einstein-Podolsky-Rosen (EPR) experiment that depends on three players' directional choices in the experiment. We define the players' moves as their directional choices in an EPR setting and then express their payoff relations in the resulting quantum game in terms of their directional choices, the entries of the payoff matrix, and the quantum probability distribution relevant to the tripartite EPR experiment.

## 1 Introduction

In the standard scheme [1, 2] of a quantized version of a non-cooperative game [3], the players share an entangled state, their strategic moves are local unitary transformations on the state, and the quantum measurement [4] generates the players' payoffs. The resulting players' payoffs in the quantum game can be understood as the expected values of the entries in the payoff matrix of the (classical) game [5–7] arising from a set of quantum probabilities [4]. The key concerns in determining the players' payoffs relations in the quantum game are a) What are the players' moves in the quantum game? b) Which set of quantum probabilities is obtained by quantum measurement? and c) How the players' strategic moves are related to the set of quantum probabilities?

This brings us to question whether the unitary transformations are really necessary in the setup of a quantum game. A proposed scheme [8–10, 12] for playing a quantum game in which players' strategic moves are not unitary transformations uses the setting of an Einstein-Podolsky-Rosen (EPR) experiment [4, 14–18]. Two players are located in spacelike-separated regions and share a singlet state. In a run of the experiment, each player decides one out of the two available directions and a quantum measurement is performed. This leads to obtaining a (normalized) set of quantum probabilities along with a listing of the directional choices the players make in each run of the experiment. As the players' directional choices determine the quantum probability distribution, the setting can be used to develop a quantum version of a two-player game. A multipartite EPR experiment would then be required for a multiplayer quantum game.

In this paper, we consider a classical three-player symmetric game, along with a reported expression for a quantum probability distribution, which is relevant to the three-partite EPR experiment. We then define players' directional choices in the experiment as their strategic moves and express players' payoff relations in the quantum game in terms of the three directional choices and the entries of the payoff matrix.

This paper thus provides a game-theoretic perspective on the peculiarity of quantum probabilities. The first perspective along game-theoretical lines on quantum probabilities that are associated to the GHZ state [4] was provided by Vaidman in Ref. [19]. Vaidman proposed a set of rules defining a game that can only be won by a team of three players when they share a GHZ

state. The present paper extends Vaidman's perspective by considering Nash equilibria in the set of symmetric games played by a team of three players in a non-cooperative game setting. Vaidman presented his game without invoking Hilbert space as is the case in the present paper.

## 2 Three-player games with mixed-strategies

Consider a three-player (noncooperative) game in which the players Alice ( $A$ ), Bob ( $B$ ), and Chris ( $C$ ) make their strategic moves simultaneously. The players are assumed located at distance and are unable to communicate to one another. They, however, can communicate to a referee who organizes the game and ensures that the rules of the game are obeyed. Each player has to decide between two choices, called the *pure strategies*, and in repeated version of the game they can also play the *mixed strategies*. Their payoff relations are made public by the referee at the start of the game. The payoff relations depend on the game matrix, the players' pure strategies, and the probability distribution on pure strategies.

To be specific, we assume that the player  $A$ 's pure strategies are  $S_1, S_2$ ; the player  $B$ 's pure strategies are  $S'_1, S'_2$ ; and the player  $C$ 's pure strategies are  $S''_1, S''_2$ . Also, the game is defined by the following pure-strategy payoff relations [21]

$$\begin{aligned} \Pi_{A,B,C}(S_1, S'_1, S''_1) &= \alpha_1, \beta_1, \gamma_1; & \Pi_{A,B,C}(S_1, S'_2, S''_2) &= \alpha_5, \beta_5, \gamma_5; \\ \Pi_{A,B,C}(S_2, S'_1, S''_1) &= \alpha_2, \beta_2, \gamma_2; & \Pi_{A,B,C}(S_2, S'_1, S''_2) &= \alpha_6, \beta_6, \gamma_6; \\ \Pi_{A,B,C}(S_1, S'_2, S''_1) &= \alpha_3, \beta_3, \gamma_3; & \Pi_{A,B,C}(S_2, S'_2, S''_1) &= \alpha_7, \beta_7, \gamma_7; \\ \Pi_{A,B,C}(S_1, S'_1, S''_2) &= \alpha_4, \beta_4, \gamma_4; & \Pi_{A,B,C}(S_2, S'_2, S''_2) &= \alpha_8, \beta_8, \gamma_8. \end{aligned} \quad (1)$$

For example,  $\Pi_{A,B,C}(S_1, S'_2, S''_1) = \alpha_3, \beta_3, \gamma_3$  states that the players  $A, B$ , and  $C$  obtain the payoffs  $\alpha_3, \beta_3$ , and  $\gamma_3$ , respectively, when they play the pure strategies  $S_1, S'_2$ , and  $S''_1$ , respectively.

In a repeated version of this game, a player can choose between his/her two pure strategies with some probability, which defines his/her mixed-strategy. We specify a mixed-strategy by  $x, y, z \in [0, 1]$  for players  $A, B$ , and  $C$ , respectively. These are the probabilities with which the players  $A, B$ , and  $C$  play the pure strategies  $S_1, S'_1$ , and  $S''_1$ , respectively. They, then, play the pure strategies  $S_2, S'_2$ , and  $S''_2$  with probabilities  $(1 - x), (1 - y)$ , and  $(1 - z)$ , respectively, and the mixed-strategy payoff relations, therefore, read

$$\begin{aligned} \Pi_{A,B,C}(x, y, z) &= xyz\Pi_{A,B,C}(S_1, S'_1, S''_1) + x(1 - y)z\Pi_{A,B,C}(S_1, S'_2, S''_1) + \\ &xy(1 - z)\Pi_{A,B,C}(S_1, S'_1, S''_2) + x(1 - y)(1 - z)\Pi_{A,B,C}(S_1, S'_2, S''_2) + \\ &(1 - x)yz\Pi_{A,B,C}(S_2, S'_1, S''_1) + (1 - x)(1 - y)z\Pi_{A,B,C}(S_2, S'_2, S''_1) + \\ &(1 - x)y(1 - z)\Pi_{A,B,C}(S_2, S'_1, S''_2) + (1 - x)(1 - y)(1 - z)\Pi_{A,B,C}(S_2, S'_2, S''_2), \end{aligned} \quad (2)$$

that can also be written as

$$\Pi_{A,B,C}(x, y, z) = \sum_{i,j,k=1,2} \Pr_c(S_i, S'_j, S''_k) \Pi_{A,B,C}(S_i, S'_j, S''_k), \quad (3)$$

where  $\Pr_c(S_i, S'_j, S''_k)$  are the classical factorizable probabilities and for instance,  $\Pr_c(S_1, S'_1, S''_2) = xy(1 - z)$  and  $\Pr_c(S_2, S'_2, S''_1) = (1 - x)(1 - y)z$ .

### 2.1 Symmetric three-player games

Symmetric three-player games are defined by the condition that a player's payoff is decided by his/her strategic move and not by his/her identity. Mathematically, this is expressed by the conditions

$$\Pi_A(x, y, z) = \Pi_A(x, z, y) = \Pi_B(y, x, z) = \Pi_B(z, x, y) = \Pi_C(y, z, x) = \Pi_C(z, y, x), \quad (4)$$

i.e. the player  $A$ 's payoff when s/he plays  $x$  remains the same either when player  $B$  plays  $y$  whereas player  $C$  plays  $y$  or when player  $B$  plays  $x$  whereas player  $C$  plays  $x$ . The payoff relations

(2) satisfy the conditions (4) when [21]

$$\begin{aligned}
\beta_1 &= \alpha_1, & \beta_2 &= \alpha_3, & \beta_3 &= \alpha_2, & \beta_4 &= \alpha_3, \\
\beta_5 &= \alpha_6, & \beta_6 &= \alpha_5, & \beta_7 &= \alpha_6, & \beta_8 &= \alpha_8, \\
\gamma_1 &= \alpha_1, & \gamma_2 &= \alpha_3, & \gamma_3 &= \alpha_3, & \gamma_4 &= \alpha_2, \\
\gamma_5 &= \alpha_6, & \gamma_6 &= \alpha_6, & \gamma_7 &= \alpha_5, & \gamma_8 &= \alpha_8, \\
&& \alpha_6 &= \alpha_7, & \alpha_3 &= \alpha_4.
\end{aligned} \tag{5}$$

A symmetric three-player game can, therefore, be defined by only six constants  $\alpha_1, \alpha_2, \alpha_3, \alpha_5, \alpha_6,$  and  $\alpha_8$ . In the rest of this paper we will define these six constants to be  $\alpha, \beta, \delta, \epsilon, \theta,$  and  $\omega$ , where  $\alpha_1 = \alpha, \alpha_2 = \beta, \alpha_3 = \delta, \alpha_5 = \epsilon, \alpha_6 = \theta,$  and  $\alpha_8 = \omega$ . The pure-strategy payoff relations (1) in this symmetric game are then re-expressed as

$$\begin{aligned}
\Pi_{A,B,C}(S_1, S'_1, S''_1) &= \alpha, \alpha, \alpha; & \Pi_{A,B,C}(S_1, S'_2, S''_2) &= \epsilon, \theta, \theta; \\
\Pi_{A,B,C}(S_2, S'_1, S''_1) &= \beta, \delta, \delta; & \Pi_{A,B,C}(S_2, S'_1, S''_2) &= \theta, \epsilon, \theta; \\
\Pi_{A,B,C}(S_1, S'_2, S''_1) &= \delta, \beta, \delta; & \Pi_{A,B,C}(S_2, S'_2, S''_1) &= \theta, \theta, \epsilon; \\
\Pi_{A,B,C}(S_1, S'_1, S''_2) &= \delta, \delta, \beta; & \Pi_{A,B,C}(S_2, S'_2, S''_2) &= \omega, \omega, \omega.
\end{aligned} \tag{6}$$

The mixed-strategy payoff relations in Eq. (2) then take the form

$$\begin{aligned}
\Pi_{A,B,C}(x, y, z) &= xyz(\alpha, \alpha, \alpha) + x(1-y)z(\delta, \beta, \delta) + xy(1-z)(\delta, \delta, \beta) + \\
&x(1-y)(1-z)(\epsilon, \theta, \theta) + (1-x)yz(\beta, \delta, \delta) + (1-x)(1-y)z(\theta, \theta, \epsilon) + \\
&(1-x)y(1-z)(\theta, \epsilon, \theta) + (1-x)(1-y)(1-z)(\omega, \omega, \omega).
\end{aligned} \tag{7}$$

### 3 Quantum probability distribution for a GHZ state

Consider the GHZ state

$$|\psi\rangle = (|0\rangle_1 |0\rangle_2 |0\rangle_3 + |1\rangle_1 |1\rangle_2 |1\rangle_3) / \sqrt{2}, \tag{8}$$

that is shared among three the three players, where  $|i\rangle_j$  is the  $i$ -th state of the  $j$ -th qubit and the setting of the generalized EPR experiments. Each player measures the dichotomic observable  $\vec{n} \cdot \vec{\sigma}$  where  $\vec{n} = \vec{a}, \vec{b}, \vec{c}$  and  $\vec{\sigma}$  is a vector the components of which are standard Pauli matrices. The family of observables  $\vec{n} \cdot \vec{\sigma}$  covers all possible dichotomic observables for a qubit system [4].

Kaszlikowski and Żukowski [20] show that the probability of obtaining the result  $m = \pm 1$  for the player  $A$ , when s/he plays the strategy  $\vec{a}$ , the result  $l = \pm 1$  for the player  $B$ , when s/he plays the strategy  $\vec{b}$  and the result  $k = \pm 1$  for the player  $C$ , when s/he plays the strategy  $\vec{c}$  is given by

$$\Pr_{QM}(m, l, k; \vec{a}, \vec{b}, \vec{c}) = \frac{1}{8} \left[ 1 + mla_3b_3 + mka_3c_3 + lkb_3c_3 + mlk \sum_{r,p,s=1}^3 M_{rps} a_r b_p c_s \right], \tag{9}$$

where  $a_r, b_p, c_s$  are components of vectors  $\vec{a}, \vec{b}, \vec{c}$  and where nonzero elements of the tensor  $M_{rps}$  are  $M_{111} = 1, M_{122} = -1, M_{212} = -1, M_{221} = -1$ . In view of this, the only terms in the product  $a_r b_p c_s$  that contribute towards the probability  $\Pr_{QM}(m, l, k; \vec{a}, \vec{b}, \vec{c})$  are  $a_1 b_1 c_1, a_1 b_2 c_2, a_2 b_1 c_2,$  and  $a_2 b_2 c_1$ . Eq. (9) can therefore be written as

$$\Pr_{QM}(m, l, k; \vec{a}, \vec{b}, \vec{c}) = \frac{1}{8} [1 + mla_3b_3 + mka_3c_3 + lkb_3c_3 + mlk(a_1 b_1 c_1 - a_1 b_2 c_2 - a_2 b_1 c_2 - a_2 b_2 c_1)]. \tag{10}$$

Note that Eq. (9) gives a quantum probability distribution without reference to the undelying Hilbert space, unitary transformations, or quantum measurement.

We consider playing a three-player quantum game in which the players Alice, Bob, and Chris (henceforth, labelled as player  $A$ , player  $B$ , and player  $C$ ) moves consist of choosing the directions  $\vec{a}, \vec{b},$  and  $\vec{c}$ , respectively. The players's payoff relations are then expressed in terms of the quantum probability distribution given in Eq. (9).

### 3.1 Players sharing a GHZ state and when choosing a direction is a player's move

Let  $\vec{a} = \vec{a}(a_1, a_2, a_3)$ ,  $\vec{b} = \vec{b}(b_1, b_2, b_3)$ ,  $\vec{c} = \vec{c}(c_1, c_2, c_3)$  be the players' directional choices that we consider as their strategies. Denoting the quantum probabilities by  $\Pr_Q$ , the set of quantum probabilities can be obtained from Eq. (10) as follows

$$\begin{aligned}
\Pr_Q(S_1, S'_1, S''_1) &= \Pr_Q[(\vec{a}, m = +1), (\vec{b}, l = +1), (\vec{c}, k = +1)] \\
&= \frac{1}{8} [1 + a_3 b_3 + a_3 c_3 + b_3 c_3 + \Delta]; \\
\Pr_Q(S_1, S'_2, S''_1) &= \Pr_Q[(\vec{a}, m = +1), (\vec{b}, l = -1), (\vec{c}, k = +1)] \\
&= \frac{1}{8} [1 - a_3 b_3 + a_3 c_3 - b_3 c_3 - \Delta]; \\
\Pr_Q(S_1, S'_1, S''_2) &= \Pr_Q[(\vec{a}, m = +1), (\vec{b}, l = +1), (\vec{c}, k = -1)] \\
&= \frac{1}{8} [1 + a_3 b_3 - a_3 c_3 - b_3 c_3 - \Delta]; \\
\Pr_Q(S_1, S'_2, S''_2) &= \Pr_Q[(\vec{a}, m = +1), (\vec{b}, l = -1), (\vec{c}, k = -1)] \\
&= \frac{1}{8} [1 - a_3 b_3 - a_3 c_3 + b_3 c_3 + \Delta]; \\
\Pr_Q(S_2, S'_1, S''_1) &= \Pr_Q[(\vec{a}, m = -1), (\vec{b}, l = +1), (\vec{c}, k = +1)] \\
&= \frac{1}{8} [1 - a_3 b_3 - a_3 c_3 + b_3 c_3 - \Delta]; \\
\Pr_Q(S_2, S'_2, S''_1) &= \Pr_Q[(\vec{a}, m = -1), (\vec{b}, l = -1), (\vec{c}, k = +1)] \\
&= \frac{1}{8} [1 + a_3 b_3 - a_3 c_3 - b_3 c_3 + \Delta]; \\
\Pr_Q(S_2, S'_1, S''_2) &= \Pr_Q[(\vec{a}, m = -1), (\vec{b}, l = +1), (\vec{c}, k = -1)] \\
&= \frac{1}{8} [1 - a_3 b_3 + a_3 c_3 - b_3 c_3 + \Delta]; \\
\Pr_Q(S_2, S'_2, S''_2) &= \Pr_Q[(\vec{a}, m = -1), (\vec{b}, l = -1), (\vec{c}, k = -1)] \\
&= \frac{1}{8} [1 + a_3 b_3 + a_3 c_3 + b_3 c_3 - \Delta]; \tag{11}
\end{aligned}$$

where  $\Delta = a_1 b_1 c_1 - a_1 b_2 c_2 - a_2 b_1 c_2 - a_2 b_2 c_1$ . We define players  $A$ 's,  $B$ 's,  $C$ 's payoff relations in the quantum game as follows

$$\Pi_{A,B,C}(\vec{a}, \vec{b}, \vec{c}) = \sum_{i,j,k=1}^2 \Pr_Q(S_i, S'_j, S''_k) \Pi_{A,B,C}(S_i, S'_j, S''_k), \tag{12}$$

i.e. these are obtained as the expectation of payoff entries (6) on the set of quantum probabilities (11). For the symmetric game defined in Eq. (6), the payoffs to the players  $A$ ,  $B$ , and  $C$ , given in (12), can then be expanded as follows:

$$\begin{aligned}
\Pi_{A,B,C}(\vec{a}, \vec{b}, \vec{c}) &= \\
&\frac{1}{8} \{ [1 + a_3 b_3 + a_3 c_3 + b_3 c_3 + \Delta] (\alpha, \alpha, \alpha) + [1 - a_3 b_3 + a_3 c_3 - b_3 c_3 - \Delta] (\delta, \beta, \delta) + \\
&[1 + a_3 b_3 - a_3 c_3 - b_3 c_3 - \Delta] (\delta, \delta, \beta) + [1 - a_3 b_3 - a_3 c_3 + b_3 c_3 + \Delta] (\epsilon, \theta, \theta) + \\
&[1 - a_3 b_3 - a_3 c_3 + b_3 c_3 - \Delta] (\beta, \delta, \delta) + [1 + a_3 b_3 - a_3 c_3 - b_3 c_3 + \Delta] (\theta, \theta, \epsilon) + \\
&[1 - a_3 b_3 + a_3 c_3 - b_3 c_3 + \Delta] (\theta, \epsilon, \theta) + [1 + a_3 b_3 + a_3 c_3 + b_3 c_3 - \Delta] (\omega, \omega, \omega) \}. \tag{13}
\end{aligned}$$

Let  $a_3 = b_3 = c_3 = 0$  i.e. when the players' unit vectors are confined to the X-Y plane, the payoff relations (13) can be written as

$$\begin{aligned} \Pi_{A,B,C}(\vec{a}, \vec{b}, \vec{c}) = \\ \frac{1}{8}\{(1+\Delta)(\alpha, \alpha, \alpha) + (1-\Delta)(\delta, \beta, \delta) + (1-\Delta)(\delta, \delta, \beta) + (1+\Delta)(\epsilon, \theta, \theta) + \\ (1-\Delta)(\beta, \delta, \delta) + (1+\Delta)(\theta, \theta, \epsilon) + (1+\Delta)(\theta, \epsilon, \theta) + (1-\Delta)(\omega, \omega, \omega)\}. \end{aligned} \quad (14)$$

It is apparent from above that the resulting payoff relations (14) in the quantum game cannot be put into a form that is same as for the classical mixed-strategy game i.e. Eq. (7). This raises the question whether there exist constraints that can be placed on the players' directional choices, i.e. the unit vectors  $\vec{a}$ ,  $\vec{b}$ , and  $\vec{c}$ , such that the payoff relations (13) in the quantum game are reduced to the players' payoffs in the classical game allowing mixed strategies (7). In order to find an answer to this we set

$$\Pi_{A,B,C}(\vec{a}, \vec{b}, \vec{c}) = \Pi_{A,B,C}(x, y, z), \quad (15)$$

and equate the right sides of Eqs. (13, 7) i.e.

$$\frac{1}{8}[1 + a_3b_3 + a_3c_3 + b_3c_3 + \Delta] = xyz, \quad (16)$$

$$\frac{1}{8}[1 - a_3b_3 + a_3c_3 - b_3c_3 - \Delta] = x(1-y)z, \quad (17)$$

$$\frac{1}{8}[1 + a_3b_3 - a_3c_3 - b_3c_3 - \Delta] = xy(1-z), \quad (18)$$

$$\frac{1}{8}[1 - a_3b_3 - a_3c_3 + b_3c_3 + \Delta] = x(1-y)(1-z), \quad (19)$$

$$\frac{1}{8}[1 - a_3b_3 - a_3c_3 + b_3c_3 - \Delta] = (1-x)yz, \quad (20)$$

$$\frac{1}{8}[1 + a_3b_3 - a_3c_3 - b_3c_3 + \Delta] = (1-x)(1-y)z, \quad (21)$$

$$\frac{1}{8}[1 - a_3b_3 + a_3c_3 - b_3c_3 + \Delta] = (1-x)y(1-z), \quad (22)$$

$$\frac{1}{8}[1 + a_3b_3 + a_3c_3 + b_3c_3 - \Delta] = (1-x)(1-y)(1-z). \quad (23)$$

Now, by adding Eqs. (16) and (17) we obtain

$$\frac{1}{4}(1 + a_3c_3) = xz, \quad (24)$$

adding Eqs. (16) and (18) gives

$$\frac{1}{4}(1 + a_3b_3) = xy, \quad (25)$$

adding Eqs. (16) and (20) gives

$$\frac{1}{4}(1 + b_3c_3) = yz, \quad (26)$$

adding Eqs. (18) and (19) gives

$$\frac{1}{4}(1 - a_3c_3) = x(1-z). \quad (27)$$

Now, we add Eqs. (24) and (27) to obtain  $x = \frac{1}{2}$ . Adding Eqs. (22) and (23) gives

$$\frac{1}{4}(1 + a_3c_3) = (1-x)(1-z), \quad (28)$$

and substitution from Eq. (24) and  $x = \frac{1}{2}$  gives  $z = \frac{1}{2}$ . Similarly, adding Eqs. (17) and (19) gives

$$\frac{1}{4}(1 - a_3 b_3) = x(1 - y), \quad (29)$$

and adding Eqs. (21) and (23) gives

$$\frac{1}{4}(1 + a_3 b_3) = (1 - x)(1 - y). \quad (30)$$

By adding Eqs. (29) and (30) we obtain  $y = \frac{1}{2}$  and thus  $(x, y, z) = (\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$  is obtained as the solution of the Eqs. (16) to (23).

Therefore, the mixed-strategy payoff relations (7) can be recovered from the payoffs relations (13) for the quantum game only for the special case when  $(x, y, z) = (\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$ . This is because the quantum probability distribution for the GHZ state, from which the payoff relations (13) are constructed, are inherently non-factorizable. In the research area of quantum games, recovering the mixed strategy classical payoff relations from the payoff relations for a quantum game is quite often considered an essential requirement. When the underlying quantum probabilities in a quantum game are obtained from the GHZ state, this requirement is not satisfied except for a very special case, i.e.  $(x, y, z) = (\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$ .

Considering the payoff relations (13) in the quantum game, a Nash equilibrium (NE) is a directional triple  $(\vec{a}^*, \vec{b}^*, \vec{c}^*)$  that satisfies the following constraints:

$$\begin{aligned} \Pi_A(\vec{a}^*, \vec{b}^*, \vec{c}^*) - \Pi_A(\vec{a}, \vec{b}^*, \vec{c}^*) &\geq 0, \\ \Pi_B(\vec{a}^*, \vec{b}^*, \vec{c}^*) - \Pi_B(\vec{a}^*, \vec{b}, \vec{c}^*) &\geq 0, \\ \Pi_C(\vec{a}^*, \vec{b}^*, \vec{c}^*) - \Pi_C(\vec{a}^*, \vec{b}^*, \vec{c}) &\geq 0, \end{aligned} \quad (31)$$

for all  $\vec{a}$ ,  $\vec{b}$ , and  $\vec{c}$ . For the symmetric game, these Nash inequalities take the form

$$\begin{aligned} \Pi_A(\vec{a}^*, \vec{b}^*, \vec{c}^*) - \Pi_A(\vec{a}, \vec{b}^*, \vec{c}^*) &= \frac{1}{8} [(a_3^* - a_3)\Delta_1\gamma_1 + \gamma_2(a_1^* - a_1)\Delta_2 - \gamma_2(a_2^* - a_2)\Delta_3] \geq 0, \\ \Pi_A(\vec{a}^*, \vec{b}^*, \vec{c}^*) - \Pi_A(\vec{a}^*, \vec{b}, \vec{c}^*) &= \frac{1}{8} [(b_3^* - b_3)\Delta'_1\gamma_1 - \gamma_2(b_2^* - b_2)\Delta'_2 + \gamma_2(b_1^* - b_1)\Delta'_3] \geq 0, \\ \Pi_A(\vec{a}^*, \vec{b}^*, \vec{c}^*) - \Pi_A(\vec{a}^*, \vec{b}^*, \vec{c}) &= \frac{1}{8} [(c_3^* - c_3)\Delta''_1\gamma_1 - \gamma_2(c_2^* - c_2)\Delta''_2 + \gamma_2(c_1^* - c_1)\Delta''_3] \geq 0, \end{aligned} \quad (32)$$

where

$$\gamma_1 = \alpha - \beta - \epsilon + \omega \text{ and } \gamma_2 = \alpha - 2\delta - \beta + \epsilon + 2\theta - \omega, \quad (33)$$

and

$$\begin{aligned} \Delta_1 &= b_3 + c_3, \quad \Delta_2 = b_1 c_1 - b_2 c_2, \quad \Delta_3 = b_1 c_2 + b_2 c_1, \\ \Delta'_1 &= a_3 + c_3, \quad \Delta'_2 = a_1 c_2 + a_2 c_1, \quad \Delta'_3 = a_1 c_1 - a_2 c_2, \\ \Delta''_1 &= a_3 + b_3, \quad \Delta''_2 = a_1 b_2 + a_2 b_1, \quad \Delta''_3 = a_1 b_1 - a_2 b_2. \end{aligned} \quad (34)$$

### 3.2 Three-player Prisoners' Dilemma

Prisoner's Dilemma (PD) is a noncooperative game [5–7] that is widely known in the areas of economics, social, and political sciences. In recent years, quantum physics has been added to this list. It was investigated early in the history of quantum games and provided significant motivation for further work in this area.

Two-player PD is about two suspects, considered here as the players in a game, who have been arrested on the allegations of having committed a crime but there not not enough available

evidence to convict them. The investigators come up with an ingenious plan to make the suspects confess their crime.

They are taken to separate cells and are not allowed to communicate. They are contacted individually and, along with being dictated a set of rules, are asked to choose between two choices (strategies): *to Confess* ( $\mathfrak{D}$ ) and *Not to Confess* ( $\mathfrak{C}$ ), where  $\mathfrak{C}$  and  $\mathfrak{D}$  stand for Cooperation and Defection. These are the well-known wordings for the available choices for them and refer to the choice they make to the fellow prisoner, and not to the authorities.

The rules state that if neither prisoner confesses, i.e. ( $\mathfrak{C}, \mathfrak{C}$ ), both are given freedom; when one prisoner confesses ( $\mathfrak{D}$ ) and the other does not ( $\mathfrak{C}$ ), i.e. ( $\mathfrak{C}, \mathfrak{D}$ ) or ( $\mathfrak{D}, \mathfrak{C}$ ), the prisoner who confesses ( $\mathfrak{D}$ ) gets freedom as well as a financial reward, while the prisoner who did not confess ends up in prison for a longer term. If both prisoners confess, i.e. ( $\mathfrak{D}, \mathfrak{D}$ ), both are given a reduced term.

In the two-player case, involving the players  $A$  and  $B$  the strategy pair ( $\mathfrak{D}, \mathfrak{D}$ ) comes out as the unique NE (and the rational outcome) of the game, leading to the situation of both ending up in jail with reduced term. The game offers a dilemma as the rational outcome ( $\mathfrak{D}, \mathfrak{D}$ ) differs from the outcome ( $\mathfrak{C}, \mathfrak{C}$ ), which is an available choice, and for which both prisoners obtain freedom.

With the above notation, the three-player PD can be defined by making the following associations

$$S_1 \sim \mathfrak{C}, S_2 \sim \mathfrak{D}, S'_1 \sim \mathfrak{C}, S'_2 \sim \mathfrak{D}, S''_1 \sim \mathfrak{C}, S''_2 \sim \mathfrak{D}, \quad (35)$$

and afterwards imposing the following conditions [22]:

a) The strategy  $S_2$  is a dominant choice [6] for each player. For Alice this requires

$$\begin{aligned} \Pi_A(S_2, S'_1, S''_1) &> \Pi_A(S_1, S'_1, S''_1), \\ \Pi_A(S_2, S'_2, S''_2) &> \Pi_A(S_1, S'_2, S''_2), \\ \Pi_A(S_2, S'_1, S''_2) &> \Pi_A(S_1, S'_1, S''_2), \end{aligned} \quad (36)$$

and similar inequalities hold for players Bob and Chris.

b) A player is better off if more of his/her opponents choose to cooperate. For Alice this requires

$$\begin{aligned} \Pi_A(S_2, S'_1, S''_1) &> \Pi_A(S_2, S'_1, S''_2) > \Pi_A(S_2, S'_2, S''_2), \\ \Pi_A(S_1, S'_1, S''_1) &> \Pi_A(S_1, S'_1, S''_2) > \Pi_A(S_1, S'_2, S''_2), \end{aligned} \quad (37)$$

and similar relations hold for Bob and Chris.

c) If one player's choice is fixed, the other two players are left in the situation of a two-player PD. For Alice this requires

$$\begin{aligned} \Pi_A(S_1, S'_1, S''_2) &> \Pi_A(S_2, S'_2, S''_2), \\ \Pi_A(S_1, S'_1, S''_1) &> \Pi_A(S_2, S'_1, S''_2), \\ \Pi_A(S_1, S'_1, S''_2) &> (1/2) \{ \Pi_A(S_1, S'_2, S''_2) + \Pi_A(S_2, S'_1, S''_2) \}, \\ \Pi_A(S_1, S'_1, S''_1) &> (1/2) \{ \Pi_A(S_1, S'_1, S''_2) + \Pi_A(S_2, S'_1, S''_1) \}, \end{aligned} \quad (38)$$

and similar relations hold for Bob and Chris.

Translating the above conditions while using the notation introduced in (6) requires

$$\begin{aligned} \text{a) } \beta &> \alpha, \quad \omega > \epsilon, \quad \theta > \delta, \\ \text{b) } \beta &> \theta > \omega, \quad \alpha > \delta > \epsilon, \\ \text{c) } \delta &> \omega, \quad \alpha > \theta, \quad \delta > (1/2)(\epsilon + \theta), \quad \alpha > (1/2)(\delta + \beta), \end{aligned} \quad (39)$$

which defines the generalized three-player PD. For example [22], by letting

$$\alpha = 7, \quad \beta = 9, \quad \delta = 3, \quad \epsilon = 0, \quad \omega = 1, \quad \theta = 5, \quad (40)$$

all of these conditions hold.

## 4 Three-player quantum Prisoners' Dilemma with GHZ state

The values in (40) give  $\gamma_1 = -1$  and  $\gamma_2 = 1$ . With the deltas given in (34), the Nash inequalities (32) take the form

$$\begin{aligned}
& \Pi_A(\vec{a}^*, \vec{b}^*, \vec{c}^*) - \Pi_A(\vec{a}, \vec{b}^*, \vec{c}^*) = \\
& \frac{1}{8} [-(a_3^* - a_3)(b_3 + c_3) + (a_1^* - a_1)(b_1c_1 - b_2c_2) - (a_2^* - a_2)(b_1c_2 + b_2c_1)] \geq 0, \\
& \Pi_A(\vec{a}^*, \vec{b}^*, \vec{c}^*) - \Pi_A(\vec{a}^*, \vec{b}, \vec{c}^*) = \\
& \frac{1}{8} [-(b_3^* - b_3)(a_3 + c_3) - (b_2^* - b_2)(a_1c_2 + a_2c_1) + (b_1^* - b_1)(a_1c_1 - a_2c_2)] \geq 0, \\
& \Pi_A(\vec{a}^*, \vec{b}^*, \vec{c}^*) - \Pi_A(\vec{a}^*, \vec{b}^*, \vec{c}) = \\
& \frac{1}{8} [-(c_3^* - c_3)(a_3 + b_3) - (c_2^* - c_2)(a_1b_2 + a_2b_1) + (c_1^* - c_1)(a_1b_1 - a_2b_2)] \geq 0.
\end{aligned} \tag{41}$$

These inequalities show that for the PD game defined in (40), no directional triplet can exist as a NE when the three players have the choice to direct their respective unit vector along any direction i.e. there are no restrictions placed on the players' directional choices.

The inequalities (32) suggest the following cases:

### 4.1 Case (a)

Consider  $a_3 = b_3 = c_3 = 0$ . Nash inequalities (32) then take the form

$$\begin{aligned}
\Pi_A(\vec{a}^*, \vec{b}^*, \vec{c}^*) - \Pi_A(\vec{a}, \vec{b}^*, \vec{c}^*) &= \frac{1}{8}\gamma_2 [(a_1^* - a_1)\Delta_2 - (a_2^* - a_2)\Delta_3] \geq 0, \\
\Pi_A(\vec{a}^*, \vec{b}^*, \vec{c}^*) - \Pi_A(\vec{a}^*, \vec{b}, \vec{c}^*) &= \frac{1}{8}\gamma_2 [-(b_2^* - b_2)\Delta'_2 + (b_1^* - b_1)\Delta'_3] \geq 0, \\
\Pi_A(\vec{a}^*, \vec{b}^*, \vec{c}^*) - \Pi_A(\vec{a}^*, \vec{b}^*, \vec{c}) &= \frac{1}{8}\gamma_2 [-(c_2^* - c_2)\Delta''_2 + (c_1^* - c_1)\Delta''_3] \geq 0,
\end{aligned} \tag{42}$$

that can also be expressed as

$$\begin{aligned}
\Pi_A(\vec{a}^*, \vec{b}^*, \vec{c}^*) - \Pi_A(\vec{a}, \vec{b}^*, \vec{c}^*) &= \frac{1}{8}\gamma_2 [a_1^*\Delta_2 - a_2^*\Delta_3 + \varsigma] \geq 0, \\
\Pi_A(\vec{a}^*, \vec{b}^*, \vec{c}^*) - \Pi_A(\vec{a}^*, \vec{b}, \vec{c}^*) &= \frac{1}{8}\gamma_2 [-b_2^*\Delta'_2 + b_1^*\Delta'_3 + \varsigma] \geq 0, \\
\Pi_A(\vec{a}^*, \vec{b}^*, \vec{c}^*) - \Pi_A(\vec{a}^*, \vec{b}^*, \vec{c}) &= \frac{1}{8}\gamma_2 [-c_2^*\Delta''_2 + c_1^*\Delta''_3 + \varsigma] \geq 0,
\end{aligned} \tag{43}$$

where

$$\varsigma = a_1b_2c_2 + a_2b_1c_2 - a_1b_1c_1 + a_2b_2c_1. \tag{44}$$

Consider the case when  $\gamma_2 > 0$ , then for given  $a_1^*$ ,  $b_1^*$ , and  $c_1^*$ , the restrictions on the directions that the unit vectors  $\vec{a}$ ,  $\vec{b}$ , and  $\vec{c}$  can take can be determined. For instance, for  $a_2^* = b_1^* = c_1^* = 1$ , i.e. then  $a_2^* = b_2^* = c_2^* = 0$ , these constraints become

$$\Delta_2 + \varsigma \geq 0, \quad \Delta'_3 + \varsigma \geq 0, \quad \Delta''_3 + \varsigma \geq 0. \tag{45}$$

## 4.2 Case (b)

Consider  $\gamma_2 = 0$  and  $a_3 = b_3 = c_3 = 0$ . With these constraints, the allowed directions are confined to the X-Y plane and any directional triplet then exists as a NE. In this case, from Eqs. (34) we then have  $\Delta_1 = \Delta'_1 = \Delta''_1 = 0$ . As  $\vec{a}$ ,  $\vec{b}$ , and  $\vec{c}$  are unit vectors, we also have  $a_2 = \pm\sqrt{1 - a_1^2}$ ,  $b_2 = \pm\sqrt{1 - b_1^2}$ , and  $c_2 = \pm\sqrt{1 - c_1^2}$ .

## 5 Discussion

We present an analysis of the three-partite EPR experiments that use a GHZ state and is considered as a three-player non-cooperative quantum game. The players' strategic choices are the three directions  $\vec{a}$ ,  $\vec{b}$ , and  $\vec{c}$  along which the dichotomic observables  $\vec{n} \cdot \vec{\sigma}$  are measured, where  $\vec{n} = \vec{a}, \vec{b}, \vec{c}$  and  $\vec{\sigma}$  is a vector whose components are the standard Pauli matrices. Using Kaszlikowski and Żukowski's results [20] for the quantum probabilities involved in such experiments, we develop a three-player quantum game, with the underlying three-partite EPR experiment. This extends an approach to quantum games by Vaidman [19] that does not involve Hilbert space, and/or quantum measurement, and shows how three-player quantum games with EPR experiments can be developed. Players' strategies are their directions in terms of which their payoffs are expressed using Eq. (10). Nash inequalities are used to obtain Nash equilibria as direction triples and the players' payoffs are then compared to their payoffs for the Nash equilibria in the classical game.

For a three-player Prisoners' Dilemma game, defined in (40), we conclude that no directional triplet can exist as a NE when no restrictions are placed on the players' directional choices. A directional triplet, however, can exist as a NE under constraints placed on the directions allowed to the players. This is in accordance with Eisert et al.'s result in Ref. [1] showing that a pair of unitary transformations  $(\hat{Q}, \hat{Q})$ , where  $\hat{Q} \sim \hat{U}(0, \pi/2)$ , exists as a NE in PD when the players' allowed actions are restricted to certain subsets of the set SU(2) consisting of all unitary transformations.

As is known [23, 24] that the particular subset of unitary transformations that Eisert et al. used in order to obtain the NE of  $(\hat{Q}, \hat{Q})$  in two-player quantum Prisoners' Dilemma is not even closed under composition. In particular, in Eisert et al.'s protocol for  $2 \times 2$  quantum games [1], the new Nash equilibria, and the classical-quantum transitions that occur, are the outcomes of the particular strategy space chosen that is a two-parameter subset of single qubit unitary operators. By choosing a different, but equally plausible, two-parameter strategy a different Nash equilibria with different classical-quantum transitions can arise.

Using an EPR setting, and a shared GHZ state, for a three-player quantum Prisoners' Dilemma game, we present an approach that is driven along purely probabilistic lines with only an implicit reference to the mathematical formalism of quantum theory, and showing the constraints on the players' directional choices under which a particular triplet can exist as a NE in the game.

## References

- [1] Eisert, J., Wilkens, M., and Lewenstein, M., Quantum games and quantum strategies, *Phy. Rev. Lett.* **83**, 3077 (1999).
- [2] Eisert, J., Wilkens, M., Quantum games, *J. Mod. Opt.* **47**, 2543 (2000).
- [3] The URL <https://scholar.google.com.au/citations?user=wkfPcaQAAAAJ&hl=en> contains an extensive list of publications on the subject of quantum games.
- [4] Peres, A., *Quantum Theory: Concepts and Methods*, Kluwer Academic Publishers (1995).
- [5] Binmore, K., *Game Theory: A Very Short Introduction*, Oxford University Press, USA, (2007).
- [6] Rasmusen, E., *Games & Information: An Introduction to Game Theory*, Blackwell Publishers Ltd., Oxford, (2001) 3rd Edition.

- [7] Osborne, M. J., *An Introduction to Game Theory*, Oxford University Press, USA (2003).
- [8] Iqbal, A., Weigert, S., Quantum correlation games, *J. Phys. A: Math. & Gen.* **37**, 5873-5885 (2004).
- [9] Iqbal, A., Chappell, J.M., Li, Q., Pearce, Charles E.M., and Abbott, D., A probabilistic approach to quantum Bayesian games of incomplete information, *Quant. Inform. Process.* **13**, 2783-2800 (2014).
- [10] Iqbal, A., Chappell, J.M., and Abbott, D., Social optimality in quantum Bayesian games, *Physica A: Statistical Mechanics and its Applications* **436**, 798-805 (2015).
- [11] Iqbal, A., Abbott, D., Constructing quantum games from a system of Bell's inequalities, *Physics Letters A* **374**, 3155-3163 (2010).
- [12] Chappell, J.M., Iqbal, A., and Abbott, D., Constructing quantum games from symmetric non-factorizable joint probabilities, *Physics Letters A* **374**, 4104-4111 (2010).
- [13] Iqbal, A., Abbott, D., Quantum matching pennies game, *Journal of the Physical Society of Japan* **78**, 014803 (2009).
- [14] Bell, J. On the Einstein-Podolsky-Rosen paradox. *Physics* **1**, 195–200 (1964).
- [15] Bell, J. *Speakable and Unspeakable in Quantum Mechanics*, Cambridge University Press (1987).
- [16] Bell, J. On the problem of hidden variables in quantum mechanics, *Reviews of Modern Physics* **38**, 447–452 (1966).
- [17] Aspect, A., Dalibard, J., Roger, G. Experimental test of Bell's inequalities using time-varying analyzers. *Phys. Rev. Lett.* **49**, 1804–1807 (1982).
- [18] Clauser, J.F., Horne, M.A., Shimony, A., Holt, R.A., Proposed experiment to test local hidden-variable theories, *Phys. Rev. Lett.* **23**, 880–884 (1969).
- [19] Vaidman, L., Variations on the theme of the Greenberger-Horne-Zeilinger proof, *Found. Phys.* **29**, 615-630 (1999).
- [20] D. Kaszlikowski and M. Żukowski, Three qubit GHZ correlations and generalized Bell experiments, *International Journal of Theoretical Physics* **42**/5, pp 1023-1035 (2003).
- [21] A. Iqbal and A. H. Toor, Darwinism in quantum systems?, *Phys. Lett. A* **294**/5-6, 261 (2002).
- [22] M. J. O'Donnell, The three-player Prisoner's Dilemma, Available from the URL: <http://www.classes.cs.uchicago.edu/archive/1998/fall/CS105/Project/node6.html>, Department of Computer Science, The University of Chicago (1998).
- [23] Benjamin S.C., and Hayden, Patrick M., Comment on “Quantum Games and Quantum Strategies”, *Phys. Rev. Lett.* **87**, 069801 (2001).
- [24] Flitney, A. P., Hollenberg, Lloyd, C. L., Nash equilibria in quantum games with generalized two-parameter strategies, *Phys. Lett. A* **363**, Issues 5–6, 381-388 (2007).