



# **Hypoxia-Inducible Factor-1 $\alpha$ mRNA Contains An Internal Ribosome Entry Site That Allows Efficient Translation During Normoxia And Hypoxia**

**Kenneth Lang, B.Sc., B Health Sc. (Hons)**

A thesis submitted to the University of Adelaide  
for the degree of Doctor of Philosophy



Division of Human Immunology  
Hanson Centre for Cancer Research  
Through the Department of Medicine  
University of Adelaide  
Adelaide, South Australia

March, 2003

# Table of Contents

Table of Contents.....	
Summary.....	
Statement of Originality.....	
Acknowledgements.....	
Publications arising from this thesis.....	
Abbreviations.....	

## Chapter 1

<b>Introduction.....</b>	<b>1</b>
1.1 Expression and Regulation of genes.....	1
1.1.1 Transcriptional regulation of gene expression.....	1
1.1.2 Post-transcriptional mechanisms regulating gene expression.....	2
1.1.2.1 Nonsense-mediated mRNA decay.....	3
1.1.2.2 mRNA splicing.....	4
1.1.2.3 mRNA stability.....	4
1.1.2.4 Translation.....	5
1.2 Eukaryotic Translation Initiation.....	5
1.2.1 Modification of messenger RNA's.....	6
1.2.2 Cap-dependent initiation.....	6
1.2.3 Formation of the 43S pre-initiation complex.....	7
1.2.4 Binding of mRNA to ribosomes.....	7
1.2.5 Migration of the 43S pre-initiation complex to the initiation codon.....	7
1.2.6 Elongation and Termination.....	8
1.2.7 Regulation of translation initiation.....	8
1.2.7.1 Poly(A) tail.....	9
1.2.7.2 Regulation of initiation factor activity by phosphorylation.....	9
1.2.7.2.1 eIF2.....	9
1.2.7.2.2 eIF4E.....	10
1.2.7.2.3 4E-BP.....	11
1.2.7.2.4 eIF4G.....	12
1.2.7.3 Structural features regulating translation.....	13
1.2.8 Variations of the cap-dependent scanning model.....	14
1.2.8.1 Leaky Initiation.....	14
1.2.8.2 Re-Initiation.....	15
1.2.8.3 Ribosome shunting.....	15
1.2.9 Internal Ribosome Entry.....	16
1.2.10 Cellular IRES elements.....	17
1.2.11 Vascular Endothelial Growth Factor IRES.....	18
1.2.12 c-myc IRES.....	21

1.3 Angiogenesis.....	22
1.3.1 Angiogenesis and cancer.....	23
1.3.2 Role of VEGF in angiogenesis.....	23
1.3.3 Activation of VEGF by hypoxia.....	24
1.4 Hypoxia-Inducible Factor.....	25
1.4.1 HIF-1 target genes.....	26
1.4.2 HIF-1 $\beta$ / ARNT.....	27
1.4.3 HIF-1 $\alpha$ mRNA expression.....	28
1.4.4 The HIF-1 $\alpha$ protein.....	28
1.4.5 Dimerisation and DNA binding domains.....	29
1.4.6 Protein stability and degradation.....	29
1.4.7 Degradation of HIF-1 $\alpha$ is mediated by VHL.....	30
1.4.8 Proline hydroxylation determines HIF-1 $\alpha$ protein stability.....	31
1.4.9 Transactivation properties of HIF-1 $\alpha$ .....	33
1.4.10 Asparagine hydroxylation regulates HIF-1 $\alpha$ transactivation potential.....	33
1.4.11 Activation of HIF-1 $\alpha$ by other mechanisms.....	35
1.4.11.1 ERK / MAPK pathway.....	35
1.4.11.2 p53.....	35
1.4.11.3 p14 <sup>ARF</sup> Tumour Suppressor.....	36
1.4.11.4 PI3 Kinase signaling pathway.....	37
1.4.12 Role of HIF-1 $\alpha$ in development.....	38
1.4.13 Role of HIF-1 in disease.....	39
1.4.14 Role of HIF-1 in cancer.....	39
1.5 Project Rationale.....	42

## **Chapter 2**

<b>Materials and Methods.....</b>	<b>44</b>
2.1 Chemicals, reagents and consumables.....	44
2.2 Enzymes and protein molecular weight markers.....	44
2.3 Oligonucleotides.....	45
2.4 Radiochemicals.....	46
2.5 Buffers and Solutions.....	46
2.6 <i>E. coli</i> Bacterial strains.....	50
2.7 Plasmids.....	50
2.8 Eukaryotic cell lines.....	50
2.9 Miniprep purification of plasmid DNA (Alkaline lysis method).....	51
2.10 Large scale preparation of plasmid DNA.....	51
2.11 Purification of DNA fragments using GeneClean kit.....	53
2.12 DNA ligation.....	53
2.13 Preparation of competent bacteria for calcium chloride mediated transformation.....	53
2.14 Transformation of bacterial competent cells.....	54
2.15 Sequencing.....	54
2.16 Polymerase chain reaction.....	54
2.17 Transient transfection of NIH3T3, HeLa, HEK293 and MCF7 cells.....	55
2.18 Stimulation of fos promoter in pFGH.....	55
2.19 Secreted alkaline phosphatase assay.....	56
2.20 Dual luciferase assay.....	57
2.21 <i>in vitro</i> transcription to generate RNA probes.....	57
2.21.1 Preparation of plasmid templates.....	57
2.21.2 <i>in vitro</i> transcription.....	58

2.21.3 Purification of the RNA probes by PAGE.....	58
2.21.4 Elution of the RNA probes from polyacrylamide.....	59
2.21.5 Hybridising the probe with RNA.....	59
2.22 RNase protection assay.....	60
2.22.1 Digestion with RNase A.....	60
2.22.2 Electrophoresis of RNase A protection samples.....	60
2.22.3 Quantitation and analysis of specific mRNAs.....	61
2.23 Northern blot assay using radio-labelled DNA probe.....	61
2.23.1 Northern transfer of RNA from agarose gel to nylon membrane by downward capillary transfer.....	61
2.23.2 Hybridisation.....	62
2.23.3 Labelling of DNA probes.....	62
2.23.4 Washing of nylon filters for visualisation of bands.....	62
2.24 <sup>35</sup> S-methionine incorporation.....	62
2.25 Polysome analysis.....	63
2.26 Western blot.....	64

### **Chapter 3**

#### **Hypoxia-Inducible Factor-1 $\alpha$ mRNA contains an Internal Ribosome Entry site.....65**

3.1 Introduction.....	65
3.2 Construction of HIF-1 $\alpha$ reporter plasmids for analysis in mammalian cells.....	66
3.3 Characteristics of the HIF-1 $\alpha$ 5'UTR.....	69
3.4 System to determine the function of the HIF-1 $\alpha$ 5'UTR.....	70
3.5 The HIF-1 $\alpha$ 5'UTR allows efficient translation.....	70
3.6 The HIF-1 $\alpha$ 5'UTR contains an Internal Ribosome Entry Site.....	71
3.7 Relative efficiency of the HIF-1 $\alpha$ IRES compared with other cellular IRESs.....	73
3.8 Mapping of the HIF-1 $\alpha$ IRES.....	74
3.9 Summary and Discussion.....	79

### **Chapter 4**

#### **Hypoxia-Inducible Factor -1 $\alpha$ mRNA is spared the general reduction of translation under hypoxia, which is mediated by the internal ribosome entry site.....83**

4.1 Introduction.....	83
4.2 Total translation is reduced during hypoxia but translation of HIF-1 $\alpha$ mRNA is unaffected.....	84
4.3 HIF-1 $\alpha$ IRES activity during hypoxia and serum deprivation.....	85
4.4 The effect of hypoxia and serum deprivation on the VEGF and c-myc IRES.....	86
4.5 Summary and Discussion.....	88

### **Chapter 5**

#### **Activation of the phosphatidylinositol 3-kinase signaling pathway increases hypoxia-inducible factor-1 $\alpha$ internal ribosome entry site activity.....91**

5.1 Introduction.....	91
5.2 Overexpression Akt constructs.....	92
5.3 Heregulin stimulation increases HIF-1 $\alpha$ protein, but not mRNA levels.....	93
5.4 Cap-dependent translation and HIF-1 $\alpha$ IRES activity are concomitantly up-regulated in response to heregulin stimulation.....	94

5.5 Stimulation of serum-starved MCF7 cells with heregulin does not alter firefly luciferase mRNA levels.....	96
5.6 The PI3K signaling pathway regulates HIF-1 $\alpha$ IRES activity.....	97
5.7 Summary and Discussion.....	99

## **Chapter 6**

<b>Long G + C rich 5'UTRs containing internal ribosome entry sites may be more susceptible to mRNA instability.....</b>	<b>103</b>
6.1 Introduction.....	103
6.2 Constructs of reporter plasmids to determine IRES activity versus mRNA instability.....	104
6.3 VEGF 5'UTR mRNA destabilisation correlates with IRES activity.....	106
6.4 The XIAP IRES cannot destabilise fGH mRNA.....	109
6.5 G + C rich 5'UTRs containing IRES elements may also contain mRNA destabilisation elements.....	110
6.6 Summary and Discussion.....	111

## **Chapter 7**

<b>Final Discussion.....</b>	<b>115</b>
7.1 Final Discussion.....	115

## **Chapter 8**

<b>References.....</b>	<b>123</b>
------------------------	------------

## Summary

HIF-1 $\alpha$  is the regulated subunit of the HIF-1 transcription factor, which induces transcription of a number of genes involved in the cellular response to hypoxia. The HIF-1 $\alpha$  protein is rapidly degraded in cells supplied with adequate oxygen, but is stabilised in hypoxic cells. Using polysome profile analysis, I found that translation of HIF-1 $\alpha$  mRNA in NIH3T3 cells is spared the general reduction in translation rate that occurs during hypoxia. To assess whether the 5'UTR of the HIF-1 $\alpha$  mRNA contains an internal ribosome entry site (IRES), I constructed a dicistronic reporter with the HIF-1 $\alpha$  5'UTR inserted between the two reporter coding regions. I found the HIF-1 $\alpha$  5'UTR promoted translation of the downstream reporter, indicating the presence of an IRES. IRES activity was not affected by hypoxic conditions that caused a reduction in cap-dependent translation, and IRES activity was less affected by serum-starvation than was cap-dependent translation. This indicates that the presence of the IRES in the HIF-1 $\alpha$  5'UTR is to allow translation to be maintained under conditions that are inhibitory to cap-dependent translation.

Several reports have implicated the 5'UTR to be important for HIF-1 $\alpha$  protein synthesis upon activation of the phosphatidylinositol 3-kinase (PI3K) signal transduction pathway. I hypothesised that this specific induction of HIF-1 $\alpha$  protein synthesis is mediated by the HIF-1 $\alpha$  IRES. I found HIF-1 $\alpha$  IRES activity was increased in response to activation of the PI3K signaling pathway, with a concomitant increase in cap-dependent translation. Results also indicate that activation of the IRES is independent of Akt (a downstream target of PI3K), implicating other signaling pathways in the regulation of HIF-1 $\alpha$  IRES function.

I also demonstrate that GC rich 5'UTRs containing an IRES may stimulate mRNA destabilisation. This mode of regulation was found in three IRESs tested (vascular endothelial growth factor, HIF-1 $\alpha$  and c-myc), but is not a mechanism of all cellular IRESs, as the X-linked inhibitor of apoptosis 5'UTR could not destabilise a reporter

mRNA. Data presented in this thesis indicates that the HIF-1 $\alpha$  5'UTR is subject to multiple modes of regulation, including translation by an IRES, and this is regulated by signal transduction pathways, and mRNA instability.

## **Statement of Originality**

This thesis contains no material that has been accepted for the award of any degree or diploma by any other university. To the best of my knowledge it contains no material that has previously been published by any other person, except where due reference has been made in the text. I consent to this thesis, when deposited in the university library, being available for photocopying and loan.

**Kenneth Lang**

## Acknowledgements

This epic journey would have gone nowhere without the help and assistance of a number of people. I must firstly thank Greg Goodall, whose guidance and knowledge was invaluable. I also thank Greg for allowing me "to do my own thing", and for graciously reading numerous versions of thesis chapters. Thankyou for using red pen somewhat sparingly. A big thanks goes to the people who provided me with plasmids to undertake some of this work. To Anne Willis for providing the dual luciferase reporters and the c-myc 5'UTR, Andreas Kappel, who cloned the HIF-1 $\alpha$  5'UTR; and Martin Holcik for the XIAP 5'UTRs. To Mark Guthridge and Angel Lopez for donating their time in an attempt to get me to understand the PI3K signalling pathway, as well as advice and reagents.

I also need to extend my thanks to past and present members of the Goodall Lab. To Justin Dibbens for much advice and for making the VEGF 5'UTR deletions in fGH and performing the instability assays. John Harvey... what can I say? At least I know for sure that flatulence really is funny! To Steven Polyak for a continual supply of "humourous" e-mails as well as some experimental advice. To the current members, Leeanne Coles, Rosemary Sladic, Antonetta Bartley, Andrew Bert, Rod Putland, Cathy Lagnado and Lidia Lambrusco for endless helpful comments and advice. I'm sure most of the controls would have been left out if not for you guys! Last, but certainly not least, to my "evil" bay companion, Peter Diamond, who provided me with entertainment on a daily basis. Seriously though, thanks for the scientific advice, and for the listening ear on occasion. To the lab in general, thanks for listening to my zany ideas, as well as some of my not so politically correct and line-crossing ones.

An extended thanks to the division of human immunology for interesting chats in the tearoom, and for providing a friendly place to work.

Of course, none of this would have been possible without the help of three of the most important people in my life. To my wife, Tamara, who not only gave invaluable financial support, placed beliefs in me that I was doing the right thing in going on to do a PhD, and that I could actually accomplish it. And to my two daughters, Caitlyn and Jacinta, who provided an endless supply of entertainment (and sometimes annoyance) particularly during the writing period. They certainly provided me with the will to keep going, even if it killed me (and it nearly did, or was that my wife that did?). A big thanks to Mum and Dad for continual belief and support throughout my university life. I would love to mention everyone, but apparently the acknowledgements are not meant to be longer than the actual thesis, so in that case, a big thanks goes to all of my family and friends (you all know who you are) for just being there. You really did help you just probably don't know it.

Finally, I must thank coca-cola, especially for putting coke on special almost every week during the writing period. I'm so glad you put caffeine in it.

## Publications arising from work presented in this thesis

Lang, K.J.D., Goodall, G.J. (2001) Translation of hypoxia-inducible factor-1 $\alpha$  is mediated by an internal ribosome entry site 13<sup>th</sup> *Lorne Cancer Conference*.

Lang, K.J.D., Kappel, A., Goodall, G.J. (2002) Hypoxia inducible factor-1 $\alpha$  mRNA contains an internal ribosome entry site that allows efficient translation during hypoxia 23<sup>rd</sup> *Lorne Genome Conference on the Organisation and Expression of the Genome*.

Lang, K.J.D., Kappel, A., Goodall, G.J. (2002) Hypoxia inducible factor-1 $\alpha$  mRNA contains an internal ribosome entry site that allows efficient translation during normoxia and hypoxia *Mol Biol Cell* **13**: 1792-1801.

Lang, K.J.D., Goodall, G.J. (2002) HIF-1 $\alpha$  mRNA contains an IRES that allows efficient translation during normoxia and hypoxia *Australian Society for Medical Research*

Lang, K.J.D., Kappel, A., Goodall, G.J. (2002) Hypoxia inducible factor-1 $\alpha$  mRNA contains an internal ribosome entry site that allows efficient translation during normoxia and hypoxia *ComBio 2002*.

Goodall, G.J., Coles, L.S., Bartley, M.A., Lang, K.J.D., (2003) Post-transcriptional regulation of VEGF. *Genetics of Angiogenesis* BIOS Scientific Ltd. Oxford, U.K.

## ABBREVIATIONS

All abbreviations used throughout this thesis are in accordance with those described in the *Journal of Biological Chemistry*; additional and alternate abbreviations are shown below.

APS	ammonium persulphate
Amp	ampicillin
ARE	AU-rich element
ATP	adenosine triphosphate
bp	base pairs
BSA	bovine serum albumin
CHX	cycloheximide
CIP	calf intestinal phosphatase
DEPC	diethyl pyrocarbonate
DMEM	Dulbecco's modified Eagle's Medium
DMSO	dimethylsulphoxide
DNA	deoxyribonucleic acid
DNase	deoxyribonuclease
DTT	dithiothreitol
EDTA	ethylenediaminetetra-acetic acid
eIF	eukaryotic initiation factor
FCS	foetal calf serum
FF	firefly
GAPDH	glyceraldehyde-3-phosphate dehydrogenase
HEPES	N-2-hydroxyethylpiperazine-N'-2-ethane-sulfonic acid
HIF	hypoxia-inducible factor
IRES	Internal ribosome entry site
kDa	kilodalton
mA	milliamps
MEN	Mops EDTA sodium acetate
Met	methionine
mins	minutes
mRNA	messenger RNA
NP-40	Nonident ® P 40
nt	nucleotides
OD	optical density
PAGE	polyacrylamide gel electrophoresis

PBS	phosphate buffered saline
PCR	polymerase chain reaction
PPT	polypyrimidine tract
RL	renilla
RNA	ribonucleic acid
RNase A	ribonuclease A
rpm	revolutions per minute
rRNA	ribosomal RNA
SeAP	Secreted alkaline phosphatase
SDS	sodium dodecyl sulphate
secs	seconds
SEM	standard error of the mean
SSC	Saline sodium citrate
TAE	tris-acetic acid EDTA
TBE	tris-boric acid EDTA
TE	tris-EDTA
TEMED	N,N,N',N'-tetramethyl-ethene-diamine
tRNA	transfer RNA
UTR	untranslated region
UV	ultra-violet
VEGF	vascular endothelial growth factor
VHL	von-Hippel Lindau protein
XIAP	X-linked inhibitor of apoptosis