

ACCEPTED VERSION

Fei Han, Varghese Mathai

T-duality in an H-Flux: exchange of momentum and winding

Communications in Mathematical Physics, 2018; 363(1):333-350

© Springer-Verlag GmbH Germany, part of Springer Nature 2018.

*This is a post-peer-review, pre-copyedit version of an article published in **Communications in Mathematical Physics**. The final authenticated version is available online at:*

<http://dx.doi.org/10.1007/s00220-018-3112-1>

PERMISSIONS

<https://www.springer.com/gp/open-access/publication-policies/self-archiving-policy>

Self-archiving for articles in subscription-based journals

Springer journals' [policy on preprint sharing](#).

By signing the Copyright Transfer Statement you still retain substantial rights, such as self-archiving:

*Author(s) are permitted to self-archive a pre-print and an author's **accepted manuscript** version of their Article.*

.....

b. An Author's Accepted Manuscript (AAM) is the version accepted for publication in a journal following peer review but prior to copyediting and typesetting that can be made available under the following conditions:

(i) Author(s) retain the right to make an AAM of their Article available on their own personal, self-maintained website immediately on acceptance,

(ii) Author(s) retain the right to make an AAM of their Article available for public release on any of the following 12 months after first publication ("Embargo Period"): their employer's internal website; their institutional and/or funder repositories. AAMs may also be deposited in such repositories immediately on acceptance, provided that they are not made publicly available until after the Embargo Period.

An acknowledgement in the following form should be included, together with a link to the published version on the publisher's website: "This is a post-peer-review, pre-copyedit version of an article published in [insert journal title]. The final authenticated version is available online at: [http://dx.doi.org/\[insert DOI\]](http://dx.doi.org/[insert DOI])".

When publishing an article in a subscription journal, without open access, authors sign the Copyright Transfer Statement (CTS) which also details Springer's self-archiving policy.

See Springer Nature [terms of reuse](#) for archived author accepted manuscripts (AAMs) of subscription articles.

26 August 2020

<http://hdl.handle.net/2440/117581>

T-DUALITY IN AN H-FLUX: EXCHANGE OF MOMENTUM AND WINDING

FEI HAN AND VARGHESE MATHAI

ABSTRACT. Using our earlier proposal for Ramond-Ramond fields in an H-flux on loop space [13], we extend the Hori isomorphism in [6, 7] from invariant differential forms, to invariant exotic differential forms such that the *momentum* and *winding numbers* are exchanged, filling in a gap in the literature. We also extend the compatibility of the action of invariant exact Courant algebroids on the T-duality isomorphism in [10], to the T-duality isomorphism on exotic invariant differential forms.

CONTENTS

Introduction	1
1. Exact Courant algebroids and exotic differential forms	5
2. T-duality and exotic Hori formulae	7
2.1. Review of T-duality	7
2.2. Exotic theories	9
2.3. Twisted integration along the fiber	11
2.4. The exotic Hori formulae	12
2.5. The trivial bundles case	17
References	17

INTRODUCTION

In [6, 7], T-duality in a background flux was studied for the first time, and we summarise it here to begin with. Upon compactifying spacetime in one direction to a principal circle bundle $\mathbb{T} \rightarrow Z \xrightarrow{\pi} X$ with background \mathbb{T} -invariant flux H which is a closed 3-form on Z . Then there is a T-dual circle bundle $\hat{\mathbb{T}} \rightarrow \hat{Z} \xrightarrow{\hat{\pi}} X$ with T-dual background $\hat{\mathbb{T}}$ -invariant flux \hat{H} which is a closed 3-form on \hat{Z} such that $c_1(Z) = \hat{\pi}_*[\hat{H}]$ and $c_1(\hat{Z}) = \pi_*[H]$, and the constraint that $[H] = [\hat{H}]$ on the correspondence space $Z \times_X \hat{Z}$ ensures that $[\hat{H}]$ is uniquely defined. The slogan that,

the Chern class is exchanged with the background flux

encapsulates T-duality in a background flux, so there is a change in topology if either the Chern class or the background flux is topologically nontrivial.

2010 *Mathematics Subject Classification*. Primary 55N91, Secondary 58D15, 58A12, 81T30, 55N20.

Choosing \mathbb{T} -invariant connection 1-forms A on Z and $\hat{\mathbb{T}}$ -invariant connection \hat{A} on \hat{Z} respectively, the rules for transforming the RR fields can be encoded in the twisted Fourier-Mukai transform [6, 7]

$$(0.1) \quad T_* G = \int_{\mathbb{T}} G e^{-A \wedge \hat{A}},$$

where $G \in \Omega^\bullet(Z)^\mathbb{T}$ is an invariant differential form on spacetime which is the total Ramond-Ramond field-strength,

$$\begin{aligned} G &\in \Omega^{\bar{k}}(Z)^\mathbb{T} && \text{for Type IIA;} \\ G &\in \Omega^{\bar{k}+1}(Z)^\mathbb{T} && \text{for Type IIB,} \end{aligned}$$

for $\bar{k} = k \bmod 2$, and where the right hand side of equation (0.1) is an invariant differential form on $Z \times_X \hat{Z}$, and the integration is along the \mathbb{T} -fiber of Z .

Define the Riemannian metrics on Z and \hat{Z} by

$$g_Z = \pi^* g_X + R^2 A \odot A, \quad g_{\hat{Z}} = \hat{\pi}^* g_X + R^{-2} \hat{A} \odot \hat{A}.$$

Under the above choices of Riemannian metrics and flux forms, the twisted Fourier-Mukai transform is an isometry

$$(0.2) \quad T: \Omega^{\bar{k}}(Z)^\mathbb{T} \rightarrow \Omega^{\bar{k}+1}(\hat{Z})^{\hat{\mathbb{T}}},$$

for $\bar{k} = k \bmod 2$, inducing isometries on the spaces of twisted harmonic forms.

In particular, R goes to $1/R$, which is an important feature of T-duality, and there is an induced degree-shifting isomorphism between twisted cohomology groups,

$$(0.3) \quad T_* : H^{\bar{k}}(Z, H) \cong H^{\bar{k}+1}(\hat{Z}, \hat{H}).$$

for $\bar{k} = k \bmod 2$, where $H^*(Z, H) := \{\Omega^*(Z)^\mathbb{T}, d + H\}$ and $H^*(\hat{Z}, \hat{H}) := \{\Omega^*(\hat{Z})^{\hat{\mathbb{T}}}, d + \hat{H}\}$. These twisted cohomology groups were first defined in [18] and their relation to D-branes in an H-flux and their charges were further explored in [5, 17].

One of the deficiencies of the T-duality isomorphism in equation (0.2) is that it is only defined on the smaller configuration space of invariant differential forms on spacetime Z , and therefore is not easy to formulate the exchange of momentum and winding in this framework. We rectify this in our paper as follows.

Let $\xi, \hat{\xi}$ denote the complex line bundles associated to the circle bundles Z, \hat{Z} and the standard representation of the circle on complex plane respectively. Define the **exotic differential forms** by

$$\begin{aligned} \mathcal{A}^{\bar{k}}(Z)^\mathbb{T} &= \bigoplus_{n \in \mathbb{Z}} \mathcal{A}_n^{\bar{k}}(Z)^\mathbb{T} := \bigoplus_{n \in \mathbb{Z}} \Omega^{\bar{k}}(Z, \pi^*(\xi^{\otimes n}))^\mathbb{T}, \\ \mathcal{A}^{\bar{k}}(\hat{Z})^{\hat{\mathbb{T}}} &= \bigoplus_{n \in \mathbb{Z}} \mathcal{A}_n^{\bar{k}}(\hat{Z})^{\hat{\mathbb{T}}} := \bigoplus_{n \in \mathbb{Z}} \Omega^{\bar{k}}(\hat{Z}, \hat{\pi}^*(\xi^{\otimes n}))^{\hat{\mathbb{T}}} \end{aligned}$$

for $\bar{k} = k \bmod 2$. for $\bar{k} = k \bmod 2$.

An analogous space of exotic differential forms and equivariantly flat superconnection [16] was first defined on loop space in our earlier paper [13], which was inspired by and generalises

some of the results in [2], and we now briefly outline here. It motivates the definition of the spaces of exotic differential forms above.

Let $(H, B_\alpha, F_{\alpha\beta}, L_{\alpha\beta})$ denote a gerbe with connection on Z (cf. [9]), where $(H, B_\alpha, F_{\alpha\beta})$ denotes the Deligne class of the closed integral 3-form H with respect to a Brylinski open cover (cf. [13]) and $L_{\alpha\beta}$ denotes the line bundles on double overlaps that determines the gerbe \mathcal{G} . The holonomy of the gerbe is then a line bundle \mathcal{L} with connection $\tau(B_\alpha)$ having curvature $\tau(H)$ on loop space LZ , where τ denotes the transgression map. It $\iota_0 : Z \rightarrow LZ$ denotes the embedding of Z into LZ as the constant loops, then noting that $\iota_0^*(\mathcal{L})$ is canonically trivial, we proved that

$$(0.4) \quad \iota_0^* : \Omega^{\bar{k}}(LZ, \mathcal{L})^{S^1} \longrightarrow \Omega^{\bar{k}}(Z)$$

intertwines the equivariantly flat superconnection on the left hand side and $d + H$ on the right hand side, where the left hand side was called there the exotic differential forms on loop space.

The precise relation between [13] and this paper is that when Z is the total space of a principal circle bundle, then there is a natural infinite sequence of embeddings $\iota_n : Z \rightarrow LZ$ defined by $\iota_n(x) : S^1 \ni t \mapsto \gamma_x(t) = t^n \cdot x$, for all $n \in \mathbb{Z}$. We consider such sequence of embeddings motivated by the fact that there are \mathbb{Z} many connected components in the loop space $L\mathbb{T}$. We have $\iota_n^*(\mathcal{L}) \cong \pi^*(\hat{\xi})^{\otimes n}$ since they have the same Chern class. The loop space LZ has the natural circle action by rotating loops, and Z has a circle action as the total space of circle bundle. To tell the difference of these two circle actions, we use S^1 for the circle action by rotating loops. We have that for $n \neq 0$,

$$\iota_n^* : \Omega^{\bar{k}}(LZ, \mathcal{L})^{S^1} \longrightarrow \Omega^{\bar{k}}(Z, \pi^*(\hat{\xi}^{\otimes n}))^{\mathbb{T}}$$

intertwines the equivariantly flat superconnections on both spaces. We would like to point out that we don't automatically see the \mathbb{T} -invariance things on Z for $n = 0$ in the map (0.4). This comes from the effects of our embedding ι_n . However this point of view does motivate us to develop the exotic theories on Z and eventually relates to the Fourier expansion as discussed below.

Define the subspace of weight $-n$ differential forms on Z ,

$$(0.5) \quad \Omega_{-n}^*(Z) = \{\omega \in \Omega^*(Z) \mid L_v \omega = -n\omega\}.$$

It is easy to see that

$$\Omega_0^{\bar{k}}(Z) = \Omega^{\bar{k}}(Z)^{\mathbb{T}}, \quad \mathcal{A}_0^{\bar{k}+1}(\hat{Z})^{\hat{\mathbb{T}}} = \Omega^{\bar{k}+1}(\hat{Z})^{\hat{\mathbb{T}}}.$$

Under the above choices of Riemannian metrics and flux forms, our results show that the Fourier-Mukai transform T can be extended to a sequence of isometries,

$$(0.6) \quad \tau_n : \Omega_{-n}^{\bar{k}}(Z) \rightarrow \mathcal{A}_n^{\bar{k}+1}(\hat{Z})^{\hat{\mathbb{T}}},$$

for $\bar{k} = k \pmod{2}$, and is defined by the *exotic Hori formula* from Z to \hat{Z} given in equation (2.24). When $n = 0$, $\tau_0 = T$. Theorem 2.5 shows that the twisted de Rham differential $d + H$

maps to the differential $-(\hat{\pi}^*\nabla^{\xi^{\otimes n}} - \iota_{n\hat{v}} + \hat{H})$. One similarly has a sequence of isometries,

$$(0.7) \quad \sigma_n: \mathcal{A}_n^{\bar{k}}(Z)^{\mathbb{T}} \rightarrow \Omega_{-n}^{\bar{k}+1}(\hat{Z}),$$

for $\bar{k} = k \pmod{2}$, and is defined by the *inverse exotic Hori formula* from Z to \hat{Z} given in equation (2.28) and the differential $\pi^*\nabla^{\hat{\xi}^{\otimes n}} - \iota_{nv} + H$ maps to the twisted de Rham differential $-(d + \hat{H})$. Note that $\sigma_0 = T$. Similarly, one can define the sequences of isometries $\hat{\tau}_n, \hat{\sigma}_n$ on \hat{Z} . Although the extension of the Fourier-Mukai transform to all differential forms on Z is slightly asymmetric, one has the crucial identities, verified in Theorem 2.5,

$$(0.8) \quad -\text{Id} = \hat{\sigma}_n \circ \tau_n: \Omega_{-n}^{\bar{k}}(Z) \longrightarrow \Omega_{-n}^{\bar{k}}(Z),$$

$$(0.9) \quad -\text{Id} = \hat{\tau}_n \circ \sigma_n: \mathcal{A}_n^{\bar{k}}(Z)^{\mathbb{T}} \longrightarrow \mathcal{A}_n^{\bar{k}}(Z)^{\mathbb{T}}.$$

This is interpreted as saying that T-duality when applied twice, returns one to minus of the identity. It was verified in the special case when $n = 0$ in [6, 7]. We would like to point out that the minus sign comes from the convention of integration along the fiber.

This shows that for each of either Z or \hat{Z} , there are two theories (at degree 0 the two theories coincide), and there are also graded isomorphisms between the two theories of both sides.

Theorem 2.2 tells us that when $n \neq 0$, the complex $(\mathcal{A}_n^{\bar{k}+1}(\hat{Z})^{\hat{\mathbb{T}}}, \hat{\pi}^*\nabla^{\xi^{\otimes n}} - \iota_{n\hat{v}} + \hat{H})$ has vanishing cohomology. Therefore, when $n \neq 0$, the complex $(\Omega_{-n}^{\bar{k}}(Z), d + H)$ also has vanishing cohomology. In Corollary 2.6, we construct a homotopy to show this by taking advantage of the homotopy operator previously constructed in Theorem 2.2.

For a general form $\omega \in \Omega^*(Z)$, not necessarily \mathbb{T} -invariant, one can perform the family Fourier expansion (see Section 2.4) to get

$$\Omega^*(Z) \ni \omega = \sum_{n=-\infty}^{\infty} \omega_n \in \bigoplus_{n=-\infty}^{\infty} \Omega_n^*(Z)$$

with $\omega_n \in \Omega_n^{\bar{k}}(Z)$ and one takes the Fréchet space completion of the direct sum above. Since H is \mathbb{T} -invariant, if $(d + H)\omega = 0$, then $(d + H)\omega_n = 0$. Corollary 2.6 shows that ω_n must be $(d + H)$ -exact when $n \neq 0$. Therefore the cohomology of $(\Omega^*(Z), d + H)$ is concentrated on the $n = 0$ part, i.e. the \mathbb{T} -invariant part. This indicates why in the previous study, we only consider the \mathbb{T} -invariant forms.

Now we are able to define momentum and winding in the much larger configuration space of invariant exotic differential forms, $\mathcal{A}^{\bar{k}}(Z)^{\mathbb{T}}$ as follows. The multiple w of the infinitesimal generator v of the circle action on Z is the *winding*, as it agrees with winding around the circle direction when the circle bundle is a product, cf. [15]. The tensor power m of the line bundle ξ is the *momentum*, as it agrees with momentum when the circle bundle is a product, cf. [15]. In Theorem 2.1 and section 2.4, we show that the momentum on the spacetime Z needs to be equal to the winding number on the T-dual spacetime \hat{Z} , in order that the exotic differential is an equivariantly flat superconnection. The T-dual side also exhibits this phenomena. Thus our slogan here is,

The momentum (on spacetime Z) gets exchanged with the winding number (on the T -dual spacetime \hat{Z}) and the winding number (on the spacetime Z) gets exchanged with the momentum (on the T -dual spacetime \hat{Z}).

Finally, the invariant exact Courant algebroid $(TZ \oplus T^*Z)_H^{\mathbb{T}}$, together with the usual Dorfman bracket, $u \circ_H v$, has a representation (or action) on the exotic differential forms via a novel exotic Lie derivative $L_{X+\alpha}^\xi$ (c.f. Theorem 1.2) and where the Courant bracket is still as before, namely the antisymmetrization of the Dorfmann bracket. For the relationship between invariant Courant algebroids and T -duality, see [10].

Acknowledgements. The first author was partially supported by the grant AcRF R-146-000-218-112 from National University of Singapore. The second author was partially supported by funding from the Australian Research Council, through Discovery Project grant DP150100008 and by the Australian Laureate Fellowship FL170100020. The first author also thanks Qin Li for helpful discussions. The second author thanks Maxim Zabzine (Uppsala), who posed the question (private communication) that is answered in this paper. Both authors thank Peter Bouwknegt for sharing some insights, and also the Chern Institute of Mathematics (Tianjin) for providing a stimulating research environment on our visit.

1. EXACT COURANT ALGEBROIDS AND EXOTIC DIFFERENTIAL FORMS

In this section, we consider invariant exact Courant algebroids and their actions on invariant differential forms with coefficients in a line bundle.

Let M be a smooth manifold. Consider the generalized tangent bundle $\mathcal{T}M = TM \oplus T^*M$. On sections of $\mathcal{T}M$, there is a natural field of non-degenerate symmetric bilinear form, namely for $X + \alpha, Y + \beta \in \Gamma(TM \oplus T^*M)$, we put

$$(1.1) \quad \langle X + \alpha, Y + \beta \rangle = \frac{1}{2}(\iota_X \beta + \iota_Y \alpha).$$

The Clifford algebra $\text{Cliff}(TM \oplus T^*M)$ is the algebra with generators $\gamma_u, \mathcal{U} \in \Gamma(TM \oplus T^*M)$ and relations

$$(1.2) \quad \{\gamma_u, \gamma_v\} = 2\langle \mathcal{U}, \mathcal{V} \rangle.$$

Further assume that M admits a smooth \mathbb{T} -action and let ξ be a \mathbb{T} -equivariant complex line bundle over M . Let ∇^ξ be a \mathbb{T} -invariant connection on E and $H \in \Omega_{cl}^3(M)$ a closed 3-form such that

$$(1.3) \quad (\nabla^\xi - \iota_v + H)^2 + L_v^\xi = 0,$$

where v is the Killing vector field of the \mathbb{T} -action. $\nabla^\xi - \iota_v + H$ and L_v^ξ are operators acting on $\Omega^*(M, \xi)$, the space of smooth differential forms with coefficients in ξ .

It is known that the Clifford algebra $\text{Cliff}(TM \oplus T^*M)$ has a natural action on $\Omega^*(M)$. One can easily extend this action to $\Omega^*(M, \xi)$.

Lemma 1.1. *We have a representation of the Clifford algebra $\text{Cliff}(TM \oplus T^*M)$ on $\Omega^*(M, \xi)$ given by*

$$\gamma_{X+\alpha} \cdot \varphi = \iota_X \varphi + \alpha \wedge \varphi, \quad X + \alpha \in \Gamma(TM \oplus T^*M), \quad \varphi \in \Omega^*(M, \xi).$$

For $X + \alpha, Y + \beta \in \Gamma(TM \oplus T^*M)$ and $H \in \Omega_{cl}^3(M)$, define the (twisted) Dorfmann bracket or Loday bracket by

$$(1.4) \quad (X + \alpha) \circ_H (Y + \beta) = [X, Y] + L_X \beta - \iota_Y d\alpha + \iota_X \iota_Y H.$$

It is related to the (twisted) Courant bracket by

$$(1.5) \quad \mathcal{U} \circ_H \mathcal{V} = [[\mathcal{U}, \mathcal{V}]]_H + d\langle \mathcal{U}, \mathcal{V} \rangle, \quad \mathcal{U}, \mathcal{V} \in \Gamma(TM \oplus T^*M),$$

or conversely,

$$(1.6) \quad [[\mathcal{U}, \mathcal{V}]]_H = \frac{1}{2}(\mathcal{U} \circ_H \mathcal{V} - \mathcal{V} \circ_H \mathcal{U}).$$

For $\mathcal{U} = X + \alpha \in \Gamma(TM \oplus T^*M)$, we introduce an **exotic twisted Lie derivative along \mathcal{U}** on $\Omega^*(M, \xi)$ by

$$(1.7) \quad \mathcal{L}_{\mathcal{U}}^\xi = L_X^\xi - \mu_X^\xi + (d\alpha - \iota_v \alpha + \iota_X H) \wedge,$$

where L_X^ξ is the Lie derivative along the direction X and μ_X^ξ is the moment of the \mathbb{T} -invariant connection ∇^ξ along the direction X . Evidently, $\mathcal{L}_{\mathcal{U}}^\xi$ depends on ∇^ξ, v and H .

We have the following relations.

Theorem 1.2. *Let $\mathcal{U}, \mathcal{V} \in \Gamma(TM \oplus T^*M)$. Then on $\Omega^*(M, \xi)$, we have*

$$(1.8) \quad \begin{aligned} \{\gamma_{\mathcal{U}}, \gamma_{\mathcal{V}}\} &= 2\langle \mathcal{U}, \mathcal{V} \rangle, \\ \{\nabla^\xi - \iota_v + H, \gamma_{\mathcal{U}}\} &= \mathcal{L}_{\mathcal{U}}^\xi, \\ \{\nabla^\xi - \iota_v + H, \mathcal{L}_{\mathcal{U}}^\xi\} &= 0 \text{ on } \Omega^*(M, \xi)^\mathbb{T}, \\ [\mathcal{L}_{\mathcal{U}}^\xi, \gamma_{\mathcal{V}}] &= \gamma_{\mathcal{U} \circ_H \mathcal{V}}, \\ [\mathcal{L}_{\mathcal{U}}^\xi, \mathcal{L}_{\mathcal{V}}^\xi] &= \mathcal{L}_{\mathcal{U} \circ_H \mathcal{V}}^\xi = \mathcal{L}_{[[\mathcal{U}, \mathcal{V}]]_H}^\xi \text{ on } \Omega^*(M, \xi)^\mathbb{T}. \end{aligned}$$

Proof. The first relation can be proved in a verbatim way as the situation without the presence of ξ .

To prove the second relation, we have

$$(1.9) \quad \begin{aligned} &\{\nabla^\xi - \iota_v + H, \gamma_{\mathcal{U}}\} \\ &= \{\nabla^\xi - \iota_v + H, \iota_X + \alpha \wedge\} \\ &= \{\nabla^\xi, \iota_X\} + \{\nabla^\xi, \alpha \wedge\} - \{\iota_v, \iota_X\} + \{\iota_v, \alpha \wedge\} + \{H, \iota_X\} + \{H, \alpha \wedge\} \\ &= \{\nabla^\xi, \iota_X\} + d\alpha \wedge - \iota_v \alpha \wedge + \iota_X H \wedge \\ &= L_X^\xi - \mu_X^\xi + (d\alpha - \iota_v \alpha + \iota_X H) \wedge. \end{aligned}$$

To prove the third relation, simply notice that equation (1.3) tells us that on $\Omega^*(M, \xi)^\mathbb{T}$, $\{\nabla^\xi - \iota_v + H, \nabla^\xi - \iota_v + H\} = 0$ and apply the second relation.

To prove the fourth relation, we have

$$\begin{aligned}
& [L_X^\xi - \mu_X^\xi + (d\alpha - \iota_v \alpha + \iota_X H) \wedge, \iota_Y + \beta \wedge] \\
&= [L_X^\xi, \iota_Y] + [L_X^\xi, \beta \wedge] - [\mu_X^\xi, \iota_Y] - [\mu_X^\xi, \beta \wedge] + [d\alpha \wedge, \iota_Y] + [d\alpha \wedge, \beta \wedge] \\
(1.10) \quad & - [\iota_v \alpha \wedge, \iota_Y] - [\iota_v \alpha \wedge, \beta \wedge] + [\iota_X H \wedge, \iota_Y] - [\iota_X H \wedge, \beta \wedge] \\
&= \iota_{[X, Y]} + (L_X \beta) \wedge - 0 - 0 - \iota_Y (d\alpha) \wedge + 0 - 0 - 0 - \iota_Y (\iota_X H) - 0 \\
&= \iota_{[X, Y]} + (L_X \beta - \iota_Y (d\alpha) + \iota_X \iota_Y H) \wedge,
\end{aligned}$$

and this shows that

$$[\mathcal{L}_U^\xi, \gamma_V] = \gamma_{U \circ_H V}.$$

The last relation can be deduced by combining the second, the third and the fourth relation. \square

Antisymmetrizing the fourth relation, we get that

Corollary 1.3. $\forall \varphi \in \Omega^*(M, \xi)^\mathbb{T}$, the following identity holds,

$$\begin{aligned}
& \gamma_U \gamma_V \cdot ((\nabla^\xi - \iota_v + H)\varphi) \\
(1.11) \quad &= (\nabla^\xi - \iota_v + H)(\gamma_U \gamma_V \cdot \varphi) + \gamma_V \cdot ((\nabla^\xi - \iota_v + H)(\gamma_U \cdot \varphi)) \\
& - \gamma_U \cdot ((\nabla^\xi - \iota_v + H)(\gamma_V \cdot \varphi)) + \gamma_{[[U, V]]_H} \cdot \varphi.
\end{aligned}$$

2. T-DUALITY AND EXOTIC HORI FORMULAE

2.1. Review of T-duality. First we review the results in [6, 7], where the following situation is studied. We give more details here than the brief review in the introduction.

In [6, 7], spacetime Z was compactified in one direction. More precisely, Z is a principal \mathbb{T} -bundle over X

$$(2.1) \quad \begin{array}{ccc} \mathbb{T} & \longrightarrow & Z \\ & & \pi \downarrow \\ & & X \end{array}$$

classified up to isomorphism by its first Chern class $c_1(Z) \in H^2(X, \mathbb{Z})$. Assume that spacetime Z is endowed with an H -flux which is a representative in the degree 3 Deligne cohomology of Z , that is $H \in \Omega^3(Z)$ with integral periods (for simplicity, we drop factors of $\frac{1}{2\pi i}$), together with the following data. Consider a local trivialization $U_\alpha \times \mathbb{T}$ of $Z \rightarrow X$, where $\{U_\alpha\}$ is a good cover of X . Let $H_\alpha = H|_{U_\alpha \times \mathbb{T}} = dB_\alpha$, where $B_\alpha \in \Omega^2(U_\alpha \times \mathbb{T})$ and finally, $B_\alpha - B_\beta = F_{\alpha\beta} \in \Omega^1(U_{\alpha\beta} \times \mathbb{T})$. Then the choice of H -flux entails that we are given a local trivialization as above and locally defined 2-forms B_α on it, together with closed 2-forms $F_{\alpha\beta}$ defined on double overlaps, that is, $(H, B_\alpha, F_{\alpha\beta})$. Also the first Chern class of $Z \rightarrow X$ is represented in integral cohomology by (F, A_α) where $\{A_\alpha\}$ is a connection 1-form on $Z \rightarrow X$ and $F = dA_\alpha$ is the curvature 2-form of $\{A_\alpha\}$.

The T-dual is another principal \mathbb{T} -bundle over M , denoted by \hat{Z} ,

$$(2.2) \quad \begin{array}{ccc} \hat{\mathbb{T}} & \longrightarrow & \hat{Z} \\ & & \hat{\pi} \downarrow \\ & & X \end{array}$$

To define it, we see that $\pi_*(H_\alpha) = d\pi_*(B_\alpha) = d\hat{A}_\alpha$, so that $\{\hat{A}_\alpha\}$ is a connection 1-form whose curvature $d\hat{A}_\alpha = \hat{F}_\alpha = \pi_*(H_\alpha)$ that is, $\hat{F} = \pi_*H$. So let \hat{Z} denote the principal \mathbb{T} -bundle over M whose first Chern class is $c_1(\hat{Z}) = [\pi_*H, \pi_*(B_\alpha)] \in H^2(X; \mathbb{Z})$.

The Gysin sequence [3] for Z enables us to define a T-dual H -flux $[\hat{H}] \in H^3(\hat{Z}, \mathbb{Z})$, satisfying

$$(2.3) \quad c_1(Z) = \hat{\pi}_*\hat{H},$$

where π_* and similarly $\hat{\pi}_*$, denote the pushforward maps. Note that \hat{H} is not fixed by this data, since any integer degree 3 cohomology class on X that is pulled back to \hat{Z} also satisfies the requirements. However, \hat{H} is determined uniquely (up to cohomology) upon imposing the condition $[H] = [\hat{H}]$ on the correspondence space $Z \times_X \hat{Z}$ as will be explained now.

The *correspondence space* (sometimes called the doubled space) is defined as

$$Z \times_X \hat{Z} = \{(x, \hat{x}) \in Z \times \hat{Z} : \pi(x) = \hat{\pi}(\hat{x})\}.$$

Then we have the following commutative diagram,

$$\begin{array}{ccc} & (Z \times_X \hat{Z}, [H] = [\hat{H}]) & \\ & \swarrow p \quad \searrow \hat{p} & \\ (Z, [H]) & & (\hat{Z}, [\hat{H}]) \\ & \searrow \pi \quad \swarrow \hat{\pi} & \\ & X & \end{array}$$

By requiring that

$$p^*[H] = \hat{p}^*[\hat{H}] \in H^3(Z \times_X \hat{Z}, \mathbb{Z}),$$

determines $[\hat{H}] \in H^3(\hat{Z}, \mathbb{Z})$ uniquely, via an application of the Gysin sequence. An alternate way to see this is explained below.

Let $(H, B_\alpha, F_{\alpha\beta}, L_{\alpha\beta})$ denote a gerbe with connection on Z , cf. [9], where $(H, B_\alpha, F_{\alpha\beta})$ denotes the Deligne class of the closed integral 3-form H and $L_{\alpha\beta}$ denotes the line bundles on double overlaps that determines the gerbe. We also choose a connection 1-form A on Z . Let v denote the vector field generating the \mathbb{T} -action on Z . Then define $\hat{A}_\alpha = -\iota_v B_\alpha$ on the chart U_α and the connection 1-form $\hat{A} = \hat{A}_\alpha + d\hat{\theta}_\alpha$ on the chart $U_\alpha \times \hat{\mathbb{T}}$. In this way we get a T-dual circle bundle $\hat{Z} \rightarrow X$ with connection 1-form \hat{A} .

Without loss of generality, we can assume that H is \mathbb{T} -invariant. Consider

$$\Omega = H - A \wedge F_{\hat{A}}$$

where $F_{\hat{A}} = d\hat{A}$ and $F_A = dA$ are the curvatures of A and \hat{A} respectively. One checks that the contraction $i_v(\Omega) = 0$ and the Lie derivative $L_v(\Omega) = 0$ so that Ω is a basic 3-form on Z , that is Ω comes from the base X .

Setting

$$\hat{H} = F_A \wedge \hat{A} + \Omega$$

this defines the T-dual flux 3-form. One verifies that \hat{H} is a closed 3-form on \hat{Z} . It follows that on the correspondence space, one has as desired,

$$(2.4) \quad \hat{H} = H + d(A \wedge \hat{A}).$$

Our next goal is to determine the T-dual curving or B-field. The Buscher rules imply that on the open sets $U_\alpha \times \mathbb{T} \times \hat{\mathbb{T}}$ of the correspondence space $Z \times_X \hat{Z}$, one has

$$(2.5) \quad \hat{B}_\alpha = B_\alpha + A \wedge \hat{A} - d\theta_\alpha \wedge d\hat{\theta}_\alpha,$$

Note that

$$(2.6) \quad \iota_v \hat{B}_\alpha = \iota_v (B_\alpha + A \wedge \hat{A} - d\theta_\alpha \wedge d\hat{\theta}_\alpha) = -\hat{A}_\alpha + \hat{A} - d\hat{\theta}_\alpha = 0$$

so that \hat{B}_α is indeed a 2-form on \hat{Z} and not just on the correspondence space. Obviously, $d\hat{B}_\alpha = \hat{H}$. Following the descent equations one arrives at the complete T-dual gerbe with connection, $(\hat{H}, \hat{B}_\alpha, \hat{F}_{\alpha\beta}, \hat{L}_{\alpha\beta})$. cf. [8].

Define the Riemannian metrics on Z and \hat{Z} respectively by

$$g = \pi^* g_X + R^2 A \odot A, \quad \hat{g} = \hat{\pi}^* g_X + 1/R^2 \hat{A} \odot \hat{A}.$$

where g_X is a Riemannian metric on X . Then g is \mathbb{T} -invariant and the length of each circle fibre is R ; \hat{g} is $\hat{\mathbb{T}}$ -invariant and the length of each circle fibre is $1/R$.

The rules for transforming the Ramond-Ramond (RR) fields can be encoded in the [6, 7] generalization of *Hori's formula*

$$(2.7) \quad T_* G = \int^{\mathbb{T}} e^{-A \wedge \hat{A}} G,$$

where $G \in \Omega^\bullet(Z)^\mathbb{T}$ is the total RR field-strength,

$$\begin{aligned} G &\in \Omega^{\text{even}}(Z)^\mathbb{T} && \text{for Type IIA;} \\ G &\in \Omega^{\text{odd}}(Z)^\mathbb{T} && \text{for Type IIB,} \end{aligned}$$

and where the right hand side of equation (2.7) is an invariant differential form on $Z \times_X \hat{Z}$, and the integration is along the \mathbb{T} -fiber of Z .

Recall that the twisted cohomology is defined as the cohomology of the complex

$$(2.8) \quad H^\bullet(Z, H) = H^\bullet(\Omega^\bullet(Z), d_H = d + H \wedge).$$

By the identity (2.7), T_* maps d_H -closed forms G to $d_{\hat{H}}$ -closed forms T_*G . So T-duality T_* induces a map on twisted cohomologies,

$$T : H^\bullet(Z, H) \rightarrow H^{\bullet+1}(\hat{Z}, \hat{H}).$$

2.2. Exotic theories. Let $\xi, \hat{\xi}$ be the complex line bundle determined by the circle bundles Z, \hat{Z} and the standard representation of the circle on the complex plane. Let ∇^ξ and $\nabla^{\hat{\xi}}$ be the connections on $\xi, \hat{\xi}$ induced from the connections on Z, \hat{Z} respectively. Let v, \hat{v} be the vertical tangent vector fields on Z, \hat{Z} respectively as in the above section.

Theorem 2.1. $\forall n \in \mathbb{Z}$, we have:

on $\Omega^*(Z, \pi^*(\hat{\xi}^{\otimes n}))$, the following identity holds,

$$(2.9) \quad (\pi^*\nabla^{\hat{\xi}^{\otimes n}} - \iota_{nv} + H)^2 + nL_v^{\hat{\xi}^{\otimes n}} = 0;$$

on $\Omega^*(\hat{Z}, \hat{\pi}^*(\xi^{\otimes n}))$, the following identity holds,

$$(2.10) \quad (\hat{\pi}^*\nabla^{\xi^{\otimes n}} - \iota_{n\hat{v}} + \hat{H})^2 + nL_{\hat{v}}^{\xi^{\otimes n}} = 0.$$

Proof. Let $\{\hat{s}_\alpha\}$ be local sections of the line bundle $\hat{\xi}$ such that the connection 1-form corresponding to whom is $\{\hat{A}_\alpha\}$. It is obvious that $\pi^*\hat{s}_\alpha$'s are invariant about the \mathbb{T} -action on Z . Under $\{\pi^*\hat{s}_\alpha\}$, to prove the first equality, we only need to prove that for forms on $U_\alpha \times \mathbb{T}$,

$$(2.11) \quad (d + \pi^*(n\hat{A}_\alpha) - \iota_{nv} + H)^2 + nL_v = 0.$$

But this is evident, as we have

$$\begin{aligned} (d - \iota_{nv})^2 + nL_v &= 0, \\ dH &= 0, \quad \iota_v \pi^*(\hat{A}_\alpha) = 0, \\ d\pi^*(n\hat{A}_\alpha) - n\iota_v H &= n\pi^*(d\hat{A}_\alpha) - n\iota_v H = 0. \end{aligned}$$

One can similarly prove the second equality. □

Denote

$$\begin{aligned} \mathcal{A}^*(Z) &= \bigoplus_{n \in \mathbb{Z}} \mathcal{A}_n^*(Z) := \bigoplus_{n \in \mathbb{Z}} \Omega^*(Z, \pi^*(\hat{\xi}^{\otimes n}))^{\mathbb{T}}, \\ \mathcal{A}^*(\hat{Z}) &= \bigoplus_{n \in \mathbb{Z}} \mathcal{A}_n^*(\hat{Z}) := \bigoplus_{n \in \mathbb{Z}} \Omega^*(\hat{Z}, \hat{\pi}^*(\xi^{\otimes n}))^{\hat{\mathbb{T}}}. \end{aligned}$$

For each $n \in \mathbb{Z}$, consider the complexes

$$\begin{aligned} (\mathcal{A}_n^*(Z), \pi^*\nabla^{\hat{\xi}^{\otimes n}} - \iota_{nv} + H), \\ (\mathcal{A}_n^*(\hat{Z}), \hat{\pi}^*\nabla^{\xi^{\otimes n}} - \iota_{n\hat{v}} + \hat{H}). \end{aligned}$$

We have

Theorem 2.2. *If $n \neq 0$, then*

$$(2.12) \quad H(\mathcal{A}_n^*(Z), \pi^*\nabla^{\hat{\xi}^{\otimes n}} - \iota_{nv} + H) \cong \{0\},$$

$$(2.13) \quad H(\mathcal{A}_n^*(\hat{Z}), \hat{\pi}^*\nabla^{\xi^{\otimes n}} - \iota_{n\hat{v}} + \hat{H}) \cong \{0\}.$$

Proof. For $n \neq 0$, set $\eta_n = \frac{A}{F_A - n}$, where A is the connection 1-form on Z and F_A is its curvature 2-form. Then we have

$$\begin{aligned}
& (d - \iota_{nv})\eta_n \\
&= \frac{[(d - \iota_{nv})A](F_A - n) - A[(d - \iota_{nv})(F_A - n)]}{(F_A - n)^2} \\
(2.14) \quad &= \frac{(F_A - n)^2}{(F_A - n)^2} \\
&= 1.
\end{aligned}$$

We therefore obtain a homotopy for $n \neq 0$: $\forall x \in \Omega^*(Z, \pi^*(\hat{\xi}^{\otimes n}))$.

$$\begin{aligned}
& (\pi^*\nabla^{\hat{\xi}^{\otimes n}} - \iota_{nv} + H)(\eta_n x) + \eta_n[(\pi^*\nabla^{\hat{\xi}^{\otimes n}} - \iota_{nv} + H)x] \\
(2.15) \quad &= [(d - \iota_{nv})\eta_n]x \\
&= x.
\end{aligned}$$

We can prove the second isomorphism in a verbatim way by using $\hat{A}, F_{\hat{A}}$ to construct the homotopy. □

2.3. Twisted integration along the fiber. In order to define the exotic Hori formula to be introduced in the next subsection, we first introduce a twisted version of integration along the fiber.

Let $\pi : P \rightarrow M$ be a principal circle bundle over M , and Θ a connection one form on P . Let L be a Hermitian line bundle over M such that Z is the circle bundle of L . Let ∇^L be the connection on L corresponding to Θ . Choose a good cover $\{U_\alpha\}$ on M such that $\pi^{-1}(U_\alpha) \cong U_\alpha \times S^1$. Let $\{f_\alpha\}$ be a local basis of L corresponding to the constant map $U_\alpha \rightarrow \{1\} \subset S^1$.

$\forall n \in \mathbb{Z}$, define the **twisted integration along the fiber** as follows: for $\omega \in \Omega^*(P)$,

$$(2.16) \quad \int^{P/M, n} \omega \in \Omega^*(M, L^{\otimes n}), \text{ such that } \left(\int^{P/M, n} \omega \right) \Big|_{U_\alpha} = \left(\int^{\pi^{-1}(U_\alpha)/U_\alpha} \omega_\alpha e^{2\pi i n \theta_\alpha} \right) \otimes f_\alpha^{\otimes n},$$

where $\omega_\alpha = \omega|_{\pi^{-1}(U_\alpha)}$, θ_α is the vertical coordinates of $\pi^{-1}(U_\alpha)$. Note that as on $U_{\alpha\beta} = U_\alpha \cap U_\beta$, $f_\alpha/f_\beta = e^{2\pi i(\theta_\beta - \theta_\alpha)}$ (a function on $U_{\alpha\beta}$), the above construction patches to be a global section of the bundle $\wedge^*(T^*M) \otimes L^{\otimes n}$. Moreover, it is not hard to see that this definition is independent of choice of the good cover $\{U_\alpha\}$ and local trivializations.

Theorem 2.3. *Let Y be a vector field on M and \tilde{Y} a lift of Y on P . Let H be a differential form on M . Then $\forall n \in \mathbb{Z}$*

$$(2.17) \quad (\nabla^{L^{\otimes n}} - \iota_Y + H) \int^{P/M, n} \omega = - \int^{P/M, n} (d + n\Theta - \iota_{\tilde{Y}} + H)\omega.$$

Proof. For the definition of the usual integration along the fiber, we refer to [1]. It is not hard to see that

$$(2.18) \quad H \cdot \int^{P/M,n} \omega = - \int^{P/M,n} H \cdot \omega,$$

$$(2.19) \quad \iota_Y \int^{P/M,n} \omega = - \int^{P/M,n} \iota_{\tilde{Y}} \omega.$$

One only needs to prove

$$(2.20) \quad \nabla^{L \otimes n} \left(\int^{P/M,n} \omega \right) = - \int^{P/M,n} (d + n\Theta)\omega.$$

In fact, suppose Θ_α is the connection 1-form under the local basis f_α , we have

$$(2.21) \quad \begin{aligned} & \left(\nabla^{L \otimes n} \left(\int^{P/M,n} \omega \right) \right) \Big|_{U_\alpha} \\ &= \left[d \int^{\pi^{-1}(U_\alpha)/U_\alpha} \omega_\alpha e^{2\pi i n \theta_\alpha} + (-1)^{\deg \omega - 1} \left(\int^{\pi^{-1}(U_\alpha)/U_\alpha} \omega_\alpha e^{2\pi i n \theta_\alpha} \right) n \Theta_\alpha \right] \otimes f_\alpha^{\otimes n} \\ &= \left[d \int^{\pi^{-1}(U_\alpha)/U_\alpha} \omega_\alpha e^{2\pi i n \theta_\alpha} - \int^{\pi^{-1}(U_\alpha)/U_\alpha} n \Theta_\alpha \omega_\alpha e^{2\pi i n \theta_\alpha} \right] \otimes f_\alpha^{\otimes n} \\ &= \left[- \int^{\pi^{-1}(U_\alpha)/U_\alpha} d(\omega_\alpha e^{2\pi i n \theta_\alpha}) - \int^{\pi^{-1}(U_\alpha)/U_\alpha} n \Theta_\alpha \omega_\alpha e^{2\pi i n \theta_\alpha} \right] \otimes f_\alpha^{\otimes n} \\ &= \left[- \int^{\pi^{-1}(U_\alpha)/U_\alpha} \left((d\omega_\alpha) e^{2\pi i n \theta_\alpha} + (-1)^{\deg \omega} \omega_\alpha e^{2\pi i n \theta_\alpha} (2\pi i n d\theta_\alpha) \right) \right. \\ & \quad \left. - \int^{\pi^{-1}(U_\alpha)/U_\alpha} n \Theta_\alpha \omega_\alpha e^{2\pi i n \theta_\alpha} \right] \otimes f_\alpha^{\otimes n} \\ &= - \left[\int^{\pi^{-1}(U_\alpha)/U_\alpha} [(d\omega_\alpha + n\Theta_\alpha + n2\pi i d\theta_\alpha)\omega_\alpha] e^{2\pi i n \theta_\alpha} \right] \otimes f_\alpha^{\otimes n} \\ &= - \left(\int^{P/M,n} (d + n\Theta)\omega \right) \Big|_{U_\alpha}. \end{aligned}$$

The desired equality follows. □

2.4. The exotic Hori formulae. Let us go back to the T-duality with same notions as in Section 2.1, 2.2.

Let $X + \alpha \in \Gamma(TZ \oplus T^*Z)^\mathbb{T}$. Then one can write

$$(2.22) \quad X = x + fv, \quad \alpha = \theta + gA,$$

where $x \in \Gamma(TM)$, $\theta \in \Omega^1(M)$, $f, g \in C^\infty(M)$. Define

$$\phi(X, \alpha) = (x + gv) + (\theta + fA).$$

Recall that

$$(2.23) \quad \Omega_{-n}^*(Z) = \{\omega \in \Omega^*(Z) | L_v \omega = -n\omega\}.$$

Note that $\Omega_{-n}^*(Z) = \Omega^*(Z)^\mathbb{T}$, i.e. the \mathbb{T} -invariant forms on Z .

Let $\omega_{-n} \in \Omega_{-n}^*(Z)$. Define the **exotic Hori formula** by

$$(2.24) \quad \tau_n(\omega_{-n}) = \int^{\mathbb{T}, n} \omega_{-n} e^{-A \wedge \hat{A}} \in \Omega^*(\hat{Z}, \hat{\pi}^*(\xi)^{\otimes n}),$$

where $\int^{\mathbb{T}, n}$ stands for $\int^{(Z \times_X \hat{Z})/\hat{Z}, n}$ for simplicity.

Remark 2.4. (i) Let $\{s_\alpha\}$ be local sections of the line bundle ξ and θ_α be the vertical coordinate function on $\pi^{-1}(U_\alpha)$. Then, locally, ω_{-n} must be of the form

$$(\omega_{-n, \alpha, 0} + \omega_{-n, \alpha, 1}(d\theta_\alpha + A_\alpha))e^{-2\pi i n \theta_\alpha},$$

where $\omega_{-n, \alpha, 0}$ and $\omega_{-n, \alpha, 1}$ are both forms on U_α . So if $m \neq n$,

$$(2.25) \quad \begin{aligned} & \int^{\mathbb{T}, m} \omega_{-n} e^{-A \wedge \hat{A}} \Big|_{U_\alpha} \\ &= \left(\int^{\mathbb{T}} (\omega_{-n, \alpha, 0} + \omega_{-n, \alpha, 1}(d\theta_\alpha + A_\alpha))(1 - (d\theta_\alpha + A_\alpha) \wedge \hat{A})e^{-2\pi i n \theta_\alpha} \cdot e^{2\pi i m \theta_\alpha} \right) \otimes \hat{\pi}^*(s_\alpha)^{\otimes n} \\ &= 0. \end{aligned}$$

This explains why we only define $\tau_m(\omega_{-n})$ for $m = n$.

(ii) When $n = 0$, τ_0 is just the Hori formula (0.1) in [6, 7].

Denote by ρ the tautological global section of the line bundle $(\hat{\pi} \circ \hat{p})^* \xi$ on $Z \times_X \hat{Z}$. Let $\hat{\theta}_n \in \Omega^*(\hat{Z}, \hat{\pi}^*(\xi)^{\otimes n})^{\hat{\mathbb{T}}}$. Define the **inverse exotic Hori formula** by

$$(2.26) \quad \hat{\sigma}_n(\hat{\theta}_n) = \int^{\hat{\mathbb{T}}} \hat{p}^*(\hat{\theta}_n) \cdot (\rho^{-1})^{\otimes n} \cdot e^{A \wedge \hat{A}} \in \Omega^*(Z).$$

One can dually define the exotic Hori formula $\hat{\tau}_n$ from \hat{Z} to Z and the inverse exotic Hori formula σ_n from Z to \hat{Z} . Let $\hat{\omega}_{-n} \in \Omega_{-n}^*(\hat{Z})$. Define

$$(2.27) \quad \hat{\tau}_n(\hat{\omega}_{-n}) = \int^{\hat{\mathbb{T}}, n} \hat{\omega}_{-n} e^{A \wedge \hat{A}} \in \Omega^*(Z, \pi^*(\hat{\xi})^{\otimes n}).$$

Denote by $\hat{\rho}$ the tautological global section of the line bundle $(\pi \circ p)^* \hat{\xi}$ on $Z \times_X \hat{Z}$. Let $\theta_n \in \Omega^*(Z, \pi^*(\hat{\xi})^{\otimes n})^\mathbb{T}$. Define

$$(2.28) \quad \sigma_n(\theta_n) = \int^{\mathbb{T}} p^*(\theta_n) \cdot (\hat{\rho}^{-1})^{\otimes n} \cdot e^{-A \wedge \hat{A}} \in \Omega^*(\hat{Z}).$$

We have the following results:

Theorem 2.5. (1) ϕ is orthogonal with respect to the pairing on $TZ \oplus T^*Z$, hence induces an isomorphism on Clifford algebras.

(2) ϕ preserves the twisted Courant bracket.

(3) $\tau_n(\omega_{-n}) \in \Omega^*(\hat{Z}, \hat{\pi}^*(\xi^{\otimes n}))^{\hat{\mathbb{T}}}$ and $\hat{\sigma}_n(\theta_n) \in \Omega_{-n}^*(Z)$. For $\mathcal{U} \in \Gamma(TZ \oplus T^*Z)^{\mathbb{T}}$, we have $\tau_n(\gamma_{\mathcal{U}} \cdot \{\omega_{-n}\}) = \gamma_{\phi(\mathcal{U})} \cdot \tau_n(\{\omega_{-n}\})$, for all $\{\omega_{-n}\} \in \Omega_{-n}^*(Z)$, hence induces an isomorphism of Clifford modules

$$\tau_n : \Omega_{-n}^*(Z) \rightarrow \Omega^*(\hat{Z}, \hat{\pi}^*(\xi^{\otimes n}))^{\hat{\mathbb{T}}}.$$

$\hat{\sigma}_n = -\tau_n^{-1}$ and is an isomorphism of Clifford modules

$$\hat{\sigma}_n : \Omega^*(\hat{Z}, \hat{\pi}^*(\xi^{\otimes n}))^{\hat{\mathbb{T}}} \rightarrow \Omega_{-n}^*(Z).$$

The dual results for $\hat{\tau}_n$ and σ_n are also true.

(4) The map τ_n induces a chain map on the complexes

$$(\Omega_{-n}^*(Z), d + H) \rightarrow (\Omega^*(\hat{Z}, \hat{\pi}^*(\xi^{\otimes n}))^{\hat{\mathbb{T}}}, -(\hat{\pi}^*\nabla^{\xi^{\otimes n}} - \iota_{n\hat{v}} + \hat{H}))$$

and the map $\hat{\sigma}_n$ induces a chain map on the complexes

$$(\Omega^*(\hat{Z}, \hat{\pi}^*(\xi^{\otimes n}))^{\hat{\mathbb{T}}}, \hat{\pi}^*\nabla^{\xi^{\otimes n}} - \iota_{n\hat{v}} + \hat{H}) \rightarrow (\Omega_{-n}^*(Z), -(d + H)).$$

The dual results for $\hat{\tau}_n$ and σ_n are also true.

(1), (2) in the above theorem as well as (3) and (4) without the presence of the line bundles $\xi, \hat{\xi}$ are existing results ([11, 12], c.f. [4]).

Proof. We will prove (3) and (4).

Let $\{s_\alpha\}$ be local sections of the line bundle ξ and $\{\hat{s}_\alpha\}$ local sections of the line bundle $\hat{\xi}$. Let θ_α be the vertical coordinate function of $\pi^{-1}(U_\alpha)$ and $\hat{\theta}_\alpha$ the vertical coordinate function of $\hat{\pi}^{-1}(U_\alpha)$.

Take $\{\omega_n\} \in \Omega_{-n}^*(Z)$. Locally, ω_{-n} is of the form

$$(\omega_{-n,\alpha,0} + \omega_{-n,\alpha,1}A)e^{-2\pi i n \theta_\alpha},$$

where $\omega_{-n,\alpha,0}$ and $\omega_{-n,\alpha,1}$ are both forms on U_α . Then

$$\begin{aligned} & \tau_n(\omega_{-n})|_{U_\alpha} \\ &= \int^{\mathbb{T},n} \omega_n e^{-A \wedge \hat{A}} \Big|_{U_\alpha} \\ (2.29) \quad &= \left(\int^{\mathbb{T}} (\omega_{-n,\alpha,0} + \omega_{-n,\alpha,1}A) e^{-A \wedge \hat{A}} e^{-2\pi i n \theta_\alpha} \cdot e^{2\pi i n \theta_\alpha} \right) \otimes \hat{\pi}^*(s_\alpha)^{\otimes n} \\ &= \left(\int^{\mathbb{T}} (\omega_{-n,\alpha,0} + \omega_{-n,\alpha,1}A) e^{-A \wedge \hat{A}} \right) \otimes \hat{\pi}^*(s_\alpha)^{\otimes n}, \end{aligned}$$

which is equal to

$$(-\omega_{-n,\alpha,0}\hat{A} - \omega_{-n,\alpha,1}) \otimes \hat{\pi}^*(s_\alpha)^{\otimes n}$$

if ω_{-n} is of even degree; or

$$(\omega_{-n,\alpha,0}\hat{A} + \omega_{-n,\alpha,1}) \otimes \hat{\pi}^*(s_\alpha)^{\otimes n}$$

if ω_{-n} is of odd degree. So we see that

$$\tau_n(\omega_{-n}) \in \Omega^*(\hat{Z}, \hat{\pi}^*(\xi^{\otimes n}))^{\hat{\mathbb{T}}}.$$

On the other hand, if $\hat{\theta}_n \in \Omega^*(\hat{Z}, \hat{\pi}^*(\xi)^{\otimes n})^{\hat{\mathbb{T}}}$, suppose $\hat{\theta}_n$ is locally equal to

$$(\hat{\eta}_{n,\alpha,0}\hat{A} + \hat{\eta}_{n,\alpha,1}) \otimes \hat{\pi}^*(s_\alpha)^{\otimes n},$$

where $\hat{\eta}_{n,\alpha,0}, \hat{\eta}_{n,\alpha,1}$ are forms on U_α . then $\hat{\sigma}_n(\hat{\theta}_n)$ is locally equal to

$$(2.30) \quad \int_{U_\alpha}^{\hat{\mathbb{T}}} \hat{p}^*(\hat{\theta}_n) \cdot (\rho^{-1})^{\otimes n} \cdot e^{A \wedge \hat{A}} \Big|_{U_\alpha} = \left(\int_{U_\alpha}^{\hat{\mathbb{T}}} (\hat{\eta}_{n,\alpha,0}\hat{A} + \hat{\eta}_{n,\alpha,1}) \cdot e^{A \wedge \hat{A}} \right) \cdot e^{-2\pi i n \theta_\alpha},$$

which is equal to

$$(\hat{\eta}_{n,\alpha,0} + \hat{\eta}_{n,\alpha,1}A) \cdot e^{-2\pi i n \theta_\alpha}$$

if $\hat{\theta}_n$ is of even degree; or

$$-(\hat{\eta}_{n,\alpha,0} + \hat{\eta}_{n,\alpha,1}A) \cdot e^{-2\pi i n \theta_\alpha}$$

if $\hat{\theta}_n$ is of odd degree. And so evidently $L_v \hat{\sigma}_n(\hat{\theta}_n) = -n \hat{\sigma}_n(\hat{\theta}_n)$. We therefore have

$$\hat{\sigma}_n(\hat{\theta}_n) \in \Omega_{-n}^*(Z).$$

From the above expressions (2.29), (2.30) and local nature of (3), we see from the original Hori formula that $\tau_n, \hat{\sigma}_n$ both respect the Clifford actions and

$$(2.31) \quad \hat{\sigma}_n = -\tau_n^{-1}.$$

We next prove (4). We have

$$(2.32) \quad [(d + H)\omega_{-n}]e^{-A \wedge \hat{A}} = [d + H - (H - \hat{H})](\omega_{-n}e^{-A \wedge \hat{A}}) = (d + \hat{H})(\omega_{-n}e^{-A \wedge \hat{A}}).$$

Also one has

$$(2.33) \quad (nA - \iota_{n\hat{v}})(\omega_{-n}e^{-A \wedge \hat{A}}) = nA\omega_{-n}e^{-A \wedge \hat{A}} - \iota_{n\hat{v}}(\omega_{-n}e^{-A \wedge \hat{A}}) = (-1)^{|\omega_{-n}|}\omega_{-n}(nA - nA) = 0.$$

By Theorem 2.3, we have

$$(2.34) \quad \begin{aligned} & \tau_n((d + H)\omega_{-n}) \\ &= \int^{\mathbb{T},n} [(d + H)\omega_{-n}] \cdot e^{-A \wedge \hat{A}} \\ &= \int^{\mathbb{T},n} (d + \hat{H})(\omega_{-n} \cdot e^{-A \wedge \hat{A}}) \\ &= \int^{\mathbb{T},n} (d + nA - \iota_{n\hat{v}} + \hat{H})(\omega_{-n} \cdot e^{-A \wedge \hat{A}}) \\ &= -(\hat{\pi}^* \nabla^{\xi^{\otimes n}} - \iota_{n\hat{v}} + \hat{H}) \int^{\mathbb{T},n} \omega_{-n} \cdot e^{-A \wedge \hat{A}} \\ &= -(\hat{\pi}^* \nabla^{\xi^{\otimes n}} - \iota_{n\hat{v}} + \hat{H})\tau_n(\omega_{-n}). \end{aligned}$$

As $-\hat{\sigma}_n$ is the inverse of τ_n , one deduces that $\hat{\sigma}_n$ is also a chain map. □

Combining Theorem 2.2, we have

Corollary 2.6. *If $n \neq 0$, then*

$$(2.35) \quad H(\Omega_{-n}^*(Z), d + H) = 0,$$

$$(2.36) \quad H(\Omega_{-n}^*(\hat{Z}), d + \hat{H}) = 0.$$

Actually, from the proof of Theorem 2.2, we have the following homotopy

$$(2.37) \quad (d + H)\hat{\sigma}_n[\hat{\eta}_n \cdot \tau_n(\omega_{-n})] + \hat{\sigma}_n(\hat{\eta}_n \cdot \tau_n[(d + H)\omega_{-n}]) = \omega_{-n},$$

where $\omega_{-n} \in \Omega_{-n}^*(Z)$ and $\hat{\eta}_n = \frac{\hat{A}}{F_{\hat{A}}^{-n}}$. It is interesting to see that to construct this homotopy on $(\Omega_{-n}^*(Z), d + H)$, one uses the data on the dual side \hat{Z} . The interested readers may write a homotopy for the dual case.

For a general form $\omega \in \Omega^*(Z)$, one can perform the **family Fourier expansion** as follows. Suppose locally on U_α ,

$$(2.38) \quad \omega = \sum_I f_I(x, \theta_\alpha) dx_I + \sum_J g_J(x, \theta_\alpha) dx_J \wedge A.$$

Consider the local form

$$\sum_I \left(\int^{\mathbb{T}} f_I(x, \theta_\alpha) e^{2\pi i n \theta_\alpha} d\theta_\alpha \right) dx_I + \sum_J \left(\int^{\mathbb{T}} g_J(x, \theta_\alpha) e^{2\pi i n \theta_\alpha} d\theta_\alpha \right) dx_J \wedge A.$$

As $e^{2\pi i n(\theta_\alpha - \theta_\beta)}$ is a function on the base, the form

$$\left[\sum_I \left(\int^{\mathbb{T}} f_I(x, \theta_\alpha) e^{2\pi i n \theta_\alpha} d\theta_\alpha \right) dx_I + \sum_J \left(\int^{\mathbb{T}} g_J(x, \theta_\alpha) e^{2\pi i n \theta_\alpha} d\theta_\alpha \right) dx_J \wedge A \right] e^{-2\pi i n \theta_\alpha}$$

for each α glue together to be a global form on Z . Denote this form by ω_{-n} . Since $\left(\int^{\mathbb{T}} f_I(x, \theta_\alpha) e^{2\pi i n \theta_\alpha} d\theta_\alpha \right) dx_I$, $\left(\int^{\mathbb{T}} g_J(x, \theta_\alpha) e^{2\pi i n \theta_\alpha} d\theta_\alpha \right) dx_J$ as well as A are all \mathbb{T} -invariant, we see that $L_\nu \omega_{-n} = -n\omega_{-n}$. By the Fourier expansion, $\omega = \sum_{n=-\infty}^{\infty} \omega_{-n}$.

Let s_α be the local basis of the bundle ξ . Then

$$(2.39) \quad \left[\sum_I \left(\int^{\mathbb{T}} f_I(x, \theta_\alpha) e^{2\pi i n \theta_\alpha} d\theta_\alpha \right) dx_I + \sum_J \left(\int^{\mathbb{T}} g_J(x, \theta_\alpha) e^{2\pi i n \theta_\alpha} d\theta_\alpha \right) dx_J \wedge A \right] \otimes \pi^* s_\alpha^{\otimes n}$$

patch together to be an element $\Omega_{-n} \in \Omega^*(Z, \pi^* \xi^{\otimes n})$. Let γ be the tautological global section of the bundle $\pi^* \xi$ over Z . Then

$$(2.40) \quad \omega = \sum_{n=-\infty}^{\infty} \omega_{-n} = \sum_{n=-\infty}^{\infty} \Omega_{-n} \otimes (\gamma^{-1})^{\otimes n}.$$

We call Ω_{-n} 's the **family Fourier coefficients** of ω .

The above theorem shows that

$$\tau_n(\omega_{-n}) = \tau_n(\Omega_{-n} \otimes (\gamma^{-1})^{\otimes n}) \in \Omega^*(\hat{Z}, \hat{\pi}^*(\xi)^{\otimes n})^{\hat{\mathbb{T}}}.$$

The n on the left hand side of the above equality should be the momentum of Z , as it is n -th power of ξ . Let

$$\omega_{-n} - (\iota_v \omega_{-n})A \neq 0,$$

i.e. ω_{-n} is not some kind of product. Suppose

$$(d + H)(\omega_{-n}) = 0.$$

Then from the proof of Theorem 2.5, we see that

$$(2.41) \quad (\hat{\pi}^* \nabla^{\xi^{\otimes n}} - \iota_{m\hat{v}} + \hat{H})\tau_n(\omega_{-n}) = 0 \iff m = n,$$

where m is the winding of \hat{Z} as it the multiple of \hat{v} . This clearly also applies for the trivial bundle case.

2.5. The trivial bundles case. Consider the trivial bundles case. Now

$$Z = M \times \mathbb{T}, \quad \hat{Z} = M \times \hat{\mathbb{T}}$$

and H, \hat{H} and the connections are all 0.

Pick $\omega_{-n} \in \Omega^{\text{even}}(Z)_{-n}$. It is of the form $(\lambda_0 + \lambda_1 d\theta)e^{-2\pi i n \theta}$, where λ_0, λ_1 are forms on M . Then by definition,

$$\tau_n(\omega_{-n}) = -\lambda_0 d\hat{\theta} - \lambda_1, \quad \hat{\sigma}_n(-\lambda_0 d\hat{\theta} - \lambda_1) = -\lambda_0 - \lambda_1 d\theta.$$

Suppose $d\omega_{-n} = 0$.

We have

$$d\lambda_0 = 0, \quad d\lambda_1 - n\lambda_0 = 0.$$

Then

$$(d - \iota_{n\hat{v}})\tau_n(\omega_{-n}) = -(d - \iota_{n\hat{v}})(\lambda_0 d\hat{\theta} + \lambda_1) = -(d\lambda_1 - n\lambda_0) = 0,$$

i.e. $\tau_n(\omega_{-n})$ is exotic equivariant closed (in this case equivariant closed).

If $n \neq 0$, the homotopy (2.37) shows that

$$d\left(\frac{1}{n}\lambda_1 e^{-2\pi i n \theta}\right) = (\lambda_0 + \lambda_1 d\theta)e^{-2\pi i n \theta} = \omega_{-n},$$

i.e. ω_{-n} is $(d + H)$ -exact (in this case d -exact).

One can similarly do the odd degree case.

REFERENCES

- [1] N. Berline, E. Getzler and M. Vergne, Heat kernels and Dirac operators, Grundlehren Math. Wiss., **298**, Springer-Verlag, New York, 1992. MR1215720
- [2] J-M. Bismut, Index theorem and equivariant cohomology on the loop space. Comm. Math. Phys. **98** (1985), no. 2, 213–237. MR0786574
- [3] R. Bott and L. Tu, Differential forms in algebraic topology, Graduate Texts in Mathematics, **82**. Springer-Verlag, New York-Berlin, 1982. MR0658304
- [4] P. Bouwknegt, Lectures on Cohomology, T-duality and generalized geometry, Lect. Notes Phys. **807**, (2010) 261–311.
- [5] P. Bouwknegt, A. Carey, V. Mathai, M. Murray and D. Stevenson, Twisted K-theory and K-theory of bundle gerbes, Comm. Math. Phys. **228** (2002) 17–49, MR1911247 [arXiv:hep-th/0106194].

- [6] P. Bouwknegt, J. Evslin and V. Mathai, T-duality: Topology Change from H-flux, *Comm. Math. Phys.* **249** (2004) 383–415, MR2080959 [[arXiv:hep-th/0306062](#)].
- [7] ———, On the Topology and Flux of T-Dual Manifolds, *Phys. Rev. Lett.* **92** (2004) 181601, MR2116165 [[arXiv:hep-th/0312052](#)].
- [8] P. Bouwknegt and V. Mathai, Review of T-duality (in progress).
- [9] Jean-Luc Brylinski, *Loop spaces, characteristic classes and geometric quantization*, Progress in Mathematics, **107**, Birkhauser Boston, Inc., Boston, MA, 1993. MR1197353
- [10] G. Cavalcanti and M. Gualtieri, Generalized complex geometry and T-duality. A celebration of the mathematical legacy of Raoul Bott, 341–365, CRM Proc. Lecture Notes, **50**, Amer. Math. Soc., Providence, RI, 2010. MR2648900
- [11] M. Gualtieri, Generalized complex geometry, Ph.D thesis, University of Oxford, 2003, [[arXiv:math.DG/0401221](#)]
- [12] ———, Generalized complex geometry, *Ann. of Math. (2)* **174** (2011), no. 1, 75–123. MR2811595
- [13] F. Han and V. Mathai, Exotic twisted equivariant cohomology of loop spaces, twisted Bismut-Chern character and T-duality, *Comm. Math. Phys.*, **337**, no. 1, (2015) 127–150. MR3324158 [[arXiv:1405.1320](#)].
- [14] F. Han and V. Mathai, work in progress.
- [15] K. Hori, S. Katz, A. Klemm, R. Pandharipande, R. Thomas, C. Vafa, R. Vakil, E. Zaslow, Mirror symmetry. Clay Mathematics Monographs, American Mathematical Society, Providence, RI; Clay Mathematics Institute, Cambridge, MA, 2003. MR2003030
- [16] V. Mathai and D. Quillen, Superconnections, Thom classes, and equivariant differential forms. *Topology* **25** no.1 (1986) 85–110. MR0836726
- [17] V. Mathai and D. Stevenson, Chern character in twisted K-theory: equivariant and holomorphic cases, *Comm. Math. Phys.* **236** (2003) 161–186. MR1977885 [[arXiv:hep-th/0201010](#)].
- [18] R. Rohm and E. Witten, The antisymmetric tensor field in superstring theory. *Ann. Physics* **170** (1986), no. 2, 454–489. MR0851628

DEPARTMENT OF MATHEMATICS, NATIONAL UNIVERSITY OF SINGAPORE, SINGAPORE 119076
E-mail address: mathanf@nus.edu.sg

SCHOOL OF MATHEMATICAL SCIENCES, UNIVERSITY OF ADELAIDE, ADELAIDE 5005, AUSTRALIA
E-mail address: mathai.varghese@adelaide.edu.au