



Article

The Association between Dietary Intake, Asthma, and PCOS in Women from the Australian Longitudinal Study on Women's Health

Jessica A Grieger ^{1,2,*} , Allison Hodge ^{3,4} , Gita Mishra ⁵, Anju E Joham ^{6,7} and Lisa J Moran ^{1,6}

¹ Robinson Research Institute, University of Adelaide, North Adelaide, SA 5005, Australia; Lisa.Moran@monash.edu

² Adelaide Medical School, University of Adelaide, Adelaide, SA 5005, Australia

³ Cancer Epidemiology Division, Cancer Council Victoria, Melbourne, VCT 3004, Australia; Allison.Hodge@cancervic.org.au

⁴ Centre for Epidemiology and Biostatistics, Melbourne School of Population and Global Health, University of Melbourne, Melbourne, VCT 3010, Australia

⁵ School of Public Health, Faculty of Medicine, The University of Queensland, Brisbane, QLD 4072, Australia; g.mishra@sph.uq.edu.au

⁶ Monash Centre for Health Research and Implementation, School of Public Health and Preventive Medicine, Monash University, Clayton, VCT 3168, Australia; Anju.Joham@monash.edu

⁷ Diabetes and Vascular Medicine, Monash Health, Clayton, VCT 3168, Australia

* Correspondence: Jessica.grieger@adelaide.edu.au

Received: 18 December 2019; Accepted: 12 January 2020; Published: 15 January 2020



Abstract: Dietary intake potentially modifies the prevalence or severity of asthma. The prevalence of asthma is higher in women with polycystic ovary syndrome (PCOS); it is not known if diet confounds or modifies the association between asthma and PCOS. The aims of this study were: (i) To determine if the association of PCOS and asthma is independent of dietary pattern and (ii) to determine if dietary pattern modifies the association between PCOS and asthma. Women in this study were from the Australian Longitudinal Study on Women's Health (ALSWH) cohort born between 1973 to 1978 and aged 18 to 23 years ($n = 7382$). Logistic regression was used to assess the association between PCOS and asthma, adjusting for the following: (i) Potential confounders identified a priori and (ii) dietary patterns (z-score) identified by principle component analysis. In the adjusted analysis, women with PCOS were more likely to have asthma than the women without PCOS (OR 1.35 and 95% CI, 1.02 and 1.78). This relationship was not altered by further adjustment for dietary patterns (non-core food, meats and takeaway, or Mediterranean-style pattern). In the interaction analysis, only the women consuming less than the median intake of non-core foods (i.e., lower intake of discretionary or unhealthy foods) and with PCOS were more likely to have asthma (OR 1.91 and 95% CI, 1.29 and 2.82). Dietary intake did not confound the relationship between PCOS and asthma. Other mechanistic pathways are likely responsible for the asthma and PCOS association, and further studies assessing factors such as oral contraceptive use and sex steroid hormones warrant investigation.

Keywords: asthma; polycystic ovary syndrome; dietary patterns; diet; non-core foods; women

1. Introduction

Polycystic ovary syndrome (PCOS) is an endocrine disorder with a reported overall prevalence rate of 6% to 10% [1] based on the Rotterdam diagnostic criteria of two of three features, i.e., oligo- or anovulation, hyperandrogenism (clinical or biochemical), and polycystic ovaries on ultrasound, after exclusion of other causes [2]. PCOS is associated with reproductive (hyperandrogenism,

anovulation, infertility, and pregnancy complications) [3] and metabolic features (elevated prevalence of and risk factors for type 2 diabetes and cardiovascular disease) [4,5], along with psychological impact, such as increased risk of depression and anxiety [6] and impaired quality of life [7]. Insulin resistance is a key pathophysiological contributor to PCOS. Intrinsic insulin resistance is present in PCOS, with cases being more insulin resistant as compared with women of similar weight without PCOS [8]. Excess body weight exacerbates insulin resistance and the severity of PCOS symptoms [9]. Inflammation is also proposed as a pathophysiological contributor to PCOS [10]. This is not only as a consequence of elevated obesity, as women with PCOS have higher levels of proinflammatory markers such as C-reactive protein than BMI-matched controls [11–13]. Conversely, other studies have revealed that women who have PCOS do not have higher levels of inflammation than women without PCOS [14,15].

Asthma is a heterogeneous and complex disease, involving alterations in airway structure and airway inflammation and increased systemic inflammation [16]. It is one of the most common medical conditions with a prevalence of up to 38% in children [17] but only around 4% in adults [18]. Furthermore, a higher prevalence of asthma has been documented in women with PCOS from international studies including Danish [19,20] and Australian populations [21–23] which is hypothesized to be related to the elevated inflammation associated with PCOS. Diet can also modify the presentation of asthma through mechanisms including pro or anti-inflammatory effects. Poor dietary intake, such as foods high in saturated fats and glucose, contribute to inflammation through pathways involved in oxidative stress and insulin resistance, while foods high in antioxidants (such as fruits and vegetables) or unsaturated fatty acids (such as fish and nuts) are anti-inflammatory [24,25]. In keeping with these dietary effects on inflammation, a high fat/sugar/takeaway dietary pattern was associated with an increased likelihood of uncontrolled asthma in pregnancy [26]; and a high fat intake increased neutrophilic airway inflammation in subjects with asthma [27]. Higher dietary antioxidant intake was associated with improved lung function and asthma control (reviewed in [28]). In adults with asthma, those who consumed a low antioxidant diet for 14 days, were twice as likely to have an exacerbation at any time as compared with those consuming a high antioxidant diet [29].

It is unclear whether the association between asthma and PCOS is independent of diet given that both asthma and PCOS are associated with chronic low-grade inflammation. It is also unclear whether better dietary intake such as an anti-inflammatory diet could reduce the risk of asthma in women with PCOS. Determining this has important implications for understanding the pathophysiology of asthma in PCOS. We have previously examined dietary patterns in the Australian Longitudinal Study on Women's Health (ALSWH) and identified a range of potentially anti-inflammatory (Mediterranean-style) and proinflammatory (non-core foods or meats and takeaway) dietary patterns [30]. The current analysis will build on this work to assess the following hypotheses: (i) Dietary intake confounds the relationship between asthma and PCOS, such that a more proinflammatory dietary pattern increases the risk of both PCOS and asthma and (ii) dietary intake is an effect modifier on the association between PCOS and asthma, such that following a healthier diet after a diagnosis of PCOS could reduce the risk of asthma.

2. Methods

2.1. Study Population

This study is based on ALSWH, a longitudinal population-based study of four age cohorts of Australian women. This analysis was conducted using data from women in the young cohort of ALSWH, aged 18 to 23 years at ALSWH commencement in 1996. Women were randomly selected from the national health insurance scheme (Medicare) database with intentional oversampling from rural and remote areas [31]. Further methodological details have been reported elsewhere [32,33]. The Human Research Ethics Committees of the University of Newcastle, Australia, and the University of Queensland, Australia, approved the study and informed written consent was obtained from each participant. This analysis comprised women at Survey 5 (in 2009) (aged 31 to 36 years), as PCOS status

was only identified in Surveys 4 and 5, and food frequency questionnaire (FFQ) data was only collected in Surveys 3 and 5. We analyzed data from $n = 7382$ women who responded to the question on PCOS diagnosis in Survey 4 or 5 (responded yes to ("In the last three years have you been diagnosed with or treated for polycystic ovary syndrome")), who responded to the question on asthma at Survey 5 ("In the last three years have you been diagnosed with or treated for asthma"), and who had dietary data available for analysis ($n = 550$ PCOS and $n = 6832$ non-PCOS). No specific inclusion or exclusion criteria were applied, and all women were included, irrespective of pregnancy, medication, or country of birth. At Survey 5, the women sampled were broadly representative of the Australian population when compared with the census data [34].

2.2. Anthropometric and Demographic Variables

Information on self-reported demographic characteristics and anthropometric outcomes (height, weight, and waist circumference) were collected. Body mass index (BMI) was calculated using height and weight and respondents were categorized according to the World Health Organization guidelines for underweight (BMI < 18.5 kg/m²), normal weight (BMI 18.5 to 24.9 kg/m²), overweight (BMI 25 to 29.9 kg/m²), and obese (BMI ≥ 30 kg/m²) [35]. Demographic variables including country of birth, parity (no children or ≥1 child), education (no formal or high school education, diploma, trade, or degree), occupation (no paid job, clerical, trade, associated professional, or professional), personal income (no income, low income (\$1 to 36,399 annually), medium income (\$36,400 to 77,999 annually) and high income (\$ ≥ 78,000 annually), marital status (never married, separated/divorced/widowed, or married/de facto), smoking status (no defined as never smoker or ex-smoker, or yes defined as any current smoking) and combined oral contraceptive pill use (yes/no) were collected at Survey 5 with the exception of country of birth which was obtained at Survey 1.

2.3. Dietary Intake Variables

Self-reported diet data were collected from the Dietary Questionnaire for Epidemiological Studies (DQES) Version 2. This is a FFQ developed by The Cancer Council of Victoria which was previously validated in young to middle-aged (16 to 48 year old) Australian women in comparison with a seven-day weighed food record [36]. Participants who had incomplete FFQ data (>10% of items with missing responses) or those who reported a daily energy intake of >14,700 kJ/day or <2100 kJ/day were excluded from the analyses. Intakes of 100 different foods (grams per day) were obtained from the FFQ and were assigned into 33 food groups (grams per day) for use in the dietary pattern analysis. As previously described [30], dietary patterns were derived using factor analysis with factor loadings extracted using the principal component method and varimax/orthogonal rotation. The number of dietary patterns identified was based on eigenvalues >1.5, on identification of a break point in the scree plot, and on interpretability [37]. Using these criteria, a three-factor solution was chosen and rerun with the resulting factor scores saved and converted to z-scores for analysis. Items with factor loadings ≥0.25 were considered as the items of relevance for the identified factor. These items represent the foods most highly related to the identified factor [38]. Foods that cross-loaded on several factors were retained.

2.4. Statistical Analysis

All statistical procedures were performed using Stata version 14.2 (StataCorp, 146 College Station, TX, USA). Frequencies and descriptive statistics were expressed as n (%) and as mean ± SD, respectively. All reported p values were two-tailed, and a p value <0.05 was considered to be statistically significant. Data were analyzed by independent t -test to compare continuous variables and Chi-square test to compare categorical variables between women with and without PCOS. Binary logistic regression analyses were used to assess the association between the dependent variable of asthma and the independent variable PCOS. Then, multivariable analyses were conducted adjusting for potential confounders identified a priori, including BMI, personal income, occupation, education, age, parity, smoking, marital status, country of birth, and oral contraceptive pill use. Additional multivariable

analyses were then conducted adjusting for each dietary pattern (z-score) both in separate models for each z-score and in one model with all three z-scores included. Models were constructed to avoid collinearity and assessed for standard assumptions. Interaction effects between PCOS status and each dietary pattern (assessed as a categorical variable of above or below the median of the z-score) were assessed in the final models. Where interaction effects were significant, multivariable regression analyses were performed assessing the association between asthma as the dependent variable and PCOS status as the independent variable stratified for above and below the relevant median dietary pattern z-score.

3. Results

Participant characteristics are reported in Table 1. The prevalence of asthma was higher in women with PCOS as compared with those without PCOS (14.0 vs. 10.0%, $p = 0.004$). Women with PCOS had a higher BMI, weight and waist circumference, were less likely to have ≥ 1 child, less likely to be using the oral contraceptive pill, but more likely to have asthma.

Table 1. Characteristics for women with and without polycystic ovary syndrome (PCOS).

	Non-PCOS <i>n</i> = 6832	PCOS <i>n</i> = 550	<i>p</i> -Value
Age (years)	33.7 \pm 1.5	33.5 \pm 1.4	0.005
BMI (kg/m ²)	25.6 \pm 5.8	28.5 \pm 7.4	<0.001
Underweight	183 (2.8)	11 (2.1)	<0.001
Healthy weight	3532 (53.5)	205 (38.7)	
Overweight	1677 (25.4)	119 (22.5)	
Obese	1216 (18.4)	195 (36.8)	
Weight (kg)	70.8 \pm 16.6	78.3 \pm 21.6	<0.001
Waist circumference (cm)	86.2 \pm 14.1	91.2 \pm 17.2	<0.001
Smoking (yes)	989 (14.5)	74 (13.5)	0.436
Personal income			0.241
No income	604 (9.6)	55 (10.9)	
Low (\$1–36,399/y)	2592 (41.1)	200 (39.5)	
Medium (\$36,400–77,999/y)	2215 (35.2)	166 (32.8)	
High (\$ \geq 78,000/y)	890 (14.1)	85 (16.8)	
Education			0.658
No formal/high school	1398 (20.9)	104 (19.5)	
Trade/diploma	1790 (26.7)	140 (26.2)	
Degree	3513 (52.4)	290 (54.3)	
Occupation			0.788
No paid job	1425 (21.2)	108 (20.1)	
Clerical trade	1163 (17.3)	108 (20.1)	
Associate professional	1230 (18.3)	93 (17.3)	
Professional	2892 (43.1)	242 (50.0)	
Marital status			0.549
Never married	1178 (17.3)	103 (18.8)	
Separated/divorced/widowed	373 (5.5)	26 (4.7)	
Married/de facto	5262 (77.2)	419 (76.5)	
Parity (≥ 1)	4345 (63.6)	320 (58.2)	0.011
Born in Oceania	6437 (94.8)	510 (93.2)	0.116
OCP use (yes)	1581 (23.4)	96 (17.6)	0.002
Asthma	688 (10.0)	77 (14.0)	0.004

Data are presented as mean \pm SD or *n* (%) and were analyzed by independent *t*-test (continuous variables) or Chi-square test (categorical variables). BMI, body mass index; OCP, oral contraceptive pill; and PCOS, polycystic ovary syndrome.

The dietary intakes for women with and without PCOS are reported in Table 2. Women with PCOS had higher intakes of energy and fibre and a lower glycemic index, but differences were small.

Using principal components analysis, three dietary patterns were identified that explained 27% of the variance in food intake. These were non-core foods with high factor loadings for cakes, biscuits, sweet pastries; confectionary, refined grains and also takeaway foods and crisps; meats and takeaway with high factor loadings for fish (fried, processed, canned and cooked), processed meat, red meat, and takeaway food; and Mediterranean-style with high positive factor loadings for a variety of vegetables, fruit and nuts, and fish and inverse factor loadings for crisps. In an unadjusted analysis, women with PCOS had higher scores for all three dietary patterns, indicating a greater likelihood of following them.

Table 2. Dietary intake in women with and without PCOS.

Dietary Component/Pattern	Non-PCOS <i>n</i> = 6832	PCOS <i>n</i> = 550	<i>p</i> -Value
Energy (kJ)	6746 ± 2262	6994 ± 2485	0.014
Carbohydrate (% energy)	40.3 ± 5.7	40.3 ± 5.8	0.963
Protein (% energy)	20.9 ± 3.3	21.1 ± 3.3	0.379
Fat (% energy)	37.0 ± 4.9	36.8 ± 5.1	0.566
Saturated fat (% energy)	15.4 ± 3.1	15.3 ± 3.2	0.273
Monounsaturated fat (% energy)	13.1 ± 2.1	13.1 ± 2.2	0.929
Polyunsaturated fat (% energy)	5.1 ± 1.6	5.1 ± 1.6	0.837
Glycemic load	86.9 ± 33.5	89.1 ± 35.7	0.136
Glycemic index	50.8 ± 4.0	50.4 ± 4.0	0.017
Fibre (grams)	19.1 ± 7.0	20.0 ± 7.9	0.002
Alcohol (grams)	9.4 ± 13.5	8.3 ± 13.3	0.081
Non-core foods dietary pattern	−0.03 ± 0.92	0.06 ± 1.00	0.035
Meats and take away dietary pattern	−0.04 ± 0.81	0.05 ± 0.89	0.008
Mediterranean style dietary pattern	−0.004 ± 0.99	0.12 ± 1.04	0.007

Data are presented as mean ± SD and were analyzed by independent *t*-test (continuous variables). PCOS, polycystic ovary syndrome.

In the unadjusted analysis, women with PCOS were more likely to have asthma than those without PCOS (OR 1.45 and 95% CI, 1.12 and 1.87) (Table 3). The association was retained in adjusted analysis (OR 1.35 and 95% CI, 1.02 and 1.78) (Table 3). When including the dietary patterns in separate logistic regression models, the relationship between asthma and PCOS was not altered. There was also no independent relationship between the non-core foods (OR 1.06 and 95% CI, 0.97 and 1.17), meats and takeaway (OR 0.95 and 95% CI, 0.85 and 1.05), or Mediterranean-style dietary pattern scores (OR 0.94 and 95% CI, 0.86 to 1.03) and asthma in separate adjusted models.

Table 3. Logistic regression examining the association between PCOS and asthma.

Model	OR (95% CI)
Unadjusted	1.45 (1.12, 1.87)
Adjusted ^a	1.35 (1.02, 1.78)
Adjusted ^a , Non-core foods dietary pattern	1.34 (1.01, 1.78)
Adjusted ^a , Meats and takeaway dietary pattern	1.35 (1.02, 1.79)
Adjusted ^a , Mediterranean-style dietary pattern	1.36 (1.02, 1.80)
Adjusted ^a , all three dietary patterns	1.36 (1.03, 1.80)

Data are presented as odds ratio, 95% confidence interval, and *p* values and were analyzed using logistic regression ^a, Adjusted for BMI (WHO categories), income, occupation, education, age, parity, smoking, marital status, country of birth, and oral contraceptive pill use.

The interaction between PCOS and the dietary pattern scores was assessed for asthma. There was no interaction between PCOS and the meats and takeaway (*p* = 0.449) or the Mediterranean-style dietary pattern (*p* = 0.917). There was an interaction between PCOS and the non-core foods dietary pattern (*p* = 0.015). There was an association between asthma and PCOS for women with a non-core

foods dietary pattern z-score below the median (-0.15) (OR 1.91 and 95% CI, 1.29 and 2.82) but not above the median (OR 0.95 and 95% CI, 0.63 and 1.43). The relationship between asthma and PCOS was only observed for women with an overall healthier diet with lower intake of discretionary foods. The gram intake of food groups (cakes, confectionary, refined grains, vegemite, takeaway, crisps, juice, tomato sauce, processed meat, red meat, added sugar, and wholegrains) that scored ≥ 0.25 factor loading for the non-core foods dietary pattern are listed in Table 4 for the above and below median groups. Significant differences in median intake were found for all foods in the non-core foods dietary pattern, with the largest percentage difference for intake of cakes, confectionary, juice, crisps, and condiments including vegemite.

Table 4. Food group intake for the non-core foods dietary patterns.

Food	Factor Loading ^a	Below Median (g/d)	Above Median (g/d)	Mean Difference (g/d, 95% CI ^b	% Difference from below to above Median Intakes
Cakes, biscuits, sweet pastries (g/day)	0.661	10.2 \pm 10.2	31.4 \pm 29.0	−21.2 (−22.2, −20.2)	308% higher
Confectionary (g/day)	0.629	12.4 \pm 12.0	32.6 \pm 27.9	−20.2 (−21.2, −19.2)	263% higher
Refined grains (g/day)	0.483	74.8 \pm 51.4	128.5 \pm 78.1	−53.7 (−56.7, −50.6)	174% higher
Vegemite (g/day)	0.483	0.88 \pm 1.1	2.2 \pm 2.3	−1.3 (−1.4, −1.2)	250% higher
Takeaway (g/day)	0.467	30.5 \pm 22.6	53.2 \pm 37.5	−22.7 (−24.1, −21.3)	174% higher
Crisps (g/day)	0.466	14.1 \pm 14.6	28.7 \pm 26.6	−14.6 (−15.6, −13.6)	204% higher
Juice (g/day)	0.408	36.1 \pm 54.5	90.3 \pm 116.5	−54.3 (−58.4, −50.1)	250% higher
Tomato sauce (g/day)	0.380	1.8 \pm 1.7	3.5 \pm 3.5	−1.7 (−1.8, −1.6)	194% higher
Processed meat (g/day)	0.359	17.7 \pm 16.2	28.5 \pm 24.2	−10.9 (−11.8, −9.9)	161% higher
Red meat (g/day)	0.330	54.1 \pm 44.2	76.7 \pm 53.9	−22.6 (−24.8, −20.3)	142% higher
Added sugar (g/day)	0.325	8.8 \pm 9.2	14.7 \pm 12.8	−5.8 (−6.4, −5.3)	167% higher
Wholegrains (g/day)	0.319	74.9 \pm 58.6	115.5 \pm 90.7	−40.6 (−44.1, −37.1)	154% higher

^a, Foods only included with factor loading >0.25 and ^b, all $p > 0.001$ and were analyzed using independent t -test. Data are presented as mean \pm SD or mean difference (95% confidence interval).

4. Discussion

In a community cohort of Australian women aged 18 to 33 years, we confirm prior reports of a higher prevalence of asthma for women with PCOS than without, with a 1.35 higher likelihood of asthma with PCOS, or an absolute difference of 4% prevalence. We extend our previous research to also report that this higher prevalence of asthma was maintained after adjusting for dietary intake, specifically dietary patterns high in non-core foods, meats and takeaway, or a Mediterranean-style dietary pattern. These dietary patterns were not independently associated with asthma. However, we report for the first time an interaction between PCOS and the non-core foods dietary pattern score, such that the relationship between PCOS and asthma was only observed for women with a lower score on the non-core foods dietary pattern, indicating an overall healthier diet with lower intake of discretionary foods.

Our results confirm the literature that has reported that PCOS is independently associated with asthma [20,23,39]. The relationship between asthma and PCOS is likely to be multifactorial, but there are mechanisms proposed that could explain the association. Higher circulating C-reactive protein, interleukin 6, and tumor necrosis factor alpha have been reported in PCOS which, through various mechanisms, can contribute to adipocyte dysfunction, insulin resistance, and type 2 diabetes [10,40]. Compared to women without PCOS, women with PCOS have higher free testosterone and ratio of free testosterone to free estradiol [41], and reproductive health issues including irregular menstrual periods and subfertility [7]. Similar to PCOS, asthma is proinflammatory [42]. Similar to women with PCOS, women with asthma more often have abnormal levels of progesterone and estradiol [43], experience irregular menstruation [44], and have a longer time to achieve a pregnancy [45] than women without asthma. Insulin resistance also has been shown to be a risk factor for allergic asthma in obese children and adolescents [46,47]. Although our study was based on self-reported information for PCOS and asthma, and we cannot disentangle the reproductive and metabolic disturbances for the different

clinical PCOS phenotypes, common features of inflammation, insulin resistance, and altered hormones could have a role in the causation of asthma and allergy as well as in PCOS.

Unhealthy diets are associated with higher levels of inflammation [24,25]. Studies in adolescents or children have reported a higher risk for severe asthma in those consuming fast food three or more times per week [48] or with a low intake of fruits and vegetables, fish, butter, and dairy fat [49]. In adults, a high fish and fruit diet was related to a lower incidence of asthma [50]. Additionally, studies in adults with asthma report that those consuming higher fat and lower fiber intakes had lower forced expiratory volume in 1 s and airway eosinophilia [51]. A high-fat mixed meal increased sputum neutrophils 4 h post meal [27] but also resulted in increased airway inflammation and activation of a number of genes in sputum involved in immune system processes [52]. The literature, therefore, generally indicates that a healthier diet is associated with both a lower prevalence of asthma and better management of the disease. However, we did not find an association between any of the dietary pattern scores and asthma in this population. Controlling for these dietary patterns also did not alter the association between asthma and PCOS. These results suggest dietary intake does not confound the relationship between PCOS and asthma. Other exposures, such as OCP and sex steroid hormones, which are also associated with asthma [53] and PCOS [54,55], could play a larger role in the association between asthma and PCOS.

The literature is more contradictory with regards to the association between diet and PCOS. A number of studies have reported poorer dietary intake for women with PCOS including greater intakes of high glycemic index foods, fat, animal fat, saturated fat or cheese intake, and less fibre intake [56–59]. Conversely, we previously reported better diet quality, lower saturated fat and glycemic index, and higher fibre intake in women as compared with those without PCOS from ALSWH [60]. These discrepant findings may relate to a poorer quality diet pathophysiologically contributing to PCOS through weight gain or other mechanisms, such as worsening insulin resistance, or inflammation. Alternatively, a diagnosis of PCOS may lead to improved diet quality following health professional advice or motivation to change.

In this study, women with PCOS were more likely to follow all of the different dietary patterns with either a range of unhealthier foods in the non-core foods or meats and takeaway patterns consistent with a proinflammatory diet [24,25], or healthier foods in the Mediterranean-style dietary pattern consistent with an anti-inflammatory diet [61]. Interestingly however, we have conducted similar analyses using the Dietary Inflammatory Index as the dietary exposure but found no association with PCOS or asthma (unpublished). However, when we assessed the interactions between PCOS status and each dietary pattern, we found effect modification only for the non-core foods dietary pattern. Specifically, for women with a score below the median for this pattern, indicative of a healthier diet, the association between PCOS and asthma was observed. For women with a higher score there was no association between PCOS and asthma. These results are unexpected and lead us to reject our hypothesis that following a healthier diet after a diagnosis of PCOS reduces the risk of asthma. This result could indicate behavior change bias which can occur where women adopt behaviors perceived as healthier following diagnosis of a clinical condition such as asthma or PCOS, in association with clinician feedback or self-seeking of information for management. In this study, our prior findings of higher diet quality in community-based women with PCOS as compared with those without PCOS [30,62], may be further augmented by women with two clinical conditions, namely PCOS and asthma, having an even healthier diet higher in core foods and low in discretionary foods. This is consistent with the limited literature on dietary intake studies in adults with multiple chronic diseases where adults with type 2 diabetes and obesity had better diet quality than those with only type 2 diabetes [63].

The strengths of this study include the use of a large representative population-based cohort of young women. This allowed for the inclusion of varying degrees of obesity given that women with PCOS from clinical referrals have a higher prevalence of obesity than those from unselected populations [64]. Additionally, we considered a range of potential confounders in the relationship between PCOS and asthma. While the use of self-reported data for diet, asthma, PCOS, and anthropometry is a limitation,

this is reasonable for use in a large epidemiological study. The FFQ has also been validated for use in young Australian women [36] and the self-reported BMI, as used in this study, has been validated with anthropometric measurement among the mid-age cohort from ALSWH [65]. Furthermore, we have shown that self-reported PCOS status is strongly associated with menstrual irregularity, a key diagnostic criteria for PCOS, i.e., the odds ratio for having PCOS was over 10 for women often reporting irregular menstrual cycles [66]. Limitations include that we did not assess lung function to confirm asthma status. However, it is unlikely there was misclassification bias as women would know whether they have asthma or not, and the diet exposure is unlikely to impact the probability of misclassification [67]. The absence of lung function data also meant we could not assess associations according to asthma severity. The cross-sectional nature of this study means we cannot infer causation, and thus longitudinal studies that can assess changes in modifiable risk factors, such as diet, could allow stronger inferences.

5. Conclusions

In conclusion, dietary intake did not confound the relationship between PCOS and asthma, and contrary to our hypothesis, a healthier dietary pattern did not lower the risk of asthma in PCOS but rather increased it. Although a complex relationship exists between dietary factors and risk for PCOS and asthma, our findings do not suggest that advising women with PCOS to consume more healthy diets is likely to help reduce the risk of asthma, but rather a healthy diet should be followed to support a range of general health outcomes. Other mechanistic pathways are likely responsible for the asthma and PCOS association, and further studies assessing factors, including oral contraceptive use and sex steroid hormones, warrant investigation. Further studies are also warranted to investigate if different clinical phenotypes of PCOS relate to asthma. Understanding if metabolic or reproductive features associate most with asthma would support treatment recommendations to reduce asthma prevalence.

Author Contributions: All authors have read and agree to the published version of the manuscript. Conceptualization, L.J.M., G.M., and A.E.J.; methodology, L.J.M. and A.H.; formal Analysis, L.J.M. and J.A.G.; writing—original draft preparation, J.A.G.; writing—review and editing, all authors; funding acquisition, L.J.M.

Funding: This research was funded by the National Heart Foundation Future Leader Fellowship (L.M.), the NHMRC Early Career Fellowship (A.J.), and the NHMRC Principal Research Fellowship (G.M.).

Acknowledgments: The research on which this paper is based was conducted as part of the Australian Longitudinal Study on Women's Health undertaken by The University of Newcastle and The University of Queensland. We are grateful to the Australian Government Department of Health and Ageing for funding and to the women who provided the survey data. The authors thank Professor Graham Giles of the Cancer Epidemiology Centre of Cancer Council Victoria, for permission to use the Dietary Questionnaire for Epidemiological Studies (Version 2), Melbourne: Cancer Council Victoria, 1996.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Bozdag, G.; Mumusoglu, S.; Zengin, D.; Karabulut, E.; Yildiz, B.O. The prevalence and phenotypic features of polycystic ovary syndrome: A systematic review and meta-analysis. *Hum. Reprod.* **2016**, *31*, 2841–2855. [[CrossRef](#)] [[PubMed](#)]
2. Rotterdam ESHRE/ASRM-Sponsored PCOS Consensus Workshop Group. Revised 2003 consensus on diagnostic criteria and long-term health risks related to polycystic ovary syndrome (PCOS). *Hum. Reprod.* **2004**, *19*, 41–47. [[CrossRef](#)] [[PubMed](#)]
3. Bahri Khomami, M.; Joham, A.E.; Boyle, J.A.; Piltonen, T.; Silagy, M.; Arora, C.; Misso, M.L.; Teede, H.J.; Moran, L.J. Increased maternal pregnancy complications in polycystic ovary syndrome appear to be independent of obesity—A systematic review, meta-analysis, and meta-regression. *Obes. Rev.* **2019**, *20*, 659–674. [[CrossRef](#)] [[PubMed](#)]
4. Kakoly, N.S.; Khomami, M.B.; Joham, A.E.; Cooray, S.D.; Misso, M.L.; Norman, R.J.; Harrison, C.L.; Ranasinha, S.; Teede, H.J.; Moran, L.J. Ethnicity, obesity and the prevalence of impaired glucose tolerance and type 2 diabetes in PCOS: A systematic review and meta-regression. *Hum. Reprod. Update* **2018**, *24*, 455–467. [[CrossRef](#)] [[PubMed](#)]

5. Toulis, K.A.; Goulis, D.G.; Mintziori, G.; Kintiraki, E.; Eukarpidis, E.; Mouratoglou, S.A.; Pavlaki, A.; Stergianos, S.; Poulasouchidou, M.; Tzellos, T.G.; et al. Meta-analysis of cardiovascular disease risk markers in women with polycystic ovary syndrome. *Hum. Reprod. Update* **2011**, *17*, 741–760. [CrossRef] [PubMed]
6. Brutocao, C.; Zaiem, F.; Alsawas, M.; Morrow, A.S.; Murad, M.H.; Javed, A. Psychiatric disorders in women with polycystic ovary syndrome: A systematic review and meta-analysis. *Endocrine* **2018**, *62*, 318–325. [CrossRef]
7. Teede, H.J.; Misso, M.L.; Costello, M.F.; Dokras, A.; Laven, J.; Moran, L.; Piltonen, T.; Norman, R.J. International Evidencebased Guideline for the Assessment and Management of Polycystic Ovary Syndrome. 2018. Available online: https://jeanhailes.org.au/contents/documents/Resources/Tools/PCOS_evidence-based_guideline_for_assessment_and_management_pcos.pdf (accessed on 9 September 2019).
8. Stepto, N.K.; Cassar, S.; Joham, A.E.; Hutchison, S.K.; Harrison, C.L.; Goldstein, R.F.; Teede, H.J. Women with polycystic ovary syndrome have intrinsic insulin resistance on euglycaemic-hyperinsulaemic clamp. *Hum. Reprod.* **2013**, *28*, 777–784. [CrossRef]
9. Lim, S.S.; Davies, M.J.; Norman, R.J.; Moran, L.J. Overweight, obesity and central obesity in women with polycystic ovary syndrome: A systematic review and meta-analysis. *Hum. Reprod. Update* **2012**, *18*, 618–637. [CrossRef]
10. Shorakae, S.; Teede, H.; de Courten, B.; Lambert, G.; Boyle, J.; Moran, L.J. The Emerging Role of Chronic Low-Grade Inflammation in the Pathophysiology of Polycystic Ovary Syndrome. *Semin. Reprod. Med.* **2015**, *33*, 257–269. [CrossRef]
11. Bajuk Studen, K.; Pfeifer, M. Cardiometabolic risk in polycystic ovary syndrome. *Endocr. Connect.* **2018**, *7*, R238–R251. [CrossRef]
12. Samy, N.; Hashim, M.; Sayed, M.; Said, M. Clinical significance of inflammatory markers in polycystic ovary syndrome: Their relationship to insulin resistance and body mass index. *Dis. Markers* **2009**, *26*, 163–170. [CrossRef] [PubMed]
13. Ebejer, K.; Calleja-Agius, J. The role of cytokines in polycystic ovarian syndrome. *Gynecol. Endocrinol.* **2013**, *29*, 536–540. [CrossRef] [PubMed]
14. Lindholm, A.; Blomquist, C.; Bixo, M.; Dahlbom, I.; Hansson, T.; Sundstrom Poromaa, I.; Buren, J. No difference in markers of adipose tissue inflammation between overweight women with polycystic ovary syndrome and weight-matched controls. *Hum. Reprod.* **2011**, *26*, 1478–1485. [CrossRef] [PubMed]
15. Sanchez-Ferrer, M.L.; Prieto-Sanchez, M.T.; Corbalan-Biyang, S.; Mendiola, J.; Adoamnei, E.; Hernandez-Penalver, A.I.; Carmona-Barnosi, A.; Salido-Fierrez, E.J.; Torres-Cantero, A.M. Are there differences in basal thrombophilias and C-reactive protein between women with or without PCOS? *Reprod. Biomed. Online* **2019**, *38*, 1018–1026. [CrossRef] [PubMed]
16. Holgate, S.T. Innate and adaptive immune responses in asthma. *Nat. Med.* **2012**, *18*, 673–683. [CrossRef]
17. Asher, M.I.; Montefort, S.; Bjorksten, B.; Lai, C.K.; Strachan, D.P.; Weiland, S.K.; Williams, H.; The ISAAC Phase Three Study Group. Worldwide time trends in the prevalence of symptoms of asthma, allergic rhinoconjunctivitis, and eczema in childhood: ISAAC Phases One and Three repeat multicountry cross-sectional surveys. *Lancet* **2006**, *368*, 733–743. [CrossRef]
18. Papi, A.; Brightling, C.; Pedersen, S.E.; Reddel, H.K. Asthma. *Lancet* **2018**, *391*, 783–800. [CrossRef]
19. Zierau, L.; Cortes, R.; Thomsen, S.F.; Jimenez-Solem, E.; Lindenberg, S.; Backer, V. Asthma severity and fertility outcome in women with polycystic ovary syndrome: A registry-based study. *ERJ Open Res.* **2018**, *4*, 00138–2017. [CrossRef]
20. Glintborg, D.; Hass Rubin, K.; Nybo, M.; Abrahamsen, B.; Andersen, M. Morbidity and medicine prescriptions in a nationwide Danish population of patients diagnosed with polycystic ovary syndrome. *Eur. J. Endocrinol.* **2015**, *172*, 627–638. [CrossRef]
21. Hart, R.; Doherty, D.A. The potential implications of a PCOS diagnosis on a woman’s long-term health using data linkage. *J. Clin. Endocrinol. Metab.* **2015**, *100*, 911–919. [CrossRef]
22. Doherty, D.A.; Newnham, J.P.; Bower, C.; Hart, R. Implications of polycystic ovary syndrome for pregnancy and for the health of offspring. *Obstet. Gynecol.* **2015**, *125*, 1397–1406. [CrossRef] [PubMed]
23. Htet, T.D.; Teede, H.J.; de Courten, B.; Loxton, D.; Real, F.G.; Moran, L.J.; Joham, A.E. Asthma in reproductive-aged women with polycystic ovary syndrome and association with obesity. *Eur. Respir. J.* **2017**, *49*, 1601334. [CrossRef] [PubMed]

24. Biobaku, F.; Ghanim, H.; Batra, M.; Dandona, P. Macronutrient-Mediated Inflammation and Oxidative Stress: Relevance to Insulin Resistance, Obesity, and Atherogenesis. *J. Clin. Endocrinol. Metab.* **2019**, *104*, 6118–6128. [CrossRef] [PubMed]
25. Calder, P.C.; Ahluwalia, N.; Brouns, F.; Buetler, T.; Clement, K.; Cunningham, K.; Esposito, K.; Jonsson, L.S.; Kolb, H.; Lansink, M.; et al. Dietary factors and low-grade inflammation in relation to overweight and obesity. *Br. J. Nutr.* **2011**, *106*, S5–S78. [CrossRef]
26. Grieger, J.A.; Grzeskowiak, L.E.; Wood, L.G.; Clifton, V.L. Asthma control in pregnancy is associated with pre-conception dietary patterns. *Public Health Nutr.* **2016**, *19*, 332–338. [CrossRef]
27. Wood, L.G.; Garg, M.L.; Gibson, P.G. A high-fat challenge increases airway inflammation and impairs bronchodilator recovery in asthma. *J. Allergy Clin. Immunol.* **2011**, *127*, 1133–1140. [CrossRef]
28. Grieger, J.A.; Wood, L.G.; Clifton, V.L. Improving asthma during pregnancy with dietary antioxidants: The current evidence. *Nutrients* **2013**, *5*, 3212–3234. [CrossRef]
29. Wood, L.G. Diet, Obesity, and Asthma. *Ann. Am. Thorac. Soc.* **2017**, *14*, S332–S338. [CrossRef]
30. Moran, L.J.; Grieger, J.A.; Mishra, G.D.; Teede, H.J. The Association of a Mediterranean-Style Diet Pattern with Polycystic Ovary Syndrome Status in a Community Cohort Study. *Nutrients* **2015**, *7*, 8553–8564. [CrossRef]
31. Dobson, A.J.; Hockey, R.; Brown, W.J.; Byles, J.E.; Loxton, D.J.; McLaughlin, D.; Tooth, L.R.; Mishra, G.D. Cohort Profile Update: Australian Longitudinal Study on Women's Health. *Int. J. Epidemiol.* **2015**, *44*, 1547–1547f. [CrossRef]
32. Brown, W.J.; Bryson, L.; Byles, J.E.; Dobson, A.J.; Lee, C.; Mishra, G.; Schofield, M. Women's Health Australia: Recruitment for a national longitudinal cohort study. *Women Health.* **1998**, *28*, 23–40. [CrossRef] [PubMed]
33. Lee, C. *Women's Health Australia: Progress on the Australian Longitudinal Study on Women's Health 1995–2000*; Australian Academic Press Pty Ltd: Brisbane, Australia, 2001.
34. Lee, C.; Dobson, A.J.; Brown, W.J.; Bryson, L.; Byles, J.; Warner-Smith, P.; Young, A.F. Cohort Profile: The Australian Longitudinal Study on Women's Health. *Int. J. Epidemiol.* **2005**, *34*, 987–991. [CrossRef] [PubMed]
35. World Health Organisation. Obesity and Overweight: Fact sheet. Available online: <http://www.who.int/mediacentre/factsheets/fs311/en/> (accessed on 7 August 2019).
36. Hodge, A.; Patterson, A.J.; Brown, W.J.; Ireland, P.; Giles, G. The Anti Cancer Council of Victoria FFQ: Relative validity of nutrient intakes compared with weighed food records in young to middle-aged women in a study of iron supplementation. *Aust. N. Z. J. Public Health* **2000**, *24*, 576–583. [CrossRef] [PubMed]
37. Schulze, M.B.; Hoffmann, K.; Kroke, A.; Boeing, H. An approach to construct simplified measures of dietary patterns from exploratory factor analysis. *Br. J. Nutr.* **2003**, *89*, 409–419. [CrossRef]
38. Kline, P.K. *An Easy Guide to Factor Analysis*; Routledge: London, UK, 1994.
39. Zierau, L.; Gade, E.J.; Lindenberg, S.; Backer, V.; Thomsen, S.F. Coexistence of asthma and polycystic ovary syndrome: A concise review. *Respir. Med.* **2016**, *119*, 155–159. [CrossRef]
40. Teede, H.J.; Hutchison, S.; Zoungas, S.; Meyer, C. Insulin resistance, the metabolic syndrome, diabetes, and cardiovascular disease risk in women with PCOS. *Endocrine* **2006**, *30*, 45–53. [CrossRef]
41. Insenser, M.; Murri, M.; Del Campo, R.; Martinez-Garcia, M.A.; Fernandez-Duran, E.; Escobar-Morreale, H.F. Gut Microbiota and the Polycystic Ovary Syndrome: Influence of Sex, Sex Hormones, and Obesity. *J. Clin. Endocrinol. Metab.* **2018**, *103*, 2552–2562. [CrossRef]
42. Nadif, R.; Siroux, V.; Oryszczyn, M.P.; Ravault, C.; Pison, C.; Pin, I.; Kauffmann, F.; Epidemiological study on the Genetics and Environment of Asthma. Heterogeneity of asthma according to blood inflammatory patterns. *Thorax* **2009**, *64*, 374–380. [CrossRef]
43. Balzano, G.; Fuschillo, S.; Melillo, G.; Bonini, S. Asthma and sex hormones. *Allergy* **2001**, *56*, 13–20. [CrossRef]
44. Svanes, C.; Real, F.G.; Gislason, T.; Jansson, C.; Jogi, R.; Norrman, E.; Nystrom, L.; Toren, K.; Omenaas, E. Association of asthma and hay fever with irregular menstruation. *Thorax* **2005**, *60*, 445–450. [CrossRef]
45. Grzeskowiak, L.E.; Smithers, L.G.; Grieger, J.A.; Bianco-Miotto, T.; Leemaqz, S.Y.; Clifton, V.L.; Poston, L.; McCowan, L.M.; Kenny, L.C.; Myers, J.; et al. Asthma treatment impacts time to pregnancy: Evidence from the international SCOPE study. *Eur. Respir. J.* **2018**, *51*, 1702035. [CrossRef] [PubMed]
46. Cardet, J.C.; Ash, S.; Kusa, T.; Camargo, C.A., Jr.; Israel, E. Insulin resistance modifies the association between obesity and current asthma in adults. *Eur. Respir. J.* **2016**, *48*, 403–410. [CrossRef] [PubMed]

47. Sanchez Jimenez, J.; Herrero Espinet, F.J.; Mengibar Garrido, J.M.; Roca Antonio, J.; Penos Mayor, S.; Penas Boira Mdel, M.; Roca Comas, A.; Ballester Martinez, A. Asthma and insulin resistance in obese children and adolescents. *Pediatr. Allergy Immunol.* **2014**, *25*, 699–705. [[CrossRef](#)] [[PubMed](#)]
48. Ellwood, P.; Asher, M.I.; Garcia-Marcos, L.; Williams, H.; Keil, U.; Robertson, C.; Nagel, G.; The ISAAC Phase III Study Group. Do fast foods cause asthma, rhinoconjunctivitis and eczema? Global findings from the International Study of Asthma and Allergies in Childhood (ISAAC) phase three. *Thorax* **2013**, *68*, 351–360. [[CrossRef](#)]
49. Tricon, S.; Willers, S.; Smit, H.A.; Burney, P.G.; Devereux, G.S.; Frew, A.J. Nutrition and allergic disease. *Clin. Exp. Allergy Rev.* **2006**, *6*, 117–188. [[CrossRef](#)]
50. Uddenfeldt, M.; Janson, C.; Lampa, E.; Leander, M.; Norback, D.; Larsson, L.; Rask-Andersen, A. High BMI is related to higher incidence of asthma, while a fish and fruit diet is related to a lower- Results from a long-term follow-up study of three age groups in Sweden. *Respir. Med.* **2010**, *104*, 972–980. [[CrossRef](#)]
51. Berthon, B.S.; Macdonald-Wicks, L.K.; Gibson, P.G.; Wood, L.G. Investigation of the association between dietary intake, disease severity and airway inflammation in asthma. *Respirology* **2013**, *18*, 447–454. [[CrossRef](#)]
52. Li, Q.; Baines, K.J.; Gibson, P.G.; Wood, L.G. Changes in Expression of Genes Regulating Airway Inflammation Following a High-Fat Mixed Meal in Asthmatics. *Nutrients* **2016**, *8*, E30. [[CrossRef](#)]
53. Matheson, M.C.; Burgess, J.A.; Lau, M.Y.; Lowe, A.J.; Gurrin, L.C.; Hopper, J.L.; Giles, G.G.; Johns, D.P.; Walters, E.H.; Abramson, M.J.; et al. Hormonal contraception increases risk of asthma among obese but decreases it among nonobese subjects: A prospective, population-based cohort study. *ERJ Open Res.* **2015**, *1*, 00026–2015. [[CrossRef](#)]
54. Fauser, B.C.; Tarlatzis, B.C.; Rebar, R.W.; Legro, R.S.; Balen, A.H.; Lobo, R.; Carmina, E.; Chang, J.; Yildiz, B.O.; Laven, J.S.; et al. Consensus on women's health aspects of polycystic ovary syndrome (PCOS): The Amsterdam ESHRE/ASRM-Sponsored 3rd PCOS Consensus Workshop Group. *Fertil. Steril.* **2012**, *97*, 28–38. [[CrossRef](#)]
55. Nisenblatt, V.; Norman, R.J. Androgens and polycystic ovary syndrome. *Curr. Opin. Endocrinol. Diabetes Obes.* **2009**, *16*, 224–231. [[CrossRef](#)] [[PubMed](#)]
56. Douglas, C.C.; Norris, L.E.; Oster, R.A.; Darnell, B.E.; Azziz, R.; Gower, B.A. Difference in dietary intake between women with polycystic ovary syndrome and healthy controls. *Fertil. Steril.* **2006**, *86*, 411–417. [[CrossRef](#)] [[PubMed](#)]
57. Altieri, P.; Cavazza, C.; Pasqui, F.; Morselli, A.M.; Gambineri, A.; Pasquali, R. Dietary habits and their relationship with hormones and metabolism in overweight and obese women with polycystic ovary syndrome. *Clin. Endocrinol.* **2012**, *78*, 52–59. [[CrossRef](#)] [[PubMed](#)]
58. Colombo, O.; Pinelli, G.; Comelli, M.; Marchetti, P.; Sieri, S.; Brighenti, F.; Nappi, R.E.; Tagliabue, A. Dietary intakes in infertile women a pilot study. *Nutr. J.* **2009**, *8*, 53. [[CrossRef](#)]
59. Wild, R.A.; Painter, P.C.; Coulson, P.B.; Carruth, K.B.; Ranney, G.B. Lipoprotein lipid concentrations and cardiovascular risk in women with polycystic ovary syndrome. *J. Clin. Endocrinol. Metab.* **1985**, *61*, 946–951. [[CrossRef](#)]
60. Moran, L.J.; Brown, W.J.; McNaughton, S.A.; Joham, A.E.; Teede, H.J. Weight management practices associated with PCOS and their relationships with diet and physical activity. *Hum. Reprod.* **2017**, *32*, 669–678. [[CrossRef](#)]
61. Casas, R.; Sacanella, E.; Estruch, R. The Immune Protective Effect of the Mediterranean Diet against Chronic Low-grade Inflammatory Diseases. *Endocr. Metab. Immune Disord. Drug Targets* **2016**, *14*, 245–254. [[CrossRef](#)]
62. Moran, L.J.; Ranasinha, S.; Zoungas, S.; McNaughton, S.A.; Brown, W.J.; Teede, H.J. The contribution of diet, physical activity and sedentary behaviour to body mass index in women with and without polycystic ovary syndrome. *Hum. Reprod.* **2013**, *28*, 2276–2283. [[CrossRef](#)]
63. Mangou, A.; Grammatikopoulou, M.G.; Mirkopoulou, D.; Sailer, N.; Kotzamanidis, C.; Tsigga, M. Associations between diet quality, health status and diabetic complications in patients with type 2 diabetes and comorbid obesity. *Endocrinol. Nutr.* **2012**, *59*, 109–116. [[CrossRef](#)]
64. Ezech, U.; Yildiz, B.O.; Azziz, R. Referral bias in defining the phenotype and prevalence of obesity in polycystic ovary syndrome. *J. Clin. Endocrinol. Metab.* **2013**, *98*, E1088–E1096. [[CrossRef](#)]
65. Burton, N.W.; Brown, W.; Dobson, A. Accuracy of body mass index estimated from self-reported height and weight in mid-aged Australian women. *Aust. N. Z. J. Public Health* **2010**, *34*, 620–623. [[CrossRef](#)] [[PubMed](#)]

66. Teede, H.J.; Joham, A.E.; Paul, E.; Moran, L.J.; Loxton, D.; Jolley, D.; Lombard, C. Longitudinal weight gain in women identified with polycystic ovary syndrome: Results of an observational study in young women. *Obesity* **2013**, *21*, 1526–1532. [[CrossRef](#)] [[PubMed](#)]
67. Matsui, E.C.; Keet, C.A. Weighing the evidence: Bias and confounding in epidemiologic studies in allergy/immunology. *J. Allergy Clin. Immunol.* **2017**, *139*, 448–450. [[CrossRef](#)] [[PubMed](#)]



© 2020 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).