

Pedram Hekmati, Michael K. Murray, Richard J. Szabo, Raymond F. Vozzo

Sign choices for orientifolds

Communications in Mathematical Physics, 2020; 378(3):1843-1873

© Springer-Verlag GmbH Germany, part of Springer Nature 2020

*This is a post-peer-review, pre-copyedit version of an article published in **Communications in Mathematical Physics**. The final authenticated version is available online at:*

<http://dx.doi.org/10.1007/s00220-020-03831-z>

PERMISSIONS

<https://www.springer.com/gp/open-access/publication-policies/self-archiving-policy>

Self-archiving for articles in subscription-based journals

Springer journals' [policy on preprint sharing](#).

By signing the Copyright Transfer Statement you still retain substantial rights, such as self-archiving:

*Author(s) are permitted to self-archive a pre-print and an author's **accepted manuscript** version of their Article.*

.....

b. An Author's Accepted Manuscript (AAM) is the version accepted for publication in a journal following peer review but prior to copyediting and typesetting that can be made available under the following conditions:

(i) Author(s) retain the right to make an AAM of their Article available on their own personal, self-maintained website immediately on acceptance,

(ii) Author(s) retain the right to make an AAM of their Article available for public release on any of the following 12 months after first publication ("Embargo Period"): their employer's internal website; their institutional and/or funder repositories. AAMs may also be deposited in such repositories immediately on acceptance, provided that they are not made publicly available until after the Embargo Period.

An acknowledgement in the following form should be included, together with a link to the published version on the publisher's website: "This is a post-peer-review, pre-copyedit version of an article published in [insert journal title]. The final authenticated version is available online at: [http://dx.doi.org/\[insert DOI\]](http://dx.doi.org/[insert DOI])".

When publishing an article in a subscription journal, without open access, authors sign the Copyright Transfer Statement (CTS) which also details Springer's self-archiving policy.

See Springer Nature [terms of reuse](#) for archived author accepted manuscripts (AAMs) of subscription articles.

3 August 2021

<http://hdl.handle.net/2440/128445>

SIGN CHOICES FOR ORIENTIFOLDS

PEDRAM HEKMATI, MICHAEL K. MURRAY, RICHARD J. SZABO, AND RAYMOND F. VOZZO

ABSTRACT. We analyse the problem of assigning sign choices to O-planes in orientifolds of type II string theory. We show that there exists a sequence of invariant p -gerbes with $p \geq -1$, which give rise to sign choices and are related by coboundary maps. We prove that the sign choice homomorphisms stabilise with the dimension of the orientifold and we derive topological constraints on the possible sign configurations. Concrete calculations for spherical and toroidal orientifolds are carried out, and in particular we exhibit a four-dimensional orientifold where not every sign choice is geometrically attainable. We elucidate how the K -theory groups associated with invariant p -gerbes for $p = -1, 0, 1$ interact with the coboundary maps. This allows us to interpret a notion of K -theory due to Gao and Hori as a special case of twisted KR -theory, which consequently implies the homotopy invariance and Fredholm module description of their construction.

CONTENTS

1. Introduction	2
2. Sign choices for invariant p -gerbes	5
3. Real bundle gerbes and their connections	7
3.1. Real bundle gerbes	7
3.2. Connective structures on Real bundle gerbes	8
3.3. Holonomy of orientifold B -fields	10
4. Geometric constructions of sign choices	13
4.1. Sign choice for Real functions	13
4.2. Sign choice for equivariant bundles	13
4.3. Sign choice for Real bundle gerbes	14
5. Sign choices for spherical orientifolds	17
5.1. The three-sphere $S^{3,1}$	17
5.2. The n -sphere $S^{n,1}$	19
5.3. Sign choice for equivariant bundle 2-gerbes	19
5.4. Constraints on geometric sign choices	20
6. Sign choices for toroidal orientifolds	21
6.1. The circle $S^{1,1}$	21
6.2. The two-torus $S^{1,1} \times S^{1,1}$	22

The authors acknowledge support under the Australian Research Council's *Discovery Projects* funding scheme (project numbers DP120100106, DP130102578 and DP180100383), the Consolidated Grant ST/P000363/1 from the UK Science and Technology Facilities Council, the Marsden Foundation (project number 3713803), and the Action MP1405 QSPACE from the European Cooperation in Science and Technology (COST).

We thank David Baraglia and Jonathan Rosenberg for helpful discussions.

6.3. The three-torus $S^{1,1} \times S^{1,1} \times S^{1,1}$	23
7. KR -theory with sign choices	25
7.1. The Gao–Hori construction	25
7.2. Twisted K -theory	26
References	28

1. INTRODUCTION

In this paper we study the Real Brauer group and related structures on orientifolds, that is, pairs (M, τ) consisting of a manifold M equipped with an involution $\tau: M \rightarrow M$. The fixed point set of the involution is M^τ and its connected components will be called orientifold planes or O-planes for short. Orientifolds are more commonly referred to as Real manifolds in the mathematics literature; the present terminology is borrowed from string theory where these give backgrounds which are important for model building and understanding T-duality [1, 6].

O-planes can carry a positive or negative Ramond–Ramond charge, which for consistency of the string background must be cancelled by the inclusion of suitable D-brane charges; conversely, string backgrounds with D-branes require inclusion of suitable orientifold planes. The collection of discrete O-plane charges for a given orientifold background is specified mathematically by a sign choice, which is an assignment of an element ± 1 of \mathbb{Z}_2 to each O-plane, or equivalently a cocycle in $H^0(M^\tau, \mathbb{Z}_2)$. The sign choices determine the projection of the $U(n)$ gauge groups of the Chan–Paton factors when n coincident D-branes are placed on top of an O-plane: this is $O(n)$ for sign choice $+1$ and $Sp(n)$ for sign choice -1 .

A general condition for the allowed distributions of sign choices for a given orientifold (M, τ) is not presently known. A related open problem is to define a suitable variant of KR -theory classifying D-branes on orientifolds for an arbitrarily given sign choice which is not necessarily constant over the different O-planes. The K -theory and cohomology classification of the spectrum of O-plane charges for a given orientifold is discussed in [22, 12, 14, 19, 5, 3]. Real continuous trace C^* -algebras and their sign choices were discussed in [7], where a corresponding version of KR -theory was also proposed. The purpose of this paper is to provide some detailed answers to these questions from a geometric perspective.

In orientifolds of type II string theory, the behaviour of fields in the NS–NS sector influences the allowed sign choices. At the lowest level there is the dilaton field, which is a function on spacetime M . Next, a globally defined Neveu–Schwarz B -field plays the role of a magnetic field on a D-brane and so is naturally associated to a line bundle on spacetime. More generally, a B -field with non-vanishing H -flux is associated with a bundle gerbe on spacetime; a non-trivial 2-torsion H -flux on an O-plane reverses its sign choice, as will also follow by our geometric considerations below.

We therefore wish to consider the mathematical problem of assigning a sign choice to certain geometric objects over (M, τ) . For instance in the simplest case of a Real function, that is, a function $f: M \rightarrow U(1)$ which satisfies $f \circ \tau = \bar{f}$, we have $f(m) \in \mathbb{Z}_2$ at a fixed point $m \in M^\tau$ so it defines a sign choice. More generally, we show that there is a sequence of natural geometric objects on M to which sign choices can be allocated. We also address the fundamental question

of whether every sign choice can arise from one of these geometric objects, particularly those that encode the data of a string compactification.

Of particular interest, from the string theory perspective, will be the first three classes of these geometric objects: Real functions as discussed above, \mathbb{Z}_2 -equivariant line bundles, and Real bundle gerbes. These have characteristic classes in the equivariant sheaf cohomology groups $H^0(M; \mathbb{Z}_2, \mathcal{U}^1)$, $H^1(M; \mathbb{Z}_2, \mathcal{U}^0)$ and $H^2(M; \mathbb{Z}_2, \mathcal{U}^1)$, respectively, that we will discuss below. Each of these groups comes equipped with a *sign choice map* which is a homomorphism σ_i mapping into $H^0(M^\tau, \mathbb{Z}_2)$. There are connecting homomorphisms ∂_i between these cohomology groups giving rise to the commutative diagram

$$\begin{array}{ccc}
 H^0(M; \mathbb{Z}_2, \mathcal{U}^1) & & \\
 \downarrow \partial_1 & \searrow \sigma_0 & \\
 H^1(M; \mathbb{Z}_2, \mathcal{U}^0) & \xrightarrow{\sigma_1} & H^0(M^\tau, \mathbb{Z}_2) \\
 \downarrow \partial_2 & \nearrow \sigma_2 & \\
 H^2(M; \mathbb{Z}_2, \mathcal{U}^1) & & \\
 \downarrow \partial_3 & & \\
 \vdots & &
 \end{array} \tag{1.1}$$

As the vertical arrow ∂_3 suggests, this is part of a more general story. We denote by $\Sigma_i(M)$ the image of the sign choice homomorphisms σ_i in $H^0(M^\tau, \mathbb{Z}_2)$ and call such elements *geometric sign choices*. If M has dimension d , we find that these groups coincide in degrees $d-1$ onwards. As a result,

$$\Sigma_0(M) \subseteq \Sigma_1(M) \subseteq \cdots \subseteq \Sigma_{d-1}(M) = \Sigma_d(M) = \cdots \subseteq H^0(M^\tau, \mathbb{Z}_2) ,$$

and thus the geometric sign choices are precisely $\Sigma_{d-1}(M)$. We show that not all sign choices are geometric sign choices by exhibiting a four-manifold M where $\Sigma_3(M) \neq H^0(M^\tau, \mathbb{Z}_2)$.

In Section 5.4 we derive some constraints on the possible geometric sign choices, that is, on the images $\Sigma_i(M)$ for $i = 0, 1, 2$. For a Real bundle gerbe (P, Y) over a 2-connected orientifold M with vanishing complex Dixmier–Douady class $\text{DD}(P)$, we prove that the sign choice is generically constant. More generally, we show in Proposition 5.4 that the sign choice map σ_2 is completely determined by the periods of the complex Dixmier–Douady class and the sign at a single fixed point by the formula

$$\sigma_2(P)(m) = (-1)^{\langle \mu_{m,m'}^*(\text{DD}(P)), S^3 \rangle} \sigma_2(P)(m') ,$$

where the map $\mu_{m,m'}: S^3 \rightarrow M$ is constructed from the τ -action and the fixed points $m, m' \in M^\tau$. This kind of construction depends only on knowing the sign choices for spheres and the formula extends to all invariant p -gerbes under appropriate connectedness conditions on M . In particular for the sign choice homomorphisms σ_0 and σ_1 on connected respectively 1-connected orientifolds, the Dixmier–Douady class is substituted by the winding class respectively first Chern class.

We proceed to consider some concrete examples of orientifolds which appear in conventional string theory backgrounds. An important class of compactification spaces arising in string theory with background fluxes are spheres. Recall that $S^{p,q} \subset \mathbb{R}^{p+q}$ denotes the $p+q-1$ -dimensional sphere with the involution τ which changes the sign of the first p coordinates and leaves alone the remaining q coordinates. We also carry out explicit calculations involving tori, which play

an important role in flat compactifications of string theory and provide examples of multiply connected orientifolds. The results are summarised in the following table:

(M, τ)	$S^{3,1}$	$S^{1,1}$	$S^{1,1} \times S^{1,1}$	$S^{1,1} \times S^{1,1} \times S^{1,1}$
$\dim(M)$	3	1	2	3
$H^0(M^\tau, \mathbb{Z}_2)$	\mathbb{Z}_2^2	\mathbb{Z}_2^2	\mathbb{Z}_2^4	\mathbb{Z}_2^8
$\Sigma_0(M)$	\mathbb{Z}_2	\mathbb{Z}_2^2	\mathbb{Z}_2^3	\mathbb{Z}_2^4
$H^1(M; \mathbb{Z}_2, \mathcal{U}^0)$	\mathbb{Z}_2	\mathbb{Z}_2^2	$\mathbb{Z}_2^3 \oplus \mathbb{Z}$	$\mathbb{Z}_2^4 \oplus \mathbb{Z}^3$
$\Sigma_1(M)$	\mathbb{Z}_2	\mathbb{Z}_2^2	\mathbb{Z}_2^4	\mathbb{Z}_2^7
$H^2(M; \mathbb{Z}_2, \mathcal{U}^1)$	$\mathbb{Z}_2 \oplus \mathbb{Z}$	\mathbb{Z}_2^2	$\mathbb{Z}_2^3 \oplus \mathbb{Z}$	$\mathbb{Z}_2^7 \oplus \mathbb{Z}$
$\Sigma_2(M)$	\mathbb{Z}_2^2	\mathbb{Z}_2^2	\mathbb{Z}_2^4	\mathbb{Z}_2^8

Note that $\Sigma_2(M) = H^0(M^\tau, \mathbb{Z}_2)$ in these special cases. In particular, the three-dimensional examples show that, generally, for all sign choices to arise one must consider Real bundle gerbes which have non-vanishing complex Dixmier–Douady class in the Real Brauer group $H^2(M; \mathbb{Z}_2, \mathcal{U}^1)$. We also consider higher-dimensional spheres $S^{n,1}$ with $n \geq 4$, where the sign choice map σ_2 is not surjective. In the case of tori, our geometric sign choice groups $\Sigma_1(M)$ reproduce the classification of O-plane charges from [5] using the Borel equivariant cohomology group $H_{\mathbb{Z}_2}^2(M, \mathbb{Z}_2)$ to determine the allowed globally defined B -field configurations on M .

The possible sign choices for orientifolds were also considered from a geometric perspective by Gao and Hori in [9] using constructions based directly on type II string theory in backgrounds with topologically trivial H -flux. In Section 7 we show how to interpret their geometric data in terms of our constructions, and demonstrate that their version of KR -theory can be understood as being twisted by either a Real function or an equivariant line bundle. We show that in either case it is equivalent to the KR -theory twisted by a corresponding Real bundle gerbe constructed by repeated application of the connecting homomorphisms

$$H^0(M; \mathbb{Z}_2, \mathcal{U}^1) \xrightarrow{\partial_1} H^1(M; \mathbb{Z}_2, \mathcal{U}^0) \xrightarrow{\partial_2} H^2(M; \mathbb{Z}_2, \mathcal{U}^1) .$$

In these cases the KR -theory classes are twisted by a Real bundle gerbe with trivial complex Dixmier–Douady class, whereby the possible Real structures on the bundle gerbe are given by equivariant line bundles (it is a torsor under $H^1(M; \mathbb{Z}_2, \mathcal{U}^0)$). The spectrum of O-plane charges is then determined by the fact that, on the connected components of the fixed point set of the orientifold involution, the KR -theory becomes KO -theory when the restriction of the bundle gerbe has sign $+1$ and so is Real trivial, or KSp -theory when the restriction of the bundle gerbe has 2-torsion Dixmier–Douady class with sign -1 corresponding to non-trivial torsion H -flux [13]. However, in general there are sign choices that cannot be achieved in such a way unless the complex Dixmier–Douady class (or H -flux of the string background) is non-trivial,¹ in which case the more general geometric realisation of twisted KR -theory from [13] contains the appropriate sign choices.

¹This seems to be at odds with the result of [7] who used C^* -algebra methods to show that any sign choice could arise, whereas we have found constraints. This raises the possibility that the correspondence between continuous trace C^* -algebras and groupoids for the Real case is not a straightforward generalisation of the complex case.

A summary of the contents of this paper is as follows. In Section 2 we introduce invariant p -gerbes for $p \geq -1$ and define their associated sign choice homomorphisms at the level of cohomology. We derive some general properties of the sign choice maps and show that they stabilise at degree $\dim(M) - 1$. In Section 3 we restrict our attention to the case $p = 1$, and explain how to incorporate the orientifold B -field and its holonomy. As an application, we obtain sufficient conditions for the existence of Real structures on bundle gerbes when M is 2-connected. We proceed to give geometric constructions of the sign choice maps in Section 4 for Real functions, equivariant line bundles and Real bundle gerbes. We further derive a general characterisation of the kernel and image of the sign choice map σ_2 in terms of the sheaf cohomology of the space of “physical points” M/τ of the orientifold, which exhibits obstructions to the possible sign choices. In Sections 5 and 6 we carry out detailed calculations for some spherical and toroidal orientifolds. Under certain connectedness conditions on M , we obtain a geometric constraint on the sign choices in terms of the periods of the complex characteristic classes of the p -gerbes. Finally, in Section 7 we recast the orientifold data of Gao and Hori [9] in the language of the present paper and prove that their notion of *K-theory twisted by an equivariant line bundle L* is nothing but *KR-theory twisted by the Real bundle gerbe $\partial_2(L)$* under the connecting homomorphism in (1.1).

2. SIGN CHOICES FOR INVARIANT p -GERBES

In this section we develop some general theory of sign choices for *invariant p -gerbes*. This is followed in subsequent sections by a discussion of the cases of particular geometric interest. Let (M, τ) be an orientifold and let $M^\tau \subseteq M$ denote the fixed point set of the involution.

Definition 2.1. A *sign choice* is an element of the group $H^0(M^\tau, \mathbb{Z}_2)$.

In other words, a sign choice assigns to every O-plane, that is, every connected component of M^τ , a sign ± 1 in \mathbb{Z}_2 .

We are interested in various geometric objects that determine a sign choice on an orientifold. For instance, Real functions, equivariant line bundles and Real bundle gerbes form the first three examples of invariant p -gerbes corresponding to the cases $p = -1, 0, 1$. The case $p = 2$ would correspond to equivariant bundle 2-gerbes which we briefly consider in Section 5.3. For $p > 2$ the geometric theory of p -gerbes is not well understood, but we can work instead with the corresponding cohomology groups that classify them since the sign choice maps factorise through these groups.

There are two natural \mathbb{Z}_2 -sheaves on the orientifold which in [13] we denoted by $\mathcal{U}(1)$ and $\overline{\mathcal{U}}(1)$. They are the equivariant sheaves of smooth functions into $U(1)$, where the first is endowed with a trivial \mathbb{Z}_2 -action on $U(1)$ and the second has the \mathbb{Z}_2 -action induced by complex conjugation. In the present paper it will be useful to change notation and denote these by \mathcal{U}^ϵ with $\epsilon \in \mathbb{Z}_2 = \{0, 1\}$, where $\mathcal{U}^0 = \mathcal{U}(1)$ and $\mathcal{U}^1 = \overline{\mathcal{U}}(1)$. We will continue to use the symbol $\mathcal{U}(1)$ for the sheaf of smooth $U(1)$ -valued functions. With this notation it follows that p -gerbes, up to appropriate equivalence, are classified by $H^{p+1}(M, \mathcal{U}(1))$. We denote by $H^k(M; \mathbb{Z}_2, \mathcal{U}^\epsilon)$ the Grothendieck equivariant sheaf cohomology [11] of the equivariant sheaf \mathcal{U}^ϵ .

An *invariant p -gerbe* is a p -gerbe on an orientifold that is Real if p is even and \mathbb{Z}_2 -equivariant if p is odd. They are classified up to appropriate equivalence by $H^{p+1}(M; \mathbb{Z}_2, \mathcal{U}^{[p+2]})$, where $[k]$ denotes the reduction modulo 2 of $k \in \mathbb{Z}$. Thus the first three types of invariant p -gerbes are:

- $p = -1$: Real functions from M to $U(1)$ classified by $H^0(M; \mathbb{Z}_2, \mathcal{U}^1)$.
- $p = 0$: equivariant line bundles on M classified by $H^1(M; \mathbb{Z}_2, \mathcal{U}^0)$.

- and by the long exact sequences (2.3) it follows that these are all isomorphisms, and in fact equalities. Since ι_m is a Real map, the induced maps between the long exact sequences (2.4) and (2.5) commute. But $\sigma_{p+1}(\xi)(m) = \iota_m^*(\xi)$, so it follows that the connecting homomorphisms commute with the sign choice maps and we have

Proposition 2.6. $\sigma_{p+2} \circ \partial_{p+2} = \sigma_{p+1}$ for $p = -1, 0, 1, 2, \dots$

In particular, this begins with the commutative diagram from Section 1:

$$\begin{array}{ccc}
 H^0(M; \mathbb{Z}_2, \mathcal{U}^1) & & \\
 \downarrow \partial_1 & \searrow \sigma_0 & \\
 H^1(M; \mathbb{Z}_2, \mathcal{U}^0) & \xrightarrow{\sigma_1} & H^0(M^\tau, \mathbb{Z}_2) \\
 \downarrow \partial_2 & \nearrow \sigma_2 & \\
 H^2(M; \mathbb{Z}_2, \mathcal{U}^1) & & \\
 \downarrow \partial_3 & & \\
 \vdots & &
 \end{array} \tag{2.7}$$

Let M be a d -dimensional manifold. Then $H^d(M, \mathcal{U}(1)) = H^{d+1}(M, \mathbb{Z}) = 0$, so the long exact sequences (2.3) degenerate, and it follows that ∂_d is surjective and ∂_k is an isomorphism for $k > d$. Let $\Sigma_p(M) = \text{im}(\sigma_p)$ denote the image of the sign choice homomorphism and call its elements *geometric sign choices*. We conclude that

$$\Sigma_0(M) \subseteq \Sigma_1(M) \subseteq \dots \subseteq \Sigma_{d-1}(M) = \Sigma_d(M) = \Sigma_{d+1}(M) = \dots \subseteq H^0(M^\tau, \mathbb{Z}_2) .$$

In the sequel we consider geometric constructions of sign choices in the particular cases of interest, and the general question of what sign choices on an orientifold can arise as geometric sign choices.

3. REAL BUNDLE GERBES AND THEIR CONNECTIONS

We are particularly interested in the higher structures provided by invariant p -gerbes with $p = 1$, as these are the geometric data encoding the string orientifold backgrounds with NS-NS H -flux. In this section we consider this case in more detail and describe how to properly incorporate the B -field into the picture.

3.1. Real bundle gerbes. We begin by recalling some basic facts about Real bundle gerbes, referring the reader to [13] for more details.

Let M be a manifold and $Y \xrightarrow{\pi} M$ a surjective submersion. We denote by $Y^{[p]}$ the p -fold fibre product of Y with itself, that is, $Y^{[p]} = Y \times_M Y \times_M \dots \times_M Y$. This is a simplicial space $Y^{[\bullet]}$ whose face maps are given by the projections $\pi_i: Y^{[p]} \rightarrow Y^{[p-1]}$ which omit the i -th factor. Let $\Omega^q(Y^{[p]})$ denote the space of differential q -forms on $Y^{[p]}$ and define

$$\delta: \Omega^q(Y^{[p-1]}) \longrightarrow \Omega^q(Y^{[p]}) \quad \text{by} \quad \delta = \sum_{i=1}^p (-1)^{i-1} \pi_i^* .$$

These maps form an exact complex for all $q \geq 0$ called the *fundamental complex* [17]

$$0 \longrightarrow \Omega^q(M) \xrightarrow{\pi^*} \Omega^q(Y) \xrightarrow{\delta} \Omega^q(Y^{[2]}) \xrightarrow{\delta} \Omega^q(Y^{[3]}) \xrightarrow{\delta} \dots .$$

For any function $g: Y^{[p-1]} \rightarrow U(1)$, we define $\delta(g): Y^{[p]} \rightarrow U(1)$ by $\delta(g) = \sum_{i=1}^p (-1)^{i-1} g \circ \pi_i$. Similarly, to every $U(1)$ -bundle $P \rightarrow Y^{[p-1]}$ we associate a $U(1)$ -bundle $\delta(P) \rightarrow Y^{[p]}$ by

$$\delta(P) = \pi_1^{-1}(P) \otimes (\pi_2^{-1}(P))^* \otimes \pi_3^{-1}(P) \otimes \dots .$$

One easily checks that $\delta(\delta(g)) = 1$ and that $\delta(\delta(P))$ is canonically trivial.

Definition 3.1. A *bundle gerbe* (P, Y) over M is a principal $U(1)$ -bundle $P \rightarrow Y^{[2]}$ together with a bundle gerbe multiplication defined by a bundle isomorphism over $Y^{[3]}$,

$$\pi_3^{-1}(P) \otimes \pi_1^{-1}(P) \longrightarrow \pi_2^{-1}(P) ,$$

which is associative over $Y^{[4]}$. If $P_{(y_1, y_2)}$ denotes the fibre of P over $(y_1, y_2) \in Y^{[2]}$, then associativity means that the diagram

$$\begin{array}{ccc} P_{(y_1, y_2)} \otimes P_{(y_2, y_3)} \otimes P_{(y_3, y_4)} & \longrightarrow & P_{(y_1, y_3)} \otimes P_{(y_3, y_4)} \\ \downarrow & & \downarrow \\ P_{(y_1, y_2)} \otimes P_{(y_2, y_4)} & \longrightarrow & P_{(y_1, y_4)} \end{array}$$

commutes for all $(y_1, y_2, y_3, y_4) \in Y^{[4]}$.

Bundle gerbes are classified up to stable isomorphism [18] by their complex Dixmier–Douady class in the degree two sheaf cohomology

$$\mathrm{DD}(P) \in H^2(M, \mathcal{U}(1)) .$$

Following [16], if M is endowed with an involution $\tau: M \rightarrow M$ we define a *Real structure* on a bundle gerbe (P, Y) to be a pair of maps (τ_P, τ_Y) , where $\tau_Y: Y \rightarrow Y$ is an involution covering $\tau: M \rightarrow M$, and $\tau_P: P \rightarrow P$ is a conjugate involution covering $\tau_Y^{[2]}: Y^{[2]} \rightarrow Y^{[2]}$ and commuting with the bundle gerbe multiplication. Here conjugate involution means that $\tau_P(pz) = \tau_P(p)\bar{z}$, for $p \in P$ and $z \in U(1)$, and $\tau_P^2 = \mathrm{id}_P$. In the sequel we will suppress the subscripts on τ_P and τ_Y .

Definition 3.2. A *Real bundle gerbe* over M is a bundle gerbe (P, Y) over M with a Real structure.

In [13] we proved that Real bundle gerbes are classified up to Real stable isomorphism by their *Real Dixmier–Douady class* in the degree two equivariant sheaf cohomology group:

$$\mathrm{DD}_R(P) \in H^2(M; \mathbb{Z}_2, \mathcal{U}^1) .$$

3.2. Connective structures on Real bundle gerbes. Recall from [17] that if (P, Y) is a bundle gerbe, a connection A on the $U(1)$ -bundle $P \rightarrow Y^{[2]}$ is called a *bundle gerbe connection* if it respects the bundle gerbe multiplication. If A is a bundle gerbe connection, then the curvature $F_A \in \Omega^2(Y^{[2]})$ satisfies $\delta(F_A) = 0$. Then the exactness of the fundamental complex implies that there exists $B \in \Omega^2(Y)$ such that $F_A = \delta(B)$. This two-form is called a *curving* and the pair (A, B) is a *connective structure* on (P, Y) ; in string theory parlance, a connective structure is a B -field. As δ commutes with the exterior derivative d , we have $\delta(dB) = d\delta(B) = dF_A = 0$. Hence $dB = \pi^*(H)$ for some $H \in \Omega^3(M)$; in string theory the three-form H is called an H -flux. Moreover, $\pi^*(dH) = d\pi^*(H) = d^2B = 0$, so the 3-curvature H is closed and defines a de Rham representative for the complex Dixmier–Douady class of the bundle gerbe:

$$\mathrm{DD}(P) = \left[\frac{1}{2\pi i} H \right] \in H^3(M, \mathbb{Z}) .$$

Next let us consider the *dual bundle gerbe* (P^*, Y) . This is defined by the same manifold P but with the conjugate $U(1)$ -action, $(p, z) \mapsto p\bar{z}$. If A is a bundle gerbe connection on P , then $-A$ yields a bundle gerbe connection on P^* . Indeed if $\xi \in i\mathbb{R}$, let $\iota_p(\xi) \in T_p P$ be the vertical vector generated by the action of $U(1)$, and similarly let $\iota_p^*(\xi) \in T_p P^*$. Since $\iota_p^*(\xi) = -\iota_p(\xi)$ and $A(\iota_p^*(\xi)) = -\xi$, we conclude that $-A$ is a connection on P^* with curvature $-F_A$. For any choice

of curving $B \in \Omega^2(Y)$ for A , it follows that $-B$ is a curving for $-A$ and the 3-curvature of the connective structure $(-A, -B)$ for (P^*, Y) is given by $-H$.

Let (M, τ) be an orientifold and (P, Y) a Real bundle gerbe over M . Then $\tau: P^* \rightarrow P$ is a bundle gerbe isomorphism.

Definition 3.3. A *Real connective structure* for (P, Y) or *orientifold B -field* is a B -field (A, B) satisfying $\tau^*(A, B) = (-A, -B)$.

Proposition 3.4. *Real connective structures for (P, Y) exist.*

Proof. The existence of bundle gerbe connections is shown in [17]. Real bundle gerbe connections exist because if a is any bundle gerbe connection, then

$$A = \frac{1}{2}a + \frac{1}{2}(-\tau^*(a))$$

is a Real bundle gerbe connection. Any convex combination of connection one-forms is also a connection one-form. It follows that the curvature $F_A \in \Omega^2(Y^{[2]})$ also satisfies $\tau^*(F_A) = -F_A$. Consider a choice of curving $B \in \Omega^2(Y)$ satisfying $\delta(B) = F_A$. Hence

$$\delta(\tau^*(B)) = \tau^*(F_A) = -F_A = \delta(-B) .$$

It follows that $\delta(B + \tau^*(B)) = 0$ and thus there exists $\psi \in \Omega^1(M)$ such that $B + \tau^*(B) = \pi^*(\psi)$. Applying the involution we obtain $\tau^*(B) + B = \tau^*\pi^*(\psi)$, and since pullback of forms along π is injective, we find $\tau^*(\psi) = \psi$. Hence

$$B - \frac{1}{2}\pi^*(\psi) = -\tau^*(B) + \frac{1}{2}\pi^*(\tau^*(\psi)) = -\tau^*(B - \frac{1}{2}\pi^*(\psi)) ,$$

and moreover

$$F_A = \delta(B - \frac{1}{2}\pi^*(\psi))$$

because $\delta \circ \pi^* = 0$. We conclude that a curving B for a Real bundle gerbe connection can always be chosen such that $\tau^*(B) = -B$. \square

If (A, B) is an orientifold B -field, it follows that the H -flux satisfies $\tau^*(H) = -H$, so it gives a de Rham representative for the Real Dixmier–Douady class in the Borel equivariant cohomology with local coefficients $H^3(E\mathbb{Z}_2 \times_{\mathbb{Z}_2} M, \mathbb{Z}(1))$ [13]:

$$\text{DD}_R(P) = \left[\frac{1}{2\pi i} H \right] \in H_{\mathbb{Z}_2}^3(M, \mathbb{Z}(1)) .$$

Example 3.5 (Real bundle gerbes on $S^1 \times S^2$). If G is a compact simple Lie group with maximal torus T then there is a map, called the Weyl map, $T \times G/T \rightarrow G$ defined by $(t, gT) \mapsto g t g^{-1}$ for $t \in T$ and $g \in G$. This is a finite covering with fibre the Weyl group over the open dense subset of G of regular elements. In particular, if $G = SU(2)$ we get a map $S^1 \times S^2 \rightarrow S^3$ which is a double covering over S^3 minus the north and south poles which are the points I and $-I$ in $SU(2)$, the only elements of $SU(2)$ whose eigenvalues are not distinct. Under this map the basic bundle gerbe on S^3 , whose complex Dixmier–Douady class generates $H^3(S^3, \mathbb{Z}) = \mathbb{Z}$, pulls back to a bundle gerbe on $S^1 \times S^2$ whose complex Dixmier–Douady class is twice a generator in $H^3(S^1 \times S^2, \mathbb{Z}) = \mathbb{Z}$. This is because its integral over S^3 is equal to 1 but the Weyl map is two-to-one on a dense open subset, so the integral of the pulled back form is equal to 2.

The generator of $H^3(S^1 \times S^2, \mathbb{Z}) = \mathbb{Z}$ is given by a bundle gerbe which is a cup product of a function and a line bundle; we call it the cup-product bundle gerbe on $S^1 \times S^2$. Let $L \rightarrow S^2$ be the Hopf bundle; the pullback of its Chern class generates $H^2(S^1 \times S^2, \mathbb{Z}) = \mathbb{Z}$. Let $Y = \mathbb{R} \times S^2 \rightarrow S^1 \times S^2$ be the covering space where $S^1 = \mathbb{R}/\mathbb{Z}$. Then $Y^{[2]}$ can be identified with

the set of points $(x, y, u) \in \mathbb{R} \times \mathbb{R} \times U(1)$ where $x - y \in \mathbb{Z}$. Define $P \rightarrow Y^{[2]}$ by $P_{(x,y,u)} = L_u^{x-y}$. The bundle gerbe multiplication is immediate from

$$P_{(x,y,u)} \otimes P_{(y,z,u)} = L_u^{x-y} \otimes L_u^{y-z} = L_u^{x-z} = P_{(x,z,u)} .$$

It is straightforward to construct a connection and curving whose 3-curvature is $H = d(2\pi i \theta) \wedge \text{vol}_{S^2}$, where $\theta \in [0, 1)$ is a local coordinate on S^1 and vol_{S^2} is the unit area form on S^2 .

The Weyl map in this case also has a simple description as

$$S^1 \times S^2 \longrightarrow S^3 , \quad ((a, b), (x, y, z)) \longmapsto (bx, by, bz, a) . \quad (3.6)$$

There is no line bundle on S^3 which pulls back to the Hopf bundle on $S^1 \times S^2$ because all line bundles on S^3 are trivial. We give the three-sphere S^3 the Real structure $(x, y, z, w) \mapsto (-x, -y, -z, w)$ making it $S^{3,1}$. It is then immediate from (3.6) that there are two lifts of this Real structure to $S^1 \times S^2$: either to $((a, b), (x, y, z)) \mapsto ((a, -b), (x, y, z))$ making it $S^{1,1} \times S^{0,3}$, or to $((a, b), (x, y, z)) \mapsto ((a, b), (-x, -y, -z))$ making it $S^{0,2} \times S^{3,0}$. Here $S^{1,1}$ is the circle S^1 with the conjugation action on complex numbers of modulus one and $S^{3,0}$ is the two-sphere S^2 with the antipodal map. These are quite different Real structures; on $S^{0,2} \times S^{3,0}$ the involution is free whereas the involution on $S^{1,1} \times S^{0,3}$ gives two orientifold planes $\{\pm 1\} \times S^2$. By [13] the only line bundles on S^2 which admit a Real structure are those with even Chern class. Then using either the long exact sequence (2.3) or the spectral sequence for the Grothendieck equivariant cohomology one can characterise the Real bundle gerbes in these two cases. The calculations are straightforward and so we only summarise the results:

- The Real Brauer group of $S^{0,2} \times S^{3,0}$ equals \mathbb{Z} and the Real Dixmier–Douady class is given by $n \mapsto 2n$. So the pullback of the basic bundle gerbe on S^3 whose complex Dixmier–Douady class is twice the generator of $H^3(S^1 \times S^2, \mathbb{Z})$ admits a Real structure, but the cup-product bundle gerbe whose complex Dixmier–Douady class is the generator and whose H -flux is $H = d(2\pi i \theta) \wedge \text{vol}_{S^2}$, satisfying the necessary condition $\tau^*(H) = -H$, does not.
- The Real Brauer group of $S^{1,1} \times S^{0,3}$ equals $\mathbb{Z}_2 \oplus \mathbb{Z}_2 \oplus \mathbb{Z}_2 \oplus \mathbb{Z}$ and the Real Dixmier–Douady class is given by $(n, m, k, l) \mapsto l$. So the cup-product bundle gerbe whose complex Dixmier–Douady class is the generator and whose H -flux is $H = d(2\pi i \theta) \wedge \text{vol}_{S^2}$, satisfying $\tau^*(H) = -H$, admits a Real structure.

3.3. Holonomy of orientifold B -fields. Let (P, Y) be a bundle gerbe over a closed oriented surface Σ and choose a connective structure (A, B) . As $H^3(\Sigma, \mathbb{Z}) = 0$, the bundle gerbe is trivial. Fix a trivialisation $R \rightarrow Y$ with isomorphism $\delta(R) \simeq P$ and let a be a connection on R . Then A and $\delta(a)$ are both bundle gerbe connections for (P, Y) . Their difference is a one-form $A - \delta(a) \in \Omega^1(Y^{[2]})$ satisfying $\delta(A - \delta(a)) = 0$. Hence there is a one-form α on Y satisfying $A = \delta(a) + \delta(\alpha) = \delta(a + \alpha)$. In other words, we may choose a connection a on R such that $\delta(a) = A$. It follows that $\delta(F_a) = F_A = \delta(B)$ and so $B - F_a = \pi^*(\mu_a)$ for some $\mu_a \in \Omega^2(\Sigma)$. If we change the connection a to $b = a + \pi^*(\delta(\rho))$ for some $\rho \in \Omega^1(\Sigma)$, then $B - F_b = B - F_a - d\pi^*(\rho)$ so that

$$\mu_a = \mu_b + d\rho .$$

The *holonomy* of (A, B) over Σ is defined by

$$\text{hol}(\Sigma; A, B) = \exp \left(\int_{\Sigma} \mu_a \right) \in U(1) .$$

We note that the integral depends on the orientation of Σ , but is independent of the choice of a . Any other trivialisation of the bundle gerbe is of the form $R \otimes \pi^*(T)$ for some line bundle

$T \rightarrow \Sigma$. The connection a tensored by the pullback of a connection ∇ on T changes μ_a to $\mu_a - F_\nabla$, so the holonomy remains unchanged,

$$\exp \left(\int_{\Sigma} \mu_a \right) \exp \left(- \int_{\Sigma} F_\nabla \right) = \exp \left(\int_{\Sigma} \mu_a \right),$$

since the curvature F_∇ is an integral two-form. Hence the holonomy depends only on the triple $(\Sigma; A, B)$.

Now let (A, B) be a B -field on M . For any smooth map $\phi: \Sigma \rightarrow M$, we define the *holonomy of (A, B) along ϕ* by

$$\text{hol}(\phi; A, B) = \text{hol}(\Sigma; \phi^*(A), \phi^*(B)).$$

It is straightforward to check that the holonomies of connective structures for a bundle gerbe and its dual are related by

$$\text{hol}(\phi; A, B) = \text{hol}(\phi; -A, -B)^{-1}.$$

Hence if (A, B) is a Real connective structure for a Real bundle gerbe (P, Y) , we conclude that

$$\text{hol}(\phi; A, B) = \text{hol}(\tau \circ \phi; -A, -B)^{-1}.$$

In applications to the orientifold constructions of type II string theory, the \mathbb{Z}_2 -action on the spacetime M generated by the involution τ is combined with orientation-reversal of the string worldsheet. In this case the surface Σ is not oriented and need not even be orientable, and the string fields ϕ are smooth maps from Σ to the quotient space $X = M/\tau$ which represents the “physical points” of the orientifold spacetime. This is achieved in the orientifold sigma-model by regarding ϕ as maps to the total space of the fibration $M \rightarrow X$, and gauging the symmetry of the string theory. For this, we introduce the orientation double cover $\widehat{\Sigma} \rightarrow \Sigma$ corresponding to the first Stiefel–Whitney class $w_1(\Sigma) \in H^1(\Sigma, \mathbb{Z}_2)$, which is canonically oriented with a canonical orientation-reversing involution $\Omega: \widehat{\Sigma} \rightarrow \widehat{\Sigma}$ permuting the sheets and preserving the fibres, called worldsheet parity. The string fields are then smooth maps $\hat{\phi}: \widehat{\Sigma} \rightarrow M$ which are equivariant,

$$\hat{\phi} \circ \Omega = \tau \circ \hat{\phi}.$$

In [20, 10] they show that in this situation it is possible to define the *unoriented surface holonomy* of a Jandl gerbe which is a square root of $\text{hol}(\widehat{\Sigma}; \hat{\phi}^*(A), \hat{\phi}^*(B))$. We may define the *orientifold B -field holonomy* $\text{hol}(\phi; A, B)$ of a Real bundle gerbe with Real connection by noting that every Real bundle gerbe is naturally a Jandl gerbe and applying the construction in [20, 10]. To see why this is true, note that in our language a Jandl structure on a gerbe (P, Y) is a \mathbb{Z}_2 -equivariant trivialisation of the natural \mathbb{Z}_2 -action on $P \otimes \tau^{-1}(P)$. For a Real bundle gerbe (P, Y) this arises because $P \otimes \tau^{-1}(P) \simeq \delta L$ for $L_{(y,x)} = P_{(y,\tau(x))}^*$. The required switch isomorphism of [20] $\hat{s}: P_{(x,\tau(y))}^* \rightarrow P_{(y,\tau(x))}^*$ is induced by $\tau: P_{(x,y)} \rightarrow P_{(\tau(x),\tau(y))}$. It is a straightforward exercise to check that this satisfies the required conditions to be a Jandl structure, so we can apply the results of [20, 10] to define $\text{hol}(\phi; A, B)$. This defines precisely the B -field amplitude which was only schematically discussed in [13]. By [20, 10] it is invariant under the combined actions of the involutions Ω and τ which define the string orientifold construction and satisfies

$$\text{hol}(\phi; A, B) = \sqrt{\text{hol}(\widehat{\Sigma}; \hat{\phi}^*(A), \hat{\phi}^*(B))}.$$

As a further application, we can turn the construction of Real holonomy around and use it to provide sufficient conditions for the existence of Real structures on bundle gerbes over (M, τ) .

Proposition 3.7. *Let (M, τ) be an orientifold, where M is 2-connected, $M^\tau \neq \emptyset$ and H is a three-form on M with integral periods satisfying $\tau^*(H) = -H$. Then there is a tautological Real bundle gerbe over M with complex Dixmier–Douady class $[\frac{1}{2\pi i} H] \in H^3(M, \mathbb{Z})$.*

Proof. First we recall the construction of a bundle gerbe (P, Y) from an integral three-form H on a 2-connected manifold M . Fix a basepoint $m \in M$ and let $Y = \mathcal{P}M$ be the space of paths based at m with endpoint evaluation as projection to M . If $p_1, p_2 \in Y$ have the same endpoint, choose an oriented surface Σ in M spanning them, that is, the boundary of Σ is p_1 concatenated with the oppositely oriented path p_2 . Then the fibre of $P \rightarrow Y^{[2]}$ consists of all triples $((p_1, p_2), \Sigma, z)$, with $z \in U(1)$, modulo the equivalence relation $((p_1, p_2), \Sigma, z) \equiv ((p_1, p_2), \Sigma', z')$ if $\text{hol}(\Sigma \cup \Sigma'; H) z = z'$. Here $\Sigma \cup \Sigma'$ is the closed surface obtained by gluing Σ to the oppositely oriented Σ' along their common boundary, while $\text{hol}(S; H)$ for any closed oriented surface $S \subset M$ is the holonomy expressed as the standard Wess–Zumino–Witten term

$$\text{hol}(S; H) = \exp \left(\int_{B_S} H \right),$$

where B_S is a three-manifold whose boundary is S . This is well-defined since H is an integral form.

Next we consider a Real version of this construction. There is a natural way to pair elements of $P_{(p_1, p_2)}$ with elements of $P_{(p_2, p_1)}$. Given $((p_1, p_2), \Sigma, z)$, it can be paired with $((p_2, p_1), \Sigma, w)$ by keeping the same surface Σ but changing its orientation. Then we declare these two to pair to give $zw \in U(1)$. More generally, we can pair $((p_1, p_2), \Sigma, z)$ and $((p_2, p_1), \Sigma', w)$ by first using the equivalence relation to replace Σ' with Σ .

Now assume that the basepoint m of M is a fixed point of τ , so that τ acts on everything. Assume further that $\tau^*(H) = -H$. We define a Real structure τ by showing that

$$((p_1, p_2), \Sigma, z) \mapsto ((\tau(p_1), \tau(p_2)), \tau(\Sigma), \bar{z})$$

descends through the equivalence relation to give a conjugate bundle gerbe isomorphism $P \rightarrow \tau^{-1}(P)$.

The map τ satisfies $\tau^2 = 1$. If $((p_1, p_2), \Sigma, z) \equiv ((p_1, p_2), \Sigma', z')$, then $\text{hol}(\Sigma \cup \Sigma'; H) z = z'$. Now consider $((\tau(p_1), \tau(p_2)), \tau(\Sigma), \bar{z})$ and $((\tau(p_1), \tau(p_2)), \tau(\Sigma'), \bar{z}')$. Then we find

$$\begin{aligned} \text{hol}(\tau(\Sigma) \cup \tau(\Sigma'); H) &= \text{hol}(\tau(\Sigma) \cup \tau(\Sigma'); -\tau^*(H)) \\ &= \text{hol}(\Sigma \cup \Sigma'; -H) \\ &= \overline{\text{hol}(\Sigma \cup \Sigma'; H)}, \end{aligned}$$

so that $\text{hol}(\tau(\Sigma) \cup \tau(\Sigma'); H) \bar{z} = \bar{z}'$ as required.

Finally, we check that τ commutes with the bundle gerbe multiplication. On the one hand, the multiplication is given by

$$((p_1, p_2), \Sigma, z) ((p_2, p_3), \Sigma', z') = ((p_1, p_3), \Sigma \cup \Sigma', z z').$$

On the other hand, under τ we obtain

$$((\tau(p_1), \tau(p_2)), \tau(\Sigma), \bar{z}) ((\tau(p_2), \tau(p_3)), \tau(\Sigma'), \bar{z}') = ((\tau(p_1), \tau(p_3)), \tau(\Sigma \cup \Sigma'), \overline{z z'})$$

as required. \square

Remark 3.8. In general, the class of the three-form H of Proposition 3.7 does not lift to the Real Dixmier–Douady class of the tautological Real bundle gerbe in Borel equivariant cohomology with local coefficients. The conditions under which it does, and hence represents the H -flux of an orientifold B -field, are discussed in [13].

4. GEOMETRIC CONSTRUCTIONS OF SIGN CHOICES

In this section we explain how the sign choice homomorphism (2.2) can be constructed geometrically from the invariant p -gerbes when $p = -1, 0, 1$, that is, for Real functions, \mathbb{Z}_2 -equivariant line bundles and Real bundle gerbes.

4.1. Sign choice for Real functions. Let g be a Real function on (M, τ) . If $m \in M^\tau$ then $g(m) = \overline{g(m)} = \pm 1$, so there is a natural sign choice homomorphism

$$\sigma_0: H^0(M; \mathbb{Z}_2, \mathcal{U}^1) \longrightarrow H^0(M^\tau, \mathbb{Z}_2)$$

defined by evaluation $\sigma_0(g)(m) = g(m)$. This map is multiplicative in the sense that $\sigma_0(gh) = \sigma_0(g)\sigma_0(h)$. If $g = f(\bar{f} \circ \tau)$ for an arbitrary function f on M and m is a fixed point of τ , then $g(m) = f(m)\overline{f(m)} = 1$ so that $\sigma_0(f(\bar{f} \circ \tau)) = 1$. In other words, the composition

$$H^0(M, \mathcal{U}(1)) \xrightarrow{f \mapsto f(\bar{f} \circ \tau)} H^0(M; \mathbb{Z}_2, \mathcal{U}^1) \xrightarrow{\sigma_0} H^0(M^\tau, \mathbb{Z}_2)$$

is equal to 1. In fact, there is a stronger result in this case given by

Proposition 4.1. *If M is 1-connected, then the sequence*

$$0 \longrightarrow H^0(M; \mathbb{Z}_2, \mathcal{U}^0) \longrightarrow H^0(M, \mathcal{U}(1)) \xrightarrow{f \mapsto f(\bar{f} \circ \tau)} H^0(M; \mathbb{Z}_2, \mathcal{U}^1) \xrightarrow{\epsilon} \mathbb{Z}_2 \longrightarrow 0$$

is exact.

Proof. If $g: M \rightarrow U(1)$ is a Real function, consider a lift $\hat{g} \in H^0(M, \mathcal{U}(1))$. Then $\hat{g} \circ \tau + \hat{g} = k$ for some $k \in \mathbb{Z}$, and the image $[k] \in \mathbb{Z}_2$ is well-defined independently of the lift \hat{g} of g . We call it $\epsilon(g)$. If $\epsilon(g) = 0$, then we can define $\hat{f} = -\frac{1}{2}\hat{g}$ with

$$\hat{f} \circ \tau - \hat{f} = \frac{1}{2}\hat{g} - \frac{1}{2}(\hat{g} \circ \tau) = \hat{g}$$

so that $(f \circ \tau)\bar{f} = g$. □

Corollary 4.2. *If M is 1-connected, then $\Sigma_0(M) \subseteq \mathbb{Z}_2$.*

Hence if M is 1-connected and M^τ contains more than one O-plane, then σ_0 is not surjective. We discuss further constraints on geometric sign choices in Section 5.4.

4.2. Sign choice for equivariant bundles. Let us first revisit a basic result from [13], namely the problem of classifying, up to isomorphism, the \mathbb{Z}_2 -structures on a $U(1)$ -bundle L which admits a \mathbb{Z}_2 -structure.

If τ is a \mathbb{Z}_2 -structure on L , then any other \mathbb{Z}_2 -structure $\hat{\tau}$ is given by $\hat{\tau} = g\tau$ for some function $g: M \rightarrow U(1)$. By $\hat{\tau}^2 = 1$ it follows that $g(g \circ \tau) = 1$ or $g = \bar{g} \circ \tau$. We say that $\hat{\tau}$ and τ are isomorphic if there is a smooth function $f: M \rightarrow U(1)$ such that $\hat{\tau}f = f\tau$; this is equivalent to $(f \circ \tau)g\tau = f\tau$ or $g = f(\bar{f} \circ \tau)$.

Proposition 4.3. *The space of non-isomorphic \mathbb{Z}_2 -structures on a bundle $L \rightarrow M$ admitting a \mathbb{Z}_2 -structure is the quotient of $H^0(M; \mathbb{Z}_2, \mathcal{U}^1)$ by the image of the homomorphism*

$$H^0(M; \mathcal{U}(1)) \xrightarrow{f \mapsto f(\bar{f} \circ \tau)} H^0(M; \mathbb{Z}_2, \mathcal{U}^1) .$$

From this description it follows that the space of \mathbb{Z}_2 -structures on L is independent of the choice of L as long as L admits at least one \mathbb{Z}_2 -structure.

If L is a \mathbb{Z}_2 -equivariant line bundle over (M, τ) , then its fibre over a fixed point $m \in M^\tau$ is simply a copy of \mathbb{C} with a \mathbb{Z}_2 -action by multiplication with ± 1 which we label $\text{sign}(L_m)$. This allows us to define a homomorphism

$$\sigma_1: H^1(M; \mathbb{Z}_2, \mathcal{U}^0) \longrightarrow H^0(M^\tau, \mathbb{Z}_2)$$

by $\sigma_1(L)(m) = \text{sign}(L_m)$. Again it is clear that this map is multiplicative in the sense that $\sigma_1(L \otimes K) = \sigma_1(L) \sigma_1(K)$. If L is an arbitrary line bundle on M , then $L \otimes \tau^{-1}(L)$ has the canonical \mathbb{Z}_2 -action induced by swapping factors of the tensor product

$$L_x \otimes L_{\tau(x)} \longrightarrow L_{\tau(x)} \otimes L_x .$$

At a fixed point this is the identity map, so it has sign choice equal to 1. It follows that the composition

$$H^1(M, \mathcal{U}(1)) \xrightarrow{L \mapsto L \otimes \tau^{-1}(L)} H^1(M; \mathbb{Z}_2, \mathcal{U}^0) \xrightarrow{\sigma_1} H^0(M^\tau, \mathbb{Z}_2)$$

is equal to 1.

4.3. Sign choice for Real bundle gerbes. If (P, Y) is a bundle gerbe and $y \in Y$, then the fibre $P_{(y,y)} = U(1)$. This follows from the fact that the multiplication map $P_{(y,y)} \otimes P_{(y,y)} \rightarrow P_{(y,y)}$ commutes with the $U(1)$ -action on $P_{(y,y)}$. So if $p \in P_{(y,y)}$ and $q \in P_{(y,y)}$, then $pq \in P_{(y,y)}$ and hence $pq = p \langle q \rangle$ for some $\langle q \rangle \in U(1)$. If we change p to pz for $z \in U(1)$, then

$$(pz)q = (pq)z = p \langle q \rangle z = (pz) \langle q \rangle$$

so the map $q \mapsto \langle q \rangle$ is independent of the choice of p . We need two results about this map. Firstly, if $r \in P_{(y',y)}$ then

$$(rpr^{-1})(rq r^{-1}) = r(pq)r^{-1} = rp \langle q \rangle r^{-1} = (rpr^{-1}) \langle q \rangle$$

so that $\langle rqr^{-1} \rangle = \langle q \rangle$. Secondly if τ is a Real structure, then $\tau(pq) = \tau(p)\tau(q)$ and $\tau(pq) = \tau(p \langle q \rangle) = \tau(p) \overline{\langle q \rangle}$, so that $\langle \tau(q) \rangle = \overline{\langle q \rangle}$.

Consider now a Real bundle gerbe $\mathcal{G} = (P, Y)$ over M , and let $m \in M^\tau$ be a fixed point of τ . Choose $y \in Y_m$, then $(y, \tau(y)) \in Y^{[2]}$. Let $f \in P_{(y, \tau(y))}$ so that $f\tau(f) \in P_{(y,y)}$, and define

$$\sigma_2(\mathcal{G})(m) = \sigma_2(P, Y)(m) = \langle f \tau(f) \rangle . \quad (4.4)$$

Proposition 4.5. (a) $\sigma_2(\mathcal{G})(m)$ is independent of the choice of f and y .

(b) $\sigma_2(\mathcal{G})(m) \in \mathbb{Z}_2$.

Proof. (a) If we change f to fz for $z \in U(1)$, then

$$\langle (fz) \tau(fz) \rangle = z \bar{z} \langle f \tau(f) \rangle = \langle f \tau(f) \rangle ,$$

so $\langle f \tau(f) \rangle \in U(1)$ is independent of the choice of $f \in P_{(y, \tau(y))}$. Changing y to $\tilde{y} \in Y_m$ and letting $g \in P_{(\tilde{y}, y)}$, then $gf\tau(g^{-1}) \in P_{(\tilde{y}, \tau(\tilde{y}))}$ and

$$\langle (gf\tau(g^{-1})) \tau(gf\tau(g^{-1})) \rangle = \langle gf\tau(g^{-1}) \tau(g) \tau(f) g^{-1} \rangle = \langle g(f\tau(f)) g^{-1} \rangle = \langle f \tau(f) \rangle .$$

(b) From (a) we know that $\langle \tau(f) f \rangle = \langle f \tau(f) \rangle$ so

$$\overline{\langle f \tau(f) \rangle} = \langle \tau(f \tau(f)) \rangle = \langle \tau(f) f \rangle = \langle f \tau(f) \rangle .$$

Hence $\sigma_2(\mathcal{G})(m) \in \mathbb{Z}_2$. □

This shows that letting m vary determines an element $\sigma_2(\mathcal{G}) \in H^0(M^\tau, \mathbb{Z}_2)$. The next result implies that the sign choice of a Real bundle gerbe depends only on its Real stable isomorphism class.

- Proposition 4.6.** (a) If \mathcal{G} is a Real bundle gerbe, then $\sigma_2(\mathcal{G}^*) = \sigma_2(\mathcal{G})$.
 (b) If \mathcal{G}_1 and \mathcal{G}_2 are Real bundle gerbes, then $\sigma_2(\mathcal{G}_1 \otimes \mathcal{G}_2) = \sigma_2(\mathcal{G}_1) \sigma_2(\mathcal{G}_2)$.
 (c) If \mathcal{G} is a trivial Real bundle gerbe, then $\sigma_2(\mathcal{G}) = 1$.
 (d) If \mathcal{G}_1 and \mathcal{G}_2 are Real stably isomorphic Real bundle gerbes, then $\sigma_2(\mathcal{G}_1) = \sigma_2(\mathcal{G}_2)$.

Proof. It is straightforward from the definition (4.4) to prove (a) and (b). Part (d) follows from (c) because \mathcal{G}_1 and \mathcal{G}_2 are Real stably isomorphic if and only if $\mathcal{G}_1 \otimes \mathcal{G}_2^*$ is Real trivial. For (c), take a Real bundle $R \rightarrow Y$ and for $(y_1, y_2) \in Y^{[2]}$ define $P_{(y_1, y_2)} = R_{y_2} \otimes R_{y_1}^*$ with the Real structure induced by that of R . Then the bundle gerbe multiplication

$$P_{(y_1, y_2)} \otimes P_{(y_2, y_3)} \longrightarrow P_{(y_1, y_3)}$$

comes from the obvious contractions of

$$R_{y_2} \otimes R_{y_1}^* \otimes R_{y_3} \otimes R_{y_2}^* \longrightarrow R_{y_3} \otimes R_{y_1}^* .$$

It is straightforward to show that $\langle r \otimes s^* \rangle = s^*(r)$ for $r \otimes s^* \in P_{(y, y)} = R_y \otimes R_y^*$.

Let $r \in R_y$ and define $f = \tau(r) \otimes r^* \in P_{(y, \tau(y))} = R_{\tau(y)} \otimes R_y^*$, where r^* is the element that satisfies $r^*(r) = 1$. Then

$$\langle f \tau(f) \rangle = \langle (\tau(r) \otimes r^*) (r \otimes \tau(r^*)) \rangle = \langle \tau(r) \otimes \tau(r)^* \rangle = 1$$

as required. \square

Thus we have a well-defined homomorphism

$$\sigma_2: H^2(M; \mathbb{Z}_2, \mathcal{U}^1) \longrightarrow H^0(M^\tau, \mathbb{Z}_2) .$$

Example 4.7 (Real bundle gerbes over a point). If $M = \text{pt}$, then there are two Real stable isomorphism classes of Real bundle gerbes (cf. Section 2). Their sign choices are +1 for the trivial one and -1 for the non-trivial one. By Proposition 4.6 (c) it is clear that the trivial bundle gerbe has sign choice +1. We construct the non-trivial sign choice as follows. Let $M = \{m\}$ and $Y = M \times \mathbb{Z}_2$ with the Real structure $\tau(m, 0) = (m, 1)$ and $\tau(m, 1) = (m, 0)$. Then $Y^{[2]} = M \times \mathbb{Z}_2 \times \mathbb{Z}_2$ and we take the trivial bundle $P \rightarrow Y^{[2]}$ equipped with the trivial multiplication, so that $p \mapsto \langle p \rangle$ is just the identity map. We define the Real structure to be conjugation from $P_{((m, 0), (m, 0))} \rightarrow P_{((m, 1), (m, 1))}$ and $P_{((m, 1), (m, 1))} \rightarrow P_{((m, 0), (m, 0))}$, and -1 times conjugation from $P_{((m, 0), (m, 1))} \rightarrow P_{((m, 1), (m, 0))}$ and $P_{((m, 1), (m, 0))} \rightarrow P_{((m, 0), (m, 1))}$.

A straightforward calculation shows that this commutes with the bundle gerbe multiplication and squares to the identity map. To calculate the sign, let $y = (m, 0)$ so that $\tau(y) = (m, 1)$ and let $f \in P_{((m, 0), (m, 1))}$ be 1. Then $\langle f \tau(f) \rangle = -1$ as required. This bundle gerbe cannot be a trivial Real bundle gerbe because it has a non-trivial sign. The fact that $H^2(\{m\}; \mathbb{Z}_2, \mathcal{U}^1) = \mathbb{Z}_2$ shows that these are the only two Real stable isomorphism classes of Real bundle gerbes over a point, and the calculation demonstrates that

$$\sigma_2: H^2(\{m\}; \mathbb{Z}_2, \mathcal{U}^1) \longrightarrow H^0(\{m\}^\tau, \mathbb{Z}_2)$$

is an isomorphism. In Section 7.1 we will generalise this construction and show that it agrees with the connecting homomorphism in (2.4).

Example 4.8 (Tautological Real bundle gerbes). Under the conditions of Proposition 3.7, we show that the sign choice of the tautological Real bundle gerbe (P, Y) is given by

$$\sigma_2(P) = \text{hol}(\Sigma \cup \tau(\Sigma); H) .$$

Let $m \in M^\tau$ be the basepoint in the proof of Proposition 3.7 and let $x \in M^\tau$ be another fixed point. Choose $p_1 = p$ to be a path from m to x and $p_2 = \tau(p)$ as per the way one

computes the sign choice, see Proposition 4.5. Picking a spanning surface Σ , we then have $[p, \tau(p), \Sigma, 1] \in P_{(p, \tau(p))}$ and

$$\tau([p, \tau(p), \Sigma, 1]) = [\tau(p), p, \tau(\Sigma), 1] = [\tau(p), p, \Sigma, \text{hol}(\Sigma \cup \tau(\Sigma); H)] .$$

These pair to give the sign choice $\text{hol}(\Sigma \cup \tau(\Sigma); H)$. This construction will be used later to exhibit a constraint imposed on the sign choices when M is 2-connected.

Let $\phi: N \rightarrow M$ be a Real map between orientifolds. Then there is an induced homomorphism $\phi^*: H^0(M^\tau, \mathbb{Z}_2) \rightarrow H^0(N^\tau, \mathbb{Z}_2)$. If (P, Y) is a Real bundle gerbe on M , then

$$\phi^*(\sigma_2(P)) = \sigma_2(\phi^{-1}(P)) .$$

In particular, if $m \in M^\tau$ is a fixed point, we can take $N = \{m\}$ and ϕ the inclusion to find $\sigma_2(P)(m) = \phi^*(\sigma_2(P)) = \sigma_2(\phi^{-1}(P))$. In other words, we can define the sign choice map by restricting $\sigma_2(P)$ to any fixed point $m \in M^\tau$, and declaring $\sigma_2(P)(m)$ to be $+1$ if the restricted Real bundle gerbe is Real trivial and -1 otherwise. Thus we conclude

Proposition 4.9. *The sign choice map (4.4) for Real bundle gerbes \mathcal{G} coincides with the sign choice of its Real Dixmier–Douady class $\text{DD}_R(\mathcal{G})$ as defined in (2.2).*

Similarly to the cases of Real functions and equivariant line bundles, we have

Proposition 4.10. *The composition*

$$H^2(M, \mathcal{U}(1)) \xrightarrow{Q \mapsto Q \otimes \tau^{-1}(Q^*)} H^2(M; \mathbb{Z}_2, \mathcal{U}^1) \xrightarrow{\sigma_2} H^0(M^\tau, \mathbb{Z}_2)$$

is equal to 1.

Proof. If (Q, X) is a bundle gerbe on M , define a Real bundle gerbe (P, Y) by $Y = X \times_M \tau^{-1}(X)$ and $P = Q \otimes \tau^{-1}(Q^*)$. A fibre point $y = (x, x') \in Y_m = X_m \times X_{\tau(m)}$ satisfies $\tau(x, x') = (x', x)$. The fibre of $P = Q \otimes \tau^{-1}(Q^*)$ at $(y_1, y_2) = ((x_1, x'_1), (x_2, x'_2))$ is $Q_{(x_1, x_2)} \otimes Q_{(x'_1, x'_2)}^*$. If $\tau(m) = m$, we can pick $y = (x, x) \in Y_m$. Any $f = q \otimes q^* \in P_{(y, \tau(y))} = Q_{(x, x)} \otimes Q_{(x, x)}^*$ satisfies $\tau(f) = q^* \otimes q$ and $f \tau(f) = q(q^*) \otimes q(q^*) = 1 \in P_{(y, y)}$, which maps to $1 \in U(1)$ as required. \square

Let $X = M/\tau$ denote the quotient space as in the orientifold constructions of type II string theory. By applying the Grothendieck spectral sequence to the fibration $\pi: M \rightarrow X$, it is possible to give a general characterisation of the kernel and image of the sign choice map σ_2 in terms of the sheaf cohomology of the space of “physical points” X of the orientifold (M, τ) .

Theorem 4.11. *There exists a five-term exact sequence*

$$0 \longrightarrow H^2(X, \pi_*^\tau \mathcal{U}^1) \longrightarrow H^2(M; \mathbb{Z}_2, \mathcal{U}^1) \xrightarrow{\sigma_2} H^0(M^\tau, \mathbb{Z}_2) \xrightarrow{d_3} H^3(X, \pi_*^\tau \mathcal{U}^1)$$

where $\pi_*^\tau \mathcal{U}^1$ is the invariant direct image sheaf.

Proof. The equivariant sheaf cohomology groups are the derived functors of the right exact functor $H^0(M; \mathbb{Z}_2, \mathcal{U}^1) = \Gamma^\tau(\mathcal{U}^1)$ of global invariant sections of the \mathbb{Z}_2 -sheaf \mathcal{U}^1 :

$$H^q(M; \mathbb{Z}_2, \mathcal{U}^1) = (R^q \Gamma^\tau)(\mathcal{U}^1) .$$

Let Γ denote the global sections functor on the quotient space X and π_*^τ the invariant direct image functor. Then $\Gamma^\tau = \Gamma \circ \pi_*^\tau$ is the composition of two right exact functors, so the Grothendieck spectral sequence gives

$$E_2^{p, q} = (R^p \Gamma)(R^q \pi_*^\tau)(\mathcal{U}^1) = H^p(X, (R^q \pi_*^\tau)(\mathcal{U}^1))$$

converging to

$$(R^{p+q} \Gamma^\tau)(\mathcal{U}^1) = H^{p+q}(M; \mathbb{Z}_2, \mathcal{U}^1) .$$

The sheaf $(R^q \pi_*^\tau)(\mathcal{U}^1)$ is trivial for q odd and equals the sheaf of constant \mathbb{Z}_2 -valued functions supported on M^τ for q even. Namely if $m \in M$ is not a fixed point, then we can find a contractible open neighbourhood $m \in U$ such that $\tau(U) \cap U = \emptyset$, so $\pi(U)$ is an open neighbourhood of $x = \pi(m) \in X$. The sheaf $(R^q \pi_*^\tau)(\mathcal{U}^1)$ is associated to the pre-sheaf $(V \subset X) \mapsto H^q(\pi^{-1}(V); \mathbb{Z}_2, \mathcal{U}^1|_{\pi^{-1}(V)})$, and for $q > 0$ we get

$$H^q(\pi^{-1}(\pi(U)); \mathbb{Z}_2, \mathcal{U}^1) = H^q(U \cup \tau(U); \mathbb{Z}_2, \mathcal{U}^1) = 0$$

since U is contractible. It follows that for $q > 0$, the sheaf $(R^q \pi_*^\tau)(\mathcal{U}^1)$ is supported on the fixed point set M^τ . Moreover, by choosing an invariant open neighbourhood around each connected component of M^τ which equivariantly retracts onto that O-plane, we conclude that the stalk of $(R^q \pi_*^\tau)(\mathcal{U}^1)$ at any point $x \in M^\tau$ is simply $H^q(\text{pt}; \mathbb{Z}_2, \mathcal{U}^1)$, which equals 0 for q odd and \mathbb{Z}_2 for q even. Since $\text{Aut}(\mathbb{Z}_2) = 1$, there is also no monodromy.

Inserted into the Grothendieck spectral sequence, the result follows by the exact sequence of low degree terms since $E_2^{p,1} = 0$, $E_2^{2,0} = H^2(X, \pi_*^\tau \mathcal{U}^1)$, and the third differential d_3 maps $E_2^{0,2} = H^0(M^\tau, \mathbb{Z}_2)$ to $E_2^{3,0} = H^3(X, \pi_*^\tau \mathcal{U}^1)$. \square

Remark 4.12. Theorem 4.11 gives a genuine obstruction to possible sign choices: the obstruction to surjectivity of σ_2 is the third differential d_3 and the obstruction class is realised by an ‘orbifold 2-gerbe’ on X with band in the bundle of groups $\pi_*^\tau \mathcal{U}^1$, which lifts to a trivialisable Real bundle 2-gerbe on M that corresponds physically to the flux of a flat three-form Ramond–Ramond field; this is an obstruction only on orientifold spacetimes which have O-planes of dimension a multiple of 4 [3, 15]. If the obstruction map d_3 is trivial, then there is a short exact sequence

$$0 \longrightarrow H^2(X, \pi_*^\tau \mathcal{U}^1) \longrightarrow H^2(M; \mathbb{Z}_2, \mathcal{U}^1) \xrightarrow{\sigma_2} H^0(M^\tau, \mathbb{Z}_2) \longrightarrow 0 \quad (4.13)$$

but it is not necessarily split. We shall encounter some concrete examples of these facts in the following sections.

5. SIGN CHOICES FOR SPHERICAL ORIENTIFOLDS

Spheres provide important examples of orientifold compactifications of string theory with background fluxes. In this section we will consider the n -spheres $S^{m,1}$ for $n > 1$ equipped with an involution with fixed points, and apply the techniques from the previous sections to derive a general formula for the geometric sign choices for a broad class of orientifolds.

5.1. The three-sphere $S^{3,1}$. One of the most common examples of a string orientifold involves the three-sphere $S^{3,1} \subset \mathbb{R}^4$ equipped with the involution $(x, y, z, w) \mapsto (-x, -y, -z, w)$. This is equivalent to the Lie group $SU(2)$ equipped with the involution $\tau: g \mapsto g^{-1}$, so it has two fixed points $\pm I$ corresponding to the north and south poles of the three-sphere. The long exact sequence (2.3) yields

$$\begin{array}{ccccccc} 0 = H^1(S^{3,1}, \mathcal{U}(1)) & \longrightarrow & H^1(S^{3,1}; \mathbb{Z}_2, \mathcal{U}^0) & \longrightarrow & H^2(S^{3,1}; \mathbb{Z}_2, \mathcal{U}^1) & \longrightarrow & \\ \text{\scriptsize \downarrow} & & & & & & \\ & \longrightarrow & H^2(S^{3,1}, \mathcal{U}(1)) & \longrightarrow & H^2(S^{3,1}; \mathbb{Z}_2, \mathcal{U}(1)) = 0 & & \end{array}$$

where the vanishing of the last group follows from the spectral sequence for Grothendieck's equivariant sheaf cohomology. Since $H^2(S^{3,1}, \mathcal{U}(1)) = H^3(S^3, \mathbb{Z}) = \mathbb{Z}$, we obtain

$$0 \longrightarrow H^1(S^{3,1}; \mathbb{Z}_2, \mathcal{U}^0) \xrightarrow{\partial_2} H^2(S^{3,1}; \mathbb{Z}_2, \mathcal{U}^1) \longrightarrow \mathbb{Z} \longrightarrow 0.$$

The first group equals \mathbb{Z}_2 since it is the group of isomorphism classes of \mathbb{Z}_2 -equivariant line bundles on $S^{3,1}$. As any line bundle on S^3 is trivial, the only possible \mathbb{Z}_2 -equivariant extensions are given by multiplication with $\tau = +1$ or $\tau = -1$ on the fibre \mathbb{C} . This gives the Real Brauer group

$$H^2(S^{3,1}; \mathbb{Z}_2, \mathcal{U}^1) = \mathbb{Z}_2 \oplus \mathbb{Z},$$

where the complex Dixmier–Douady class of (a, b) is b . This can also be verified by using the Grothendieck spectral sequence.

The group of sign choices is $H^0((S^{3,1})^\tau, \mathbb{Z}_2) = \mathbb{Z}_2 \oplus \mathbb{Z}_2$. We proceed to show that the sign choice map

$$\sigma_2: \mathbb{Z}_2 \oplus \mathbb{Z} \longrightarrow \mathbb{Z}_2 \oplus \mathbb{Z}_2$$

sends \mathbb{Z}_2 to the diagonal, and that it maps the basic bundle gerbe generating \mathbb{Z} to $(-1, 1)$.

For this, we will apply the construction of the sign choice map from Example 4.8, with H given by $2\pi i$ times the normalized volume form on S^3 . Consider the group $SU(2)$ and take the basepoint I . We regard S^3 as \mathbb{R}^3 with the identity I as $(0, 0, 0)$ and $\tau(x, y, z) = (-x, -y, -z)$. Then the fixed point $-I$ is at infinity. Let p be the path from the origin along the positive z -axis to infinity. Applying τ then gives the path from the origin along the negative z -axis to infinity. The surface $\Sigma \subset S^3$ is taken to be the set of all $(x, 0, z)$ with $x \geq 0$, and then $\tau(\Sigma)$ is the set of all $(x, 0, z)$ with $x \leq 0$. The union $S = \Sigma \cup \tau(\Sigma)$ is the (x, z) -plane, which we take to bound the ball B_S given by the half-space (x, y, z) with $y \geq 0$. The holonomy $\text{hol}(\Sigma \cup \tau(\Sigma); H)$ is then computed by integrating H over half of \mathbb{R}^3 , or in other words over half of S^3 . The integral of H over all of S^3 equals $2\pi i$, and hence the sought integral is πi and the holonomy is -1 . Thus the sign choice of the basic bundle gerbe at the point $-I \in SU(2)$ is -1 . At the other pole of the sphere we can take all paths and Σ constant, giving the sign choice $+1$.

We conclude that in the group $\mathbb{Z}_2 \oplus \mathbb{Z}$ of Real bundle gerbes on $SU(2)$, the basic bundle gerbe is $(0, 1)$, while the Real bundle gerbe coming from the connecting homomorphism, with sign choices -1 at both fixed points, is $(-1, 0)$. The geometric sign choices are $\Sigma_2(S^{3,1}) = \mathbb{Z}_2 \oplus \mathbb{Z}_2$, with the basic bundle gerbe mapping to $(-1, 1)$ and the coboundary bundle gerbe mapping to $(-1, -1)$. Hence the map from $\mathbb{Z}_2 \oplus \mathbb{Z}$ to $\mathbb{Z}_2 \oplus \mathbb{Z}_2$ is $(a, b) \mapsto (a + [b], a - [b])$, where $b \mapsto [b]$ is the reduction map $\mathbb{Z} \rightarrow \mathbb{Z}_2$. Thus the sign choice map is surjective, but $\mathbb{Z}_2 \oplus \mathbb{Z}_2$ is not a direct summand in the Real Brauer group. In particular, for all sign choices to arise one needs to consider Real bundle gerbes with non-vanishing complex Dixmier–Douady class.

Another way of seeing this is to use the Mayer–Vietoris sequence. Write $S^{3,1} = U_+ \cup U_-$ as the union of the upper and lower hemispheres of S^3 . These are equivariantly contractible and each contribute $H^2(\text{pt}; \mathbb{Z}_2, \mathcal{U}^1) = \mathbb{Z}_2$. The involution on the equator $U_+ \cap U_-$ acts freely and the quotient is the real projective plane $\mathbb{R}P^2$. Then

$$H^1(\mathbb{R}P^2; \mathbb{Z}_2, \mathcal{U}^1) \simeq H^2(\mathbb{R}P^2, \mathbb{Z}_{-1}) \simeq H^0(\mathbb{R}P^2, \mathbb{Z}) = \mathbb{Z}$$

where \mathbb{Z}_{-1} is the orientation bundle on $\mathbb{R}P^2$ and the second isomorphism is Poincaré duality. Since $H^2(\mathbb{R}P^2; \mathbb{Z}_2, \mathcal{U}^1) = H^3(\mathbb{R}P^2, \mathbb{Z}_{-1}) = 0$ for dimensional reasons, the Mayer–Vietoris sequence gives

$$0 \longrightarrow \mathbb{Z} \longrightarrow \mathbb{Z}_2 \oplus \mathbb{Z} \xrightarrow{\sigma_2} \mathbb{Z}_2 \oplus \mathbb{Z}_2 \longrightarrow 0$$

where the restriction map σ_2 sends $(0, 1)$ to $(-1, 1)$. This is an instance of the exact sequence (4.13) where σ_2 is a non-split surjection.

5.2. The n -sphere $S^{n,1}$. Let $S^{n,1} \subset \mathbb{R}^{n+1}$ denote the n -sphere with $n \geq 4$, equipped with the Real structure induced by the involution $\tau: (x_1, x_2, \dots, x_n, x_{n+1}) \mapsto (-x_1, -x_2, \dots, -x_n, x_{n+1})$. The long exact sequence (2.3) yields

$$0 = H^1(S^{n,1}, \mathcal{U}(1)) \longrightarrow H^1(S^{n,1}; \mathbb{Z}_2, \mathcal{U}^0) \longrightarrow H^2(S^{n,1}; \mathbb{Z}_2, \mathcal{U}^1) \longrightarrow H^2(S^{n,1}, \mathcal{U}(1)) = 0.$$

We have $H^1(S^{n,1}; \mathbb{Z}_2, \mathcal{U}^0) = \mathbb{Z}_2$ as the trivial line bundle on $S^{n,1}$ admits two inequivalent Real structures. It follows that the Real Brauer group of $S^{n,1}$ is $H^2(S^{n,1}; \mathbb{Z}_2, \mathcal{U}^1) = \mathbb{Z}_2$. On the other hand, the involution on $S^{n,1}$ has two fixed points, one at each pole of the sphere, so the group of sign choices is $H^0((S^{n,1})^\tau, \mathbb{Z}_2) = \mathbb{Z}_2^2$. We conclude that σ_2 is not surjective in this case.

More generally, consider a product of k such Real spheres $M = S^{n_1,1} \times \dots \times S^{n_k,1}$ with $n_i \geq 4$. Then $H^2(M; \mathbb{Z}_2, \mathcal{U}^1) = \mathbb{Z}_2$ while $H^0(M^\tau, \mathbb{Z}_2) = \mathbb{Z}_2^{2^k}$, so the sign choice map σ_2 is far from being onto. However, in these cases it is more natural to ask whether the sign choice map σ_{p+1} associated to p -gerbes with $p = n_1 + \dots + n_k - 2$ is surjective. An explicit check is hampered by our inability to geometrically define p -gerbes in general. We consider below some explicit cases for $p = 2$ where this can be addressed. The discussion would extend to a suitable geometric realisation of p -gerbes for all $p > 2$, wherein the approach sketched below would work by inductively defining σ_{p+1} in terms of σ_p .

5.3. Sign choice for equivariant bundle 2-gerbes. We briefly discuss the behaviour of the sign choice map σ_3 . Write $\mathcal{G} \Rightarrow M$ for a bundle gerbe over a manifold M . A *bundle 2-gerbe* \mathcal{G} over M consists of a surjective submersion $Y \rightarrow M$ together with a bundle gerbe $\mathcal{G} \Rightarrow Y^{[2]}$ subject to some additional conditions detailed in [21]. If (M, τ) is an orientifold, then we assume that \mathcal{G} is equivariant in the strong sense that τ lifts to act on all the spaces by involutions. The equivariant sheaf cohomology group $H^3(M; \mathbb{Z}_2, \mathcal{U}^0)$ classifies \mathbb{Z}_2 -equivariant bundle 2-gerbes, up to appropriate equivalence.

The connecting homomorphism ∂_3 sends Real bundle gerbes to \mathbb{Z}_2 -equivariant bundle 2-gerbes, and $\sigma_3 \circ \partial_3 = \sigma_2$ by Proposition 2.6. Let $m \in M^\tau$, and let $y \in Y_m$ be the corresponding fibre of Y over M , then $\tau: \mathcal{G}_{(y, \tau(y))} \rightarrow \mathcal{G}_{(\tau(y), y)}$. From the structure of the bundle 2-gerbe there is also an inversion map $\mathcal{G}_{(\tau(y), y)} \rightarrow \mathcal{G}_{(y, \tau(y))}$ which acts by conjugation. Thus composing the equivariant structure and the inverse gives a Real structure $\mathcal{G}_{(y, \tau(y))} \rightarrow \mathcal{G}_{(y, \tau(y))}$, which has a sign choice. We declare this to be the sign choice $\sigma_3(\mathcal{G})(m)$ of the equivariant bundle 2-gerbe \mathcal{G} over M at the fixed point m . To justify this definition, note that in the case of bundle gerbes the same construction would define an equivariant action $P_{(y, \tau(y))} \rightarrow P_{(y, \tau(y))}$ which in the notation of Section 4.3 would be $f \mapsto \sigma(\mathcal{G})(m) \tau(f)^{-1}$, and we would take $\sigma(\mathcal{G})(m)$ as our sign choice. But then $\sigma(\mathcal{G})(m) = \langle f \tau(f) \rangle = \sigma_2(\mathcal{G})(m)$.

Example 5.1 (σ_3 is onto). We show that for $S^{4,1}$ the sign choice homomorphism

$$\sigma_3: H^3(S^{4,1}; \mathbb{Z}_2, \mathcal{U}^0) \longrightarrow \mathbb{Z}_2 \oplus \mathbb{Z}_2$$

is surjective. Since $S^{4,1}$ is 2-connected and the complex Dixmier–Douady class of any Real bundle gerbe on $S^{4,1}$ vanishes, it follows from Proposition 5.4 below that σ_2 on such Real bundle gerbes is either $(1, 1)$ or $(-1, -1)$. Hence σ_2 on the corresponding equivariant bundle 2-gerbes is the same. To obtain other sign choices it suffices to find an equivariant bundle 2-gerbe whose sign choice is $(1, -1)$ or $(-1, 1)$, as we can get the other sign choice by tensoring with the $(-1, -1)$ bundle 2-gerbe.

We show that the bundle 2-gerbe is the Chern–Simons bundle 2-gerbe [4] for the standard $SU(2)$ -bundle $P \rightarrow S^4$. The total space of this bundle is given by

$$P = \{(w^0, w^1) \in \mathbb{H}^2 \mid w^0 \bar{w}^0 + w^1 \bar{w}^1 = 1\} = S^7 \subset \mathbb{H}^2 = \mathbb{R}^8,$$

and the cosets of the right action by the unit quaternions $Sp(1) = SU(2)$ is $P_1\mathbb{H} \simeq S^4$. Define the orientifold action on P by $\tau(w^0, w^1) = (-w^0, w^1)$, which descends to the action on $S^{4,1}$. Over $[1, 0]$ the involution τ acts freely and over $[0, 1]$ it acts trivially. The Chern–Simons bundle 2-gerbe is defined as follows. There is a map $\rho: P^{[2]} \rightarrow SU(2)$ defined by $p_1 = p_2 \rho(p_1, p_2)$. Then the bundle gerbe over $P^{[2]}$ is the pullback of the basic bundle gerbe $\mathcal{G} \Rightarrow SU(2)$, which we denote by \mathcal{H} . The action of τ on P satisfies $\rho \circ \tau = \rho$, so the equivariant structure can be taken to be the identity

$$\mathcal{H}_{(y_1, y_2)} = \mathcal{G}_{\rho(y_1, y_2)} = \mathcal{G}_{\rho(\tau(y_1), \tau(y_2))} = \mathcal{H}_{(\tau(y_1), \tau(y_2))}.$$

There are two cases:

- Let $y = (0, 1)$, then $\tau(y) = (0, 1)$ so that $\rho(y, \tau(y)) = I \in SU(2)$. Thus $\mathcal{H}_{(y, \tau(y))} = \mathcal{G}_I$. The sign choice we want is that of this Real structure on \mathcal{G}_I , which we know from Section 5.1 equals +1.
- Let $y = (1, 0)$, then $\tau(y) = (-1, 0)$ so that $\rho(y, \tau(y)) = -I \in SU(2)$. The inversion map on the bundle gerbe is the usual Real structure when identifying $SU(2) = S^{3,1}$, so the sign choice we want is that of this Real structure on \mathcal{G}_{-I} , which we know from Section 5.1 equals -1.

Example 5.2 (σ_3 is not onto). Consider the quadric $M = S^{2,1} \times S^{2,1}$. There are four fixed points in this case, so the group of sign choices is $H^0(M^\tau, \mathbb{Z}_2) = \mathbb{Z}_2^4$. An explicit calculation shows that all the geometric sign choices, including equivariant bundle 2-gerbes, are contained in the subgroup of sign choices which have an even number of -1 signs. Hence $\Sigma_3(M) \neq H^0(M^\tau, \mathbb{Z}_2)$.

5.4. Constraints on geometric sign choices. We shall now derive a criterion that can often be used to constrain the possible geometric sign choices on an orientifold (M, τ) .

Definition 5.3. Let $m \neq m'$ be fixed points in M^τ . A *Real d -span* of m and m' is a Real map $\mu_{m, m'}: S^{d,1} \rightarrow M$ with $\mu_{m, m'}(1, 0, \dots, 0) = m$ and $\mu_{m, m'}(-1, 0, \dots, 0) = m'$.

The problem of finding a Real d -span may be formulated in terms of certain constraints on (M, τ) as follows. Suppose that M is $d-1$ -connected with m and m' isolated fixed points of τ . Assume that it is possible to choose a path γ from m to m' which avoids the fixed points. Then $\tau(\gamma)$ is another path from m to m' that does not intersect γ except at m and m' , and $\gamma \cup \tau(\gamma)$ defines a map of $S^{1,1}$ into M , which is a Real 1-span since it commutes with the involutions on M and on S^1 . Next if $d > 1$, then M is 1-connected and suppose that a spanning surface Σ for γ and $\tau(\gamma)$ can be chosen such that it avoids the fixed point set. Then $\tau(\Sigma)$ also has boundary $\gamma \cup \tau(\gamma)$ and the union $\Sigma \cup \tau(\Sigma)$ determines a Real 2-span. Continuing in this way, and as long as the fixed point set can be avoided, we obtain a Real d -span.

Proposition 5.4. Let ξ be an invariant p -gerbe over an orientifold (M, τ) with complex characteristic class $C_{p+2}(\xi) \in H^{p+2}(M, \mathbb{Z})$ and sign choice $\sigma_{p+1}(\xi) \in H^0(M^\tau, \mathbb{Z}_2)$. If $\mu_{m, m'}$ is a Real $(p+2)$ -span of $m, m' \in M^\tau$, then

$$\sigma_{p+1}(\xi)(m) = (-1)^{\langle \mu_{m, m'}^*(C_{p+2}(\xi)), S^{p+2} \rangle} \sigma_{p+1}(\xi)(m'). \quad (5.5)$$

Proof. Since the span is a Real map, it pulls back the invariant p -gerbe on M to the sphere $S^{p+2,1}$. The sign of the pulled back p -gerbe at $(1, 0, \dots, 0)$ is $\sigma_{p+1}(\xi)(m)$ and at $(-1, 0, \dots, 0)$ it equals

$\sigma_{p+1}(\xi)(m')$. The complex characteristic class of the pulled back p -gerbe is $\mu_{m,m'}^*(C_{p+2}(\xi))$. The result follows by the calculation in Sections 5.1 and 5.2, which shows that an invariant p -gerbe on $S^{p+2,1}$ has signs that differ precisely by the periods of its complex characteristic class. \square

The formula (5.5) constrains the possible sign choices because once a sign choice is made at one fixed point, all the other sign choices are determined by the periods of the class of the p -gerbe. This is independent of the Real or equivariant structure on the p -gerbe and depends solely on the periods of $C_{p+2}(\xi)$. In particular, if the p -gerbe ξ has vanishing complex characteristic class $C_{p+2}(\xi) = 0$, then the sign choice map $\sigma_{p+1}(\xi)$ is constant and all sign choices are the same.

6. SIGN CHOICES FOR TOROIDAL ORIENTIFOLDS

Tori constitute important examples of flat orientifold compactifications of string theory. In this section we consider examples of multiply connected orientifolds involving tori. The allowed sign choices in these cases have been studied from different perspectives in [5, 9, 7, 8].

6.1. The circle $S^{1,1}$. We denote by $S^{1,1}$ the circle $S^1 = U(1)$ with the Real structure $\tau(u) = \bar{u}$ and fixed points ± 1 . We can identify $H^0((S^{1,1})^\tau, \mathbb{Z}_2)$ with $\mathbb{Z}_2 \oplus \mathbb{Z}_2$ by the map $f \mapsto (f(1), f(-1))$. There are two types of string theories to which this pertains to: the orientifold type IIA string theory compactified on $S^{1,1}$ is the type IA theory which under T-duality maps to type I string theory on a circle, or equivalently the orientifold type IIB theory on $S^{0,2}$, whereas the type $\widehat{\text{IA}}$ theory is the T-dual of the type IIB orientifold on $S^{2,0}$ [14, 19]. The two theories are distinguished by the fact that in the former case the two O-planes have the same sign while in the latter case they have opposite signs. Both theories arise within our present formalism if all sign choices are possible, which is guaranteed by

Proposition 6.1. *The sign choice map*

$$\sigma_2: H^2(S^{1,1}; \mathbb{Z}_2, \mathcal{U}^1) \longrightarrow H^0((S^{1,1})^\tau, \mathbb{Z}_2) = \mathbb{Z}_2^2 \quad (6.2)$$

is an isomorphism.

Proof. The long exact sequence (2.3) in this case degenerates to

$$0 \longrightarrow H^1(S^{1,1}; \mathbb{Z}_2, \mathcal{U}^0) \longrightarrow H^2(S^{1,1}; \mathbb{Z}_2, \mathcal{U}^1) \longrightarrow 0, \quad (6.3)$$

so we need to understand the space of equivariant line bundles over S^1 . Every line bundle over S^1 is trivial and the generic \mathbb{Z}_2 -action on the trivial bundle is $g\tau$, where $g: S^1 \rightarrow U(1)$ and τ is the trivial action on the trivial bundle. To compute the number of inequivalent \mathbb{Z}_2 -structures on the trivial bundle, we need to calculate the quotient in Proposition 4.3. We proceed by showing that the sequence

$$H^0(S^1, \mathcal{U}(1)) \xrightarrow{f \mapsto f(\bar{f} \circ \tau)} H^0(S^{1,1}; \mathbb{Z}_2, \mathcal{U}^1) \xrightarrow{\sigma_0} H^0((S^{1,1})^\tau, \mathbb{Z}_2) \longrightarrow 0 \quad (6.4)$$

is exact, from which it follows that

$$\sigma_1: H^1(S^{1,1}; \mathbb{Z}_2, \mathcal{U}^0) \longrightarrow H^0((S^{1,1})^\tau, \mathbb{Z}_2)$$

is an isomorphism. The result then follows by the connecting isomorphism (6.3).

To establish the exactness of (6.4), we show that for any $g \in H^0(S^{1,1}; \mathbb{Z}_2, \mathcal{U}^1)$ with $g(1) = 1$ and $g(-1) = 1$, there exists a function $f: S^1 \rightarrow U(1)$ with $g(u) = f(u)\overline{f(\bar{u})}$. Identify $S^1 = \mathbb{R}/\mathbb{Z}$ and lift g to $\hat{g}: \mathbb{R} \rightarrow \mathbb{R}$. Then $\hat{g}(t+1) = \hat{g}(t) + n$ for some $n \in \mathbb{Z}$ which is the winding number

of g . Because $g(1) = 1$ we can assume that $\hat{g}(0) = 0$, and then by $g(-1) = 1$ it follows that $\hat{g}(\frac{1}{2}) = k$ for some $k \in \mathbb{Z}$. Also because $\overline{g(u)} = g(u)$ for all $u \in S^1$, we have

$$-\hat{g}(1-t) = \hat{g}(t) + m$$

for some $m \in \mathbb{Z}$. Taking $t = \frac{1}{2}$ and $t = 0$ then shows that $m = -2k$ and $n = 2k$. Define $\hat{f}(t) = \frac{1}{2}\hat{g}(t)$, then

$$\hat{f}(t+1) = \hat{f}(t) + k$$

and

$$\hat{f}(t) - \hat{f}(1-t) = \frac{1}{2}\hat{g}(t) - \frac{1}{2}\hat{g}(1-t) = \hat{g}(t) - k.$$

We conclude that \hat{f} descends to a well-defined function $f: S^1 \rightarrow U(1)$ satisfying $f(u)\overline{f(\bar{u})} = g(u)$ as required. \square

6.2. The two-torus $S^{1,1} \times S^{1,1}$. The case of a two-torus is analogous but more complicated since now there are non-trivial line bundles on $T^2 = S^{1,1} \times S^{1,1}$. These are classified by $H^2(T^2, \mathbb{Z}) = \mathbb{Z}$. A line bundle whose Chern class is the generator can be described as follows [9]. If $\mathbf{x}, \mathbf{y} \in \mathbb{R}^2$ let

$$c(\mathbf{x}, \mathbf{y}) = \exp(2\pi i x^1(x^2 - y^2)).$$

Define an equivalence relation on $\mathbb{R}^2 \times U(1)$ by $(\mathbf{x}, u) \sim (\mathbf{y}, v)$ if $\mathbf{x} - \mathbf{y} \in \mathbb{Z}^2$ and $v = c(\mathbf{x}, \mathbf{y})u$. The sought $U(1)$ -bundle P is defined as the quotient by this equivalence relation, with the projection $\pi([\mathbf{x}, u]) = [\mathbf{x}]$ where $[\mathbf{x}]$ denotes the orbit in $S^1 \times S^1 = \mathbb{R}^2/\mathbb{Z}^2$. We lift the Real structure on $S^{1,1} \times S^{1,1}$ to P by defining $\tau([\mathbf{x}, u]) = [-\mathbf{x}, u]$. Checking that this is well-defined reduces to observing that $c(-\mathbf{x}, -\mathbf{y}) = c(\mathbf{x}, \mathbf{y})$. The fixed points of τ on T^2 are $(0, 0)$, $(\frac{1}{2}, 0)$, $(0, \frac{1}{2})$ and $(\frac{1}{2}, \frac{1}{2})$. On the corresponding fibres the Real structure acts by

$$\tau([\mathbf{x}, u]) = [-\mathbf{x}, u] = [\mathbf{x}, \exp(4\pi i x^1 x^2) u],$$

so τ acts by the identity on the four fixed points. Hence $\sigma_1(P) = (1, 1, 1, -1)$ where we denote a sign choice by its values on the four fixed points as listed above. In other words, every odd power of P admits a \mathbb{Z}_2 -structure with this same sign choice and even powers have trivial sign choice.

Next we need to study the quotient in Proposition 4.3 and consider the sequence

$$H^0(T^2, \mathcal{U}(1)) \xrightarrow{f \mapsto f(\bar{f} \circ \tau)} H^0(T^2; \mathbb{Z}_2, \mathcal{U}^1) \xrightarrow{\sigma_0} H^0((T^2)^\tau, \mathbb{Z}_2) = \mathbb{Z}_2^4.$$

Let $E \subset H^0((T^2)^\tau, \mathbb{Z}_2)$ denote the subgroup of sign choices whose product is $+1$, that is, there are an even number of -1 's. Then $E \simeq \mathbb{Z}_2^3$ as any element is determined by its value on just three of the four fixed points. We have

Proposition 6.5. *The image of σ_0 is E .*

Proof. Consider the functions $g: T^2 \rightarrow U(1)$, $(u, v) \mapsto g(u, v)$ defined by the values $\pm 1, \pm u, \pm v$ and $\pm uv$. These give all the sign choices with even numbers of negative signs. Assume now that there is a function g with $g(1, 1) = 1$ and $g(1, -1) = g(-1, 1) = -1$. We show that it must have $g(-1, -1) = 1$. We may lift g to $\hat{g}: \mathbb{R}^2 \rightarrow \mathbb{R}$ in such a way that $\hat{g}(0, 0) = 0$. Moreover we must have $\hat{g}(x+1, y) = \hat{g}(x, y) + m$, $\hat{g}(x, y+1) = \hat{g}(x, y) + n$ and $-\hat{g}(-x, -y) = \hat{g}(x, y) + k$ for some integers m, n and k . Substituting $\hat{g}(0, 0) = 0$ gives $k = 0$. In addition we have $\hat{g}(\frac{1}{2}, 0) = p + \frac{1}{2}$ and $\hat{g}(0, \frac{1}{2}) = q + \frac{1}{2}$ for some $p, q \in \mathbb{Z}$, so $\hat{g}(\frac{1}{2}, \frac{1}{2}) = -\hat{g}(-\frac{1}{2}, -\frac{1}{2})$ and $\hat{g}(\frac{1}{2}, \frac{1}{2}) = \hat{g}(-\frac{1}{2}, -\frac{1}{2}) + m + n$. Thus $\hat{g}(\frac{1}{2}, \frac{1}{2}) = \frac{1}{2}(m + n)$. Applying a similar argument gives $\hat{g}(\frac{1}{2}, 0) = \frac{m}{2} = p + \frac{1}{2}$ and $\hat{g}(0, \frac{1}{2}) = \frac{n}{2} = q + \frac{1}{2}$. Hence $\hat{g}(\frac{1}{2}, \frac{1}{2}) = p + q + 1 \in \mathbb{Z}$ giving the required result. Now any sign

choice with an odd number of negative signs is a sign choice with an even number of negative signs multiplied by $(1, -1, -1, 1)$, so none of them can occur in the image of σ_0 . \square

Proposition 6.6. *If $\sigma_0(g) = (1, 1, 1, 1)$, then there exists a smooth function $f: T^2 \rightarrow U(1)$ such that $g(u, v) = f(u, v) \overline{f(\bar{u}, \bar{v})}$.*

Proof. As in the proof of Proposition 6.5 we lift g to $\hat{g}: \mathbb{R}^2 \rightarrow \mathbb{R}$ such that $\hat{g}(x+1, y) = \hat{g}(x, y) + m$ and $\hat{g}(x, y+1) = \hat{g}(x, y) + n$. We may assume that $\hat{g}(0, 0) = 0$ and thus $-\hat{g}(-x, -y) = \hat{g}(x, y)$. Moreover $\hat{g}(\frac{1}{2}, 0) = p$, $\hat{g}(0, \frac{1}{2}) = q$ and $\hat{g}(\frac{1}{2}, \frac{1}{2}) = r$ where $p, q, r \in \mathbb{Z}$. Hence $\hat{g}(\frac{1}{2}, 0) = \hat{g}(-\frac{1}{2}, 0) + m$ and $p = \hat{g}(\frac{1}{2}, 0) = -\hat{g}(-\frac{1}{2}, 0)$ so that $m = 2p$. Similarly $n = 2q$. Define

$$\hat{f}(x, y) = \frac{1}{2} \hat{g}(x, y)$$

and notice that $\hat{f}(x+1, y) = \hat{f}(x, y) + p$ and $\hat{f}(x, y+1) = \hat{f}(x, y) + q$, so that \hat{f} descends to a well-defined function $f: T^2 \rightarrow U(1)$. Moreover

$$\hat{f}(x, y) - \hat{f}(-x, -y) = \frac{1}{2} \hat{g}(x, y) - \frac{1}{2} \hat{g}(-x, -y) = \hat{g}(x, y),$$

so that $g(u, v) = f(u, v) \overline{f(\bar{u}, \bar{v})}$ as required. \square

It follows that the sequence

$$H^1(T^2, \mathcal{U}(1)) \xrightarrow{P \mapsto P \otimes \tau^{-1}(P)} H^1(T^2; \mathbb{Z}_2, \mathcal{U}^0) \xrightarrow{\partial_2} H^2(T^2; \mathbb{Z}_2, \mathcal{U}^1) \longrightarrow 0 \quad (6.7)$$

is exact since T^2 is two-dimensional. Moreover we have

$$H^1(T^2; \mathbb{Z}_2, \mathcal{U}^0) = \mathbb{Z} \oplus \mathbb{Z}_2^3$$

as line bundles on T^2 with arbitrary Chern class admit any choice of \mathbb{Z}_2 -action. From the calculation above of the sign choice for the generator in $H^2(T^2, \mathbb{Z})$, we conclude that σ_1 maps $\mathbb{Z} \oplus \mathbb{Z}_2^3$ to $\mathbb{Z}_2 \oplus \mathbb{Z}_2^3 = \mathbb{Z}_2^4$ surjectively. This agrees with the results of [9], which we will explain in more generality in Section 7.

The image of the map $P \mapsto P \otimes \tau^{-1}(P)$ consists of the line bundles with even Chern class, so the Real Brauer group is

$$H^2(T^2; \mathbb{Z}_2, \mathcal{U}^1) = \mathbb{Z} \oplus \mathbb{Z}_2^3$$

and the action of σ_2 is again projection onto \mathbb{Z}_2^4 . In summary, we have

Proposition 6.8. *The Real Brauer group of the two-torus $T^2 = S^{1,1} \times S^{1,1}$ is*

$$H^2(T^2; \mathbb{Z}_2, \mathcal{U}^1) = \mathbb{Z} \oplus \mathbb{Z}_2^3$$

and the sign choice map σ_2 is the projection onto $H^0((T^2)^\tau, \mathbb{Z}_2) = \mathbb{Z}_2^4$ induced by reduction modulo 2 on the first factor.

6.3. The three-torus $S^{1,1} \times S^{1,1} \times S^{1,1}$. With the conjugation action on each circle factor, the involution on the three-dimensional torus T^3 has eight fixed points and hence there are 2^8 possible sign choices. Since $T^3 = \mathbb{R}^3/\mathbb{Z}^3$, the fixed points are the image of the eight points (ρ_1, ρ_2, ρ_3) where $\rho_i \in \{0, \frac{1}{2}\}$. We can regard the fixed points on the three-torus as forming a cube. Let $E \subset H^0((T^3)^\tau, \mathbb{Z}_2)$ be the subgroup of all sign choices with the property that their product around each square of the cube equals 1. Then it is straightforward to see that E contains 16 elements. We now show that the sequence

$$H^0(T^3, \mathcal{U}(1)) \xrightarrow{f \mapsto f(\bar{f} \circ \tau)} H^0(T^3; \mathbb{Z}_2, \mathcal{U}^1) \xrightarrow{\sigma_0} E \longrightarrow 0$$

is well-defined, that is, σ_0 takes values in E , and that it is exact. Consider $f: T^3 \rightarrow U(1)$ satisfying $f(z) \overline{f(\bar{z})} = 1$ for all $z \in T^3$, and lift f to the universal cover $\hat{f}: \mathbb{R}^3 \rightarrow \mathbb{R}$. If e_1, e_2, e_3

is the standard basis of \mathbb{R}^3 , then $\hat{f}(\mathbf{x} + \mathbf{e}_i) = \hat{f}(\mathbf{x}) + n_i$ for $n_i \in \mathbb{Z}$ and $\hat{f}(\mathbf{x}) + \hat{f}(-\mathbf{x}) = k \in \mathbb{Z}$, for all $\mathbf{x} \in \mathbb{R}^3$. If $\epsilon_i \in \{0, 1\}$ for $i = 1, 2, 3$, then it follows that

$$\hat{f}\left(\frac{1}{2} \sum_{i=1}^3 \epsilon_i \mathbf{e}_i\right) = \frac{1}{2} \left(k + \sum_{i=1}^3 n_i \epsilon_i\right). \quad (6.9)$$

Let i, j, k be distinct and $\epsilon \in \{0, 1\}$, then adding up around a square gives

$$\hat{f}\left(\frac{1}{2} \epsilon \mathbf{e}_k\right) + \hat{f}\left(\frac{1}{2} \epsilon \mathbf{e}_k + \frac{1}{2} \mathbf{e}_i\right) + \hat{f}\left(\frac{1}{2} \epsilon \mathbf{e}_k + \frac{1}{2} \mathbf{e}_j\right) + \hat{f}\left(\frac{1}{2} \epsilon \mathbf{e}_k + \frac{1}{2} \mathbf{e}_i + \frac{1}{2} \mathbf{e}_j\right) = 2k + 2\epsilon n_k + n_i + n_j \in \mathbb{Z}$$

and so multiplying the values of f around the corresponding square of fixed points gives 1. Hence the image of σ_0 is contained in E .

Next we show that σ_0 is surjective. For each S^1 factor there are two Real functions 1 and z , combining these makes eight Real functions and applying ± 1 makes 16. They are all distinct so σ_0 is surjective.

Finally we prove exactness in the centre. That is, if $\sigma_0(f) = 1$ we show that there is a function $g : T^3 \rightarrow U(1)$ such that $f(z) = g(z) \overline{g(\bar{z})}$. Lift f to $\hat{f} : \mathbb{R}^3 \rightarrow \mathbb{R}$ as above. Then from (6.9) we deduce that

$$\frac{1}{2} \left(k + \sum_{i=1}^3 n_i \epsilon_i\right)$$

is always an integer and hence that $k, n_1, n_2, n_3 \in 2\mathbb{Z}$. The function

$$\hat{g}(\mathbf{x}) = \frac{1}{2} \hat{f}(\mathbf{x})$$

satisfies $\hat{g}(\mathbf{x} + \mathbf{e}_i) - \hat{g}(\mathbf{x}) \in \mathbb{Z}$ and so descends to a well-defined function $g : T^3 \rightarrow U(1)$. Moreover we obtain

$$\hat{g}(\mathbf{x}) - \hat{g}(-\mathbf{x}) = \frac{1}{2} \hat{f}(\mathbf{x}) - \frac{1}{2} \hat{f}(-\mathbf{x}) = \hat{f}(\mathbf{x}) - \frac{1}{2} k$$

as required. This gives the claimed exactness. This also shows that $E \simeq \mathbb{Z}_2^4$.

Consider now the exact sequence

$$H^1(T^3, \mathcal{U}(1)) \xrightarrow{P \mapsto P \otimes \tau^{-1}(P)} H^1(T^3; \mathbb{Z}_2, \mathcal{U}^0) \xrightarrow{\partial_2} H^2(T^3; \mathbb{Z}_2, \mathcal{U}^1) \longrightarrow H^2(T^3, \mathcal{U}(1)).$$

It follows from the Künneth theorem that $H^1(T^3, \mathcal{U}(1)) = \mathbb{Z}^3$, and each of the generating line bundles is a pullback via a projection $T^3 \rightarrow T^2$ of the standard bundle constructed in Section 6.2, so each admits a \mathbb{Z}_2 -action. Hence

$$H^1(T^3; \mathbb{Z}_2, \mathcal{U}^0) \simeq \mathbb{Z}^3 \oplus E \simeq \mathbb{Z}^3 \oplus \mathbb{Z}_2^4,$$

with the sign choice map σ_1 being the projection of $\mathbb{Z}^3 \oplus \mathbb{Z}_2^4$ onto \mathbb{Z}_2^7 . The image consists of all sign choices whose product over the eight fixed points is equal to 1. The map $P \mapsto P \otimes \tau^{-1}(P)$ is multiplication by 2 on the \mathbb{Z}^3 factor.

Finally we consider Real bundle gerbes on T^3 . Let $Y = \mathbb{R}^3 \rightarrow T^3$ be the universal cover and define $g : Y^{[3]} \rightarrow U(1)$ by

$$g(\mathbf{x}, \mathbf{y}, \mathbf{z}) = \exp(2\pi i x^1 (x^2 - y^2)(y^3 - z^3)).$$

It is straightforward to show that $\delta(g) : Y^{[4]} \rightarrow U(1)$ equals 1. This gives a bundle gerbe $P = Y^{[2]} \times U(1)$ with multiplication

$$((\mathbf{x}, \mathbf{y}), u) ((\mathbf{y}, \mathbf{z}), v) = ((\mathbf{x}, \mathbf{z}), g(\mathbf{x}, \mathbf{y}, \mathbf{z}) u v).$$

There is a lift of the Real structure on T^3 to Y by $\tau(\mathbf{x}) = -\mathbf{x}$. Since $g(-\mathbf{x}, -\mathbf{y}, -\mathbf{z}) = \overline{g(\mathbf{x}, \mathbf{y}, \mathbf{z})}$, we obtain a Real structure on P given by $\tau((\mathbf{x}, \mathbf{y}), u) = ((-\mathbf{x}, -\mathbf{y}), \bar{u})$. One easily verifies that this bundle gerbe has complex Dixmier–Douady class generating $H^2(T^3, \mathcal{U}(1)) = \mathbb{Z}$. Hence

$$H^2(T^3; \mathbb{Z}_2, \mathcal{U}^1) = \mathbb{Z}_2^3 \oplus E \oplus \mathbb{Z}.$$

We calculate the sign of the Real bundle gerbe following the procedure of Section 6.2. We take $\mathbf{x} \in Y$ so that $(\mathbf{x}, -\mathbf{x}) \in Y^{[2]}$ and pick $p = (\mathbf{x}, -\mathbf{x}, 1) \in P_{(\mathbf{x}, -\mathbf{x})}$. Then $\tau(p) = (-\mathbf{x}, \mathbf{x}, 1)$ and

$$p\tau(p) = ((\mathbf{x}, \mathbf{x}), g(\mathbf{x}, -\mathbf{x}, \mathbf{x})) = ((\mathbf{x}, \mathbf{x}), \exp(-8\pi i x^1 x^2 x^3)).$$

Thus the sign choice is 1 except at $(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$ where it equals -1 .

Proposition 6.10. *The Real Brauer group of the three-torus $T^3 = S^{1,1} \times S^{1,1} \times S^{1,1}$ is*

$$H^2(T^3; \mathbb{Z}_2, \mathcal{U}^1) = \mathbb{Z}_2^7 \oplus \mathbb{Z},$$

and the sign choice map σ_2 is the projection onto $H^0((T^3)^\tau, \mathbb{Z}_2) = \mathbb{Z}_2^8$ induced by reduction modulo 2 on the last factor.

7. KR-THEORY WITH SIGN CHOICES

In this final section we make contact with the prescription of [9] for computing the sign choices in orientifold backgrounds of type II string theory whose complex Dixmier–Douady class is trivial, that is, when $H = dB$ globally. This enables a general KR -theory classification of D-brane charges in the cases when the sign choice map is not constant over the orientifold planes.

7.1. The Gao–Hori construction. From explicit worldsheet considerations, Gao and Hori in [9] gave the relevant geometric ingredients required of any type II string orientifold construction with topologically trivial H -flux, which we briefly review and then translate into the language of the present paper. Let (M, τ) be an orientifold. Then the data of [9] consists of a quadruple (B, L, α, c) where:

- $B \in \Omega^2(M)$ is a globally defined B -field on M .
- L is a line bundle over M called the twist bundle.
- α is a connection on L called the twist connection.
- c is a global section of $\tau^{-1}(L) \otimes L^*$ called the crosscap section.

The line bundle with connection (L, α) originates from requiring invariance of the B -field amplitude (see Section 3.3) under the parity transform $\hat{\phi} \mapsto \tau \circ \hat{\phi} \circ \Omega$ for all string worldsheets $\hat{\phi}: \widehat{\Sigma} \rightarrow M$, which for a globally defined B -field implies that the two-form $B + \tau^*(B)$ has integral periods, and so represents a class in $H^2(M, \mathbb{Z})$ or equivalently the curvature of a connection on a line bundle over M . Invariance of D-brane Chan–Paton factors under the parity transform then requires that the twist bundle $L \rightarrow M$ be equivariant and that the twist connection α on L is a \mathbb{Z}_2 -invariant connection. The crosscap section c is a \mathbb{Z}_2 -structure on the twist bundle, and altogether it follows that the quadruple (B, L, α, c) is subjected to the following constraints:

- (C1) $d\alpha = B + \tau^*(B)$.
- (C2) The connection $\tau^*(\alpha) - \alpha$ of $\tau^{-1}(L) \otimes L^*$ is flat and has trivial holonomy.
- (C3) c is a parallel section with respect to the connection $\tau^*(\alpha) - \alpha$, that is, $dc + (\tau^*(\alpha) - \alpha)c = 0$, such that $c\tau^*(c) = 1$.

The condition (C3) implies that $dc = 0$ and $c^2 = 1$ on the fixed point set M^τ , and so the crosscap section determines a sign choice $[c] \in H^0(M^\tau, \mathbb{Z}_2)$, which agrees with our definition of sign choice from Section 4.2. When the twist (L, α) is trivial, c is constant on M and all

O-planes have the same sign choice. For non-trivial twists, the values of c can differ from one O-plane to another.

We will now show that the twist connection α on L gives rise to a Real bundle gerbe connection on a trivial bundle gerbe $(Q, Y) = \partial_2(L)$, and the B -field gives rise to a Real curving for this connection. For this, we slightly unravel the coboundary map from the definition in [13]. Given the equivariant bundle $L \rightarrow M$, so that $\tau^{-1}(L) = L$, let $Y = M \times \mathbb{Z}_2$ with the involution $\tau: (m, x) \mapsto (\tau(m), x + 1)$ covering τ on M . Let $\pi_M: Y \rightarrow M$ be the projection. Then Y is two copies of M labelled by 0 and 1 so that any bundle $Q \rightarrow Y$ is a pair of bundles (Q_0, Q_1) on $M \times \{0\}$ and $M \times \{1\}$, respectively. For such a bundle Q we have $\tau^{-1}(Q) = (\tau^{-1}(Q_1), \tau^{-1}(Q_0))$, and if $L \rightarrow M$ is a line bundle then $\pi_M^{-1}(L) = (L, L) \rightarrow Y$. Following [13] we define the Real bundle gerbe over M which is the image of L under the connecting homomorphism as $Q \rightarrow Y^{[2]} = M \times \mathbb{Z}_2 \times \mathbb{Z}_2$ with $Q = \delta(U(1), L)$, where here $U(1)$ denotes the trivial bundle over M . Explicitly, Q is the disjoint union of four bundles over M labelled by elements of $\mathbb{Z}_2 \times \mathbb{Z}_2$ given by

$$Q_{(0,0)} = U(1), \quad Q_{(0,1)} = L, \quad Q_{(1,0)} = L^* \quad \text{and} \quad Q_{(1,1)} = U(1).$$

The bundle gerbe multiplication is made up of the obvious contractions and the Real structure follows from

$$\begin{aligned} \tau^{-1}(Q^*)_{(0,0)} &= \tau^{-1}(Q^*_{(1,1)}) = \tau^{-1}(U(1)^*) = U(1) = Q_{(0,0)}, \\ \tau^{-1}(Q^*)_{(1,0)} &= \tau^{-1}(Q^*_{(0,1)}) = \tau^{-1}(L^*) = L^* = Q_{(1,0)}, \\ \tau^{-1}(Q^*)_{(0,1)} &= \tau^{-1}(Q^*_{(1,0)}) = \tau^{-1}(L) = L = Q_{(0,1)}, \\ \tau^{-1}(Q^*)_{(1,1)} &= \tau^{-1}(Q^*_{(0,0)}) = \tau^{-1}(U(1)^*) = U(1) = Q_{(1,1)}. \end{aligned}$$

We can pick a connection α on L which is invariant by picking any connection and averaging it. Being invariant it automatically satisfies the conditions (C2) and (C3) above. Then the bundle Q inherits a connection which is a bundle gerbe connection and also Real. If we call this bundle gerbe connection A^Q , then on the four components of $Y^{[2]}$ we have

$$A^Q_{(0,0)} = 0, \quad A^Q_{(0,1)} = \alpha, \quad A^Q_{(1,0)} = -\alpha \quad \text{and} \quad A^Q_{(1,1)} = 0,$$

so its curvature $F^Q = dA^Q$ has components

$$F^Q_{(0,0)} = 0, \quad F^Q_{(0,1)} = d\alpha, \quad F^Q_{(1,0)} = -d\alpha \quad \text{and} \quad F^Q_{(1,1)} = 0.$$

We can take the curving on Y to have components $f^Q = (-d\alpha, d\alpha)$. The B -field is a two-form on M satisfying $B + \tau^*(B) = d\alpha$. If we define a two-form f on Y by $f = (f_0, f_1) = (B, -\tau^*(B))$, then to be a curving we would need $F^Q_{(i,j)} = f_j - f_i$, which is indeed satisfied.

7.2. Twisted K -theory. Gao and Hori use the orientifold data (B, L, α, c) in [9] to introduce a new variant of KR -theory groups $KR(M; c)$ accommodating non-constant sign choices, and conjecture a certain Fredholm module formulation for this K -theory. We will show that the K -theory of vector bundles *twisted by equivariant line bundles* introduced in [9] is simply KR -theory twisted by a certain Real bundle gerbe. From our general twisted KR -theory construction from [13], this implies in particular that the twisted K -theory construction of [9] is a generalised cohomology theory, and the Fredholm module description follows immediately.

First we note that there is a natural notion of modules associated to each invariant p -gerbe for $p = -1, 0, 1$. These modules form a semi-group, whose Grothendieck group completion determines a K -theory group. The respective notions of module are defined as follows.

Definition 7.1. Let (M, τ) be an orientifold.

- (a) If $c: M \rightarrow U(1)$ is a Real function, then a c -module is a vector bundle $E \rightarrow M$ with a conjugate linear map $\tau_E: E \rightarrow E$ covering τ and satisfying $\tau_E^2 = c \text{id}_E$. In particular if $c = 1$ then E is just a Real vector bundle in the sense of Atiyah [2].
- (b) If $L \rightarrow M$ is a \mathbb{Z}_2 -equivariant line bundle, then an L -module is a vector bundle $E \rightarrow M$ with a linear map $E \rightarrow \tau^{-1}(E^*) \otimes L$ such that the composition

$$E \longrightarrow \tau^{-1}(E^*) \otimes L \longrightarrow E \otimes \tau^{-1}(L^*) \otimes L$$

is $\text{id}_E \otimes c$, where $c \in \Gamma(\tau^{-1}(L^*) \otimes L)$ is the \mathbb{Z}_2 -structure on L .

- (c) If $\mathcal{G} = (P, Y) \Rightarrow M$ is a Real bundle gerbe, then a \mathcal{G} -module is a vector bundle $E \rightarrow Y$ with a bundle map $P \otimes \pi_2^{-1}(E) \rightarrow \pi_1^{-1}(E)$ over $Y^{[2]}$ satisfying the natural associativity condition that the diagram

$$\begin{array}{ccc}
 & P_{(y_1, y_2)} \otimes P_{(y_2, y_3)} \otimes E_{y_3} & \\
 \swarrow & & \searrow \\
 P_{(y_1, y_2)} \otimes E_{y_2} & & P_{(y_1, y_3)} \otimes E_{y_3} \\
 \searrow & & \swarrow \\
 & E_{y_1} &
 \end{array}$$

commutes on any triple $(y_1, y_2, y_3) \in Y^{[3]}$, together with a conjugate linear map $\tau_E: E \rightarrow E$ commuting with the Real structure on Y such that the diagram

$$\begin{array}{ccc}
 P_{(y_1, y_2)} \otimes E_{y_2} & \longrightarrow & E_{y_1} \\
 \tau \otimes \tau_E \downarrow & & \downarrow \tau_E \\
 P_{(\tau(y_1), \tau(y_2))}^* \otimes E_{\tau(y_2)}^* & \longrightarrow & E_{\tau(y_1)}^*
 \end{array}$$

commutes for every pair $(y_1, y_2) \in Y^{[2]}$.

The case of equivariant line bundle modules is the construction of Gao and Hori from [9], where the orientifold isomorphism $E \rightarrow \tau^{-1}(E^*) \otimes L$ is the action of the parity transform on the Chan–Paton bundle E . The case of Real bundle gerbe modules leads to the geometric realisation $KR_{\text{bg}}(M, P)$ of twisted KR -theory from [13].

Proposition 7.2. *There is an isomorphism between the K -theory of invariant p -gerbes \mathcal{G}_p for $p = -1, 0$, and the K -theory of their image $\partial_{p+2}\mathcal{G}_p$ under the connecting homomorphism.*

Proof. The proof is simply a matter of translating what the respective modules mean in each case and verifying that they agree.

First we consider the relationship between (a) and (b). If L is the trivial bundle on M , then it carries a natural \mathbb{Z}_2 -action and any other action is obtained by multiplication by a Real function c . Indeed this construction is the coboundary map ∂_1 in (2.4). Conversely starting with such a bundle L , the construction in (b) reduces to (a).

Next we look at the relationship between equivariant line bundles and Real bundle gerbes. Suppose that $L \rightarrow M$ is an equivariant line bundle. The coboundary map $(Q, Y) = \partial_2(L)$ was constructed explicitly in Section 7.1 above. Let F be a Real bundle gerbe module for Q . Then $\tau^{-1}(F) = F^*$, so it has the form $F = (F_0, F_1) = (E, \tau^{-1}(E^*))$ for a vector bundle $E \rightarrow M$. The

Real bundle gerbe multiplication gives isomorphisms $Q_{(i,j)} \otimes F_j \rightarrow F_i$ which are associative in the sense that the map

$$Q_{(k,i)} \otimes (Q_{(i,j)} \otimes F_j) \longrightarrow Q_{(k,i)} \otimes F_i \longrightarrow F_k$$

is equal to

$$(Q_{(k,i)} \otimes Q_{(i,j)}) \otimes F_j \longrightarrow Q_{(k,j)} \otimes F_j \longrightarrow F_k .$$

Using the definitions of Q and F , we require the isomorphisms $L \otimes \tau^{-1}(E^*) \rightarrow E$ or $E \rightarrow L \otimes \tau^{-1}(E^*)$. The condition on

$$E \longrightarrow \tau^{-1}(E^*) \otimes L \longrightarrow E \otimes \tau^{-1}(L^*) \otimes L$$

follows from the associativity conditions on the bundle gerbe module, which arise from the fact that $F_1 \simeq Q_{(1,0)} \otimes Q_{(0,1)} \otimes F_1 \simeq F_1$ is the identity map and the \mathbb{Z}_2 -action $\tau^{-1}(L^*) = L^*$. Furthermore, it is straightforward to check that this argument can be reversed, so that a module E for L defines a module $(E, \tau^{-1}(E^*))$ for the bundle gerbe Q .

In both of these cases it is clear that the bijection we have established is an isomorphism of semi-groups, and thus establishes an isomorphism of the corresponding Grothendieck groups. \square

Remark 7.3. The modules on M for an equivariant line bundle (L, c) are the same as the Real bundle gerbe modules introduced in [13] for the complex trivial bundle gerbe Q with non-trivial Real structure given by (L, c) . In other words, the Real bundle gerbe modules for the bundle gerbe Q restrict to orthogonal and quaternionic bundles on the different connected components of the fixed point set of (M, τ) , and in particular $KR_{\text{bg}}(M, Q) \simeq KR(M; c)$.

REFERENCES

- [1] C. Angelantonj and A. Sagnotti. Open strings. *Phys. Rept.*, 371:1–150, 2002.
- [2] M. F. Atiyah. K -theory and reality. *Quart. J. Math. Oxford Ser. (2)*, 17:367–386, 1966.
- [3] O. Bergman, E. G. Gimon, and S. Sugimoto. Orientifolds, RR torsion and K -theory. *J. High Energy Phys.*, no. 05, Paper 047, 30 pp, 2001.
- [4] A. L. Carey, S. Johnson, M. K. Murray, D. Stevenson, and B.-L. Wang. Bundle gerbes for Chern–Simons and Wess–Zumino–Witten theories. *Commun. Math. Phys.*, 259(3):577–613, 2005.
- [5] J. de Boer, R. Dijkgraaf, K. Hori, A. Keurentjes, J. Morgan, D. R. Morrison, and S. Sethi. Triples, fluxes, and strings. *Adv. Theor. Math. Phys.*, 4:995–1186, 2002.
- [6] J. Distler, D. S. Freed, and G. W. Moore. Orientifold précis. *Proc. Symp. Pure Math.*, 83:159–171, 2011.
- [7] C. Doran, S. Méndez-Diez, and J. Rosenberg. T-duality for orientifolds and twisted KR -theory. *Lett. Math. Phys.*, 104(11):1333–1364, 2014.
- [8] C. Doran, S. Méndez-Diez, and J. Rosenberg. String theory on elliptic curve orientifolds and KR -theory. *Commun. Math. Phys.*, 335(2):955–1001, 2015.
- [9] D. Gao and K. Hori. On the structure of the Chan–Paton factors for D-branes in Type II orientifolds. Preprint arXiv:1004.3972, 2010.
- [10] K. Gawedzki, R. R. Suszek, and K. Waldorf. Bundle gerbes for orientifold sigma-models. *Adv. Theor. Math. Phys.*, 15(3):621–688, 2011.
- [11] A. Grothendieck. Sur quelques points d’algèbre homologique II. *Tohoku Math. J. (2)*, 9(3):119–221, 1957.
- [12] S. Gukov. K -theory, reality and orientifolds. *Commun. Math. Phys.*, 210:621–639, 2000.
- [13] P. Hekmati, M. K. Murray, R. J. Szabo, and R. F. Vozzo. Real bundle gerbes, orientifolds and twisted KR -homology. *Adv. Theor. Math. Phys.*, 23(8), 2093–2159, 2019.
- [14] K. Hori. D-branes, T-duality and index theory. *Adv. Theor. Math. Phys.*, 3:281–342, 1999.
- [15] A. Keurentjes. Classifying orientifolds by flat n -gerbes. *J. High Energy Phys.*, no. 07, Paper 10, 36 pp, 2001.
- [16] E. M. Moutouou. Twisted groupoid KR -theory. Ph.D. Thesis, Université de Lorraine-Metz and Universität Paderborn, 2012.
- [17] M. K. Murray. Bundle gerbes. *J. London Math. Soc. (2)*, 54(2):403–416, 1996.
- [18] M. K. Murray and D. Stevenson. Bundle gerbes: stable isomorphism and local theory. *J. London Math. Soc. (2)*, 62(3):925–937, 2000.

- [19] K. Olsen and R. J. Szabo. Constructing D-branes from K -theory. *Adv. Theor. Math. Phys.*, 3:889–1025, 1999.
- [20] U. Schreiber, C. Schweigert and K. Waldorf. Unoriented WZW models and holonomy of bundle gerbes. *Commun. Math. Phys.*, 274(1):31–64, 2007.
- [21] D. Stevenson. Bundle 2-gerbes. *Proc. London Math. Soc. (3)*, 88(2):405–435, 2004.
- [22] E. Witten. D-branes and K -theory. *J. High Energy Phys.*, no. 12, Paper 019, 40 pp, 1998.

(Pedram Hekmati) DEPARTMENT OF MATHEMATICS, UNIVERSITY OF AUCKLAND, 1010 AUCKLAND, NEW ZEALAND

E-mail address: `p.hekmati@auckland.ac.nz`

(Michael K. Murray) SCHOOL OF MATHEMATICAL SCIENCES, UNIVERSITY OF ADELAIDE, ADELAIDE, SA 5005, AUSTRALIA

E-mail address: `michael.murray@adelaide.edu.au`

(Richard J. Szabo) DEPARTMENT OF MATHEMATICS, MAXWELL INSTITUTE FOR MATHEMATICAL SCIENCES AND THE HIGGS CENTRE FOR THEORETICAL PHYSICS, HERIOT-WATT UNIVERSITY, EDINBURGH, EH14 4AS, UNITED KINGDOM

DIPARTIMENTO DI SCIENZE E INNOVAZIONE TECNOLOGICA, AND INFN TORINO, GRUPPO COLLEGATO DI ALESSANDRIA, UNIVERSITÀ DEL PIEMONTE ORIENTALE, ALESSANDRIA, ITALY, AND ARNOLD–REGGE CENTRE, TORINO, ITALY

E-mail address: `r.j.szabo@hw.ac.uk`

(Raymond F. Vozzo) SCHOOL OF MATHEMATICAL SCIENCES, UNIVERSITY OF ADELAIDE, ADELAIDE, SA 5005, AUSTRALIA

E-mail address: `raymond.vozzo@adelaide.edu.au`