

Review

Risk factors for acquisition of multidrug-resistant Enterobacterales among international travellers: a synthesis of cumulative evidence

Luis Furuya-Kanamori^{1,*}, Jennifer Stone¹, Laith Yakob², Martyn Kirk¹, Peter Collignon^{3,4}, Deborah J Mills^{5,†} and Colleen L Lau^{1,5,†}

¹Research School of Population Health, ANU College of Health and Medicine, Australian National University, Canberra, Australia, ²Department of Disease Control, London School of Hygiene and Tropical Medicine, London, UK, ³ACT Pathology, Canberra Hospital, Canberra, Australia, ⁴ANU Medical School, ANU College of Health and Medicine, Australian National University, Canberra, Australia, and ⁵Travel Medicine Alliance Clinics, Brisbane, Australia

*To whom correspondence should be addressed. Email: luis.furuya-kanamori@anu.edu.au

†These authors contributed equally to this work

Submitted 29 August 2019; Revised 13 October 2019; Editorial Decision 21 October 2019; Accepted 21 October 2019

Abstract

Background: Recent studies have shown that over 50% of people travelling to Southeast Asia return colonized with multidrug-resistant Enterobacterales (MRE) including carbapenemase-producing Enterobacterales. Importation of MRE by travellers and subsequent spread to family members, communities and healthcare facilities poses real risks that have not yet been adequately assessed. This systematic review and meta-analysis aims to quantify the risk factors and interventions for reducing the risk of MRE acquisition among international travellers.

Methods: A systematic search was conducted in PubMed, Web of Science and Scopus for analytical epidemiological studies containing data post-2000 that assessed the risk factors to acquire and/or interventions to reduce the risk of MRE acquisition in travellers. Two researchers independently screened all the studies and extracted the information, and disagreements were resolved through consensus. The proportions of MRE acquisition by the region of destination and the odds ratio (OR) for the different risk factors and/or interventions were pooled using the inverse variance heterogeneity model.

Results: A total of 20 studies (5253 travellers from high-income countries) were included in the meta-analysis. South Asia [58.7%; 95% confidence interval (CI), 44.5–72.5%] and Northern Africa (43.9%; 95% CI 37.6–50.3%) were the travel destinations with the highest proportion of MRE acquisition. Inflammatory bowel disease (OR 2.1; 95% CI 1.2–3.8), use of antibiotics (OR 2.4; 95% CI 1.9–3.0), traveller's diarrhoea (OR 1.7; 95% CI 1.3–2.3) and contact with the healthcare system overseas (OR 1.5; 95% CI 1.1–2.2) were associated with MRE colonization. Vegetarians (OR 1.4; 95% CI 1.0–2.0) and backpackers (OR 1.5; 95% CI 1.2–1.8) were also at increased odds of MRE colonization. Few studies ($n = 6$) investigated preventive measures and found that consuming only bottled water/beverages, meticulous hand hygiene and probiotics had no protective effect on MRE colonization.

Conclusions: International travel is an important driver for MRE spread worldwide. Future research needs to identify effective interventions to reduce the risk of MRE acquisition as well as design strategies to reduce local transmission on return.

Registration: PROSPERO CRD42018076853.

Key words: Antibiotic, antimicrobial, resistance, transmission, carriage, meta-analysis, enterobacteriaceae

Introduction

There is a large clinical and public health burden associated with antimicrobial resistance (AMR) in bacteria causing human infections. This burden is increasing over time and urgent action is required.^{1,2} In a report commissioned by the UK Prime Minister, it was estimated that globally AMR could lead to 10 million excess deaths per year and a cumulative cost of US\$100 trillion by 2050.³ The problem of AMR is much more pronounced in low-income countries with poor water quality and sanitation, uncontrolled use of antibiotics, inadequate infrastructure and poor governance (i.e. corruption) all being important contributors.^{4,5}

Although carriage of multidrug-resistant Enterobacterales (MRE) is much more prevalent in communities in low-income countries (mainly in Southeast Asia, Western Pacific and Eastern Mediterranean regions),⁶ high-income countries are not immune to this problem⁷ (particularly in hospitals and long-term care facilities where there is high prevalence of MRE), and multi-resistant bacteria may also spread from country to country.⁸ Global trade of food (including live animals) and international travel play major roles in the spread and transmission of resistant organisms.⁹ The number of persons crossing international borders has continued to increase over many decades, which has provided an important opportunity for dissemination of MRE to other countries. Asymptomatic carriers can transport MRE between countries in their gut at alarming rates, e.g. studies have found that 89% of travellers returning from India and Sri Lanka to the UK were MRE carriers.¹⁰ These travellers, including medical tourists,¹¹ can transmit organisms to household contacts^{12,13} and may introduce the organisms into the community,¹⁴ healthcare facilities¹⁵ and the environment. Once established, these resistant organisms are extremely expensive and difficult to control.

In addition to the public health perspective of the risk of introduction of multidrug-resistant pathogens, special clinical considerations should be taken with returning travellers who are carriers of MRE. For example, studies have reported an association between international travellers and extended-spectrum β -lactamase-producing (ESBL) *Escherichia coli* bacteraemia post trans-rectal ultrasound guided prostate biopsy^{16,17} raising the question of the role of screening and antibiotic prophylaxis in this group of patients.

In 2015, Hassing *et al.* conducted a systematic review that compiled the evidence of MRE detection among international travellers.¹⁸ However, they did not quantitatively analyse the data. Additionally, since the publication of the systematic review, findings from the COMBAT study,¹³ the largest study on MRE in travellers with over 2000 participants, have been published. In this systematic review and meta-analysis, we update the evidence by including the most recent studies and quantifying the risk factors for MRE colonization among international travellers.

Methods

The protocol was prospectively registered in PROSPERO (CRD42018076853), and the findings of this systematic review and meta-analysis are presented according to PRISMA reporting guidelines.¹⁹

Search strategy and selection criteria

The original search strategy was designed in PubMed and converted for use in Web of Science and Scopus. The search included all articles published from 2000 until February 2019, and the search was updated in July 2019. Search terms related to 'travel', 'Enterobacteriaceae' and 'drug resistance' were included. To achieve a comprehensive evaluation of the published evidence, the systematic search was supplemented with a forwards and backwards citation search as well as retrieving the first 20 similar articles from PubMed for each of the studies included from the initial search. In addition, all references from relevant previous systematic reviews were hand searched to identify possible missed studies.

Eligible studies were analytical epidemiological studies among international travellers that reported the rate or risk (and protective) factors (i.e. compared MRE carriers vs non-carriers) for MRE acquisition (e.g. due to contaminated food), colonization (e.g. due to an underlying condition such as inflammatory bowel disease) or selection (due to use of antibiotics) on return, where data were available in an extractable format. Among the risk factors of interest were demographic characteristics, travel characteristics, past medical history, medication use and medical problems while overseas, food exposure and reason for travel. Potential protective factors included vaccines, probiotics and hand hygiene. MRE included ESBL, AmpC β -lactamase or carbapenemase-producing Enterobacterales. For a study to be included, it must have reported the country/region of travel and screened all participants on return for MRE and not just targeting symptomatic participants (e.g. with diarrhoea or fever on return). Given that the aim of the review was to describe the current situation of MRE among international travellers, only studies containing data post-2000 were included.

Studies reporting MRE acquisition among military personnel, refugees and asylum seekers, adoptees, travellers seeking medical attention, international inpatient transfers or during mass gatherings (e.g. Hajj and Olympic games) were not included as these travellers were considered at higher risk and not deemed representative of the average traveller. Exclusion criteria also included studies conducted on animals, *in vitro* studies, conference abstracts or proceedings, descriptive studies (e.g. case series) and ecological studies. No language restriction was imposed.

Study selection and data extraction

Titles and abstracts of all papers were extracted and uploaded to the Rayyan platform (<http://rayyan.qcri.org/>)²⁰ for screening. Two authors (LFK and JS) independently screened the titles and abstracts on the Rayyan platform. The same authors examined the full-text papers for eligibility in accordance with the review protocol. Any disagreements were resolved through consensus.

Data from the included studies were extracted and summarized in a spreadsheet, and the recorded fields included (i) first author and year of publication; (ii) country and study setting; (iii) study population characteristics (e.g. mean/median age, proportion of females and number of travellers); (iv) travel characteristics (e.g. travel destination and duration of travel); (v) rate of MRE detection (i.e. number of travellers that tested positive for MRE on return among the number of travellers that tested negative for MRE before travel); (vi) risk (and protective)

factors for MRE acquisition, colonization or selection; and (vii) MRE isolate characteristics (e.g. type of bacteria and resistance genes). Adjusted effect estimates and 95% confidence intervals (CIs) of the risk factors were extracted. If adjusted effect estimates were not available, unadjusted estimates were extracted or computed based on the sample size and number of events (i.e. MRE post-travel) in the exposed and unexposed groups.

Statistical analysis

Two effect measures of interest were examined, the proportion of MRE detected and the odds ratio (OR) for risk factors. Previous reviews^{18,21,22} have reported large heterogeneity in the proportion of detection across studies, suggesting the absence of a common worldwide effect estimate for the proportion of MRE detection. Countries were grouped into geographical regions [i.e. South Asia, Asia (excluding countries in South Asia), Northern Africa, sub-Saharan Africa, South and Central America, North America, Europe and Oceania] as in a previous systematic review by Hassing *et al.*¹⁸ The detailed list of countries per region can be found in the [supplementary material \(S1\)](#) (available as Supplementary data at *JTM* online). A meta-regression was fitted to examine if region of destination was a predictor for the observed heterogeneity. It was found that region of destination was the best predictor (accounted for over 45% of the observed variance) of MRE detection. Therefore, the proportions of MRE detection were pooled by the region of destination using the inverse variance heterogeneity (IVhet) model.²³ Adjusted and unadjusted (when adjusted were not available) estimates along with their 95% CIs were pooled for the different risk factors using the IVhet model. The secondary objective of the systematic review was to describe the most common bacteria isolated from the travellers as well as report the distribution of resistance genes. Proportion meta-analysis using the IVhet model was applied to pool the data from the types of bacteria and resistance genes in travellers. Statistical heterogeneity was assessed using the I^2 . Publication bias was assessed using the Doi plot and LFK index.²⁴ All the analyses were conducted in MetaXL version 5.3 (EpiGear Int Pty Ltd; Sunrise Beach; Australia; <http://www.epigear.com>).

Results

Studies identified

The search identified 725 publications; 177 articles remained after the title and abstract screening. Full-text review of 177 publications was conducted, and 28 met the eligibility criteria. It was noted that there was overlap in participants in five sets of publications. Four^{13,25–27} and two^{28,29} publications used data from the COMBAT and VOYAG-R studies, respectively. Three publications^{30–32} reported results from a Finnish cohort of travellers, two publications^{33,34} from a cohort of Australian healthcare volunteers and two publications^{35,36} from a cohort of Dutch travellers. Therefore, 20 datasets (from 28 publications) with 5253 travellers were included in this meta-analysis [supplementary material \(S2\)](#).

Characteristics of the included studies

All the studies were conducted in Europe (The Netherlands,^{12,13,35,37,38} Sweden,^{39–42} Switzerland,^{43,44} France,^{28,45}

Germany,^{38,46} Finland,³⁰ UK¹⁰ and Denmark⁴⁷) except for three that were conducted in Australia,³³ Japan^{48,49} and the USA.⁴⁹ The median age of the participants ranged from 25 to 66 years. Overall more females ($n=3028$, 57.6%) were included in the studies. The sample size ranged from 18 to 1847 participants. The median duration of travel ranged from 7 to 45 days, with 75% of the studies having a median duration of travel between 14 and 21 days. Thirteen studies collected stool samples,^{10,13,28,30,35,38–40,42,43,46,48,49} five studies used rectal swabs^{12,33,41,44,45} and two studies used a combination of both types of samples.^{37,47} Collection of the post-travel sample was done within 1 month after return in 70% of the studies. The majority of studies ($n=19$, 95%) used selective media for MRE detection that were mainly followed by genotyping confirmation, while one study only used meta-genomic approach for MRE detection³⁵ (Table 1).

Quantitative analysis

Proportion of MRE detection by region of destination. Overall, there was a wide variability in the proportion of MRE detected, ranging from 13% to 88% and considerable heterogeneity across studies was observed. Heterogeneity substantially decreased when estimates were pooled by regions. South Asia (58.7%; 95% CI 44.5–72.5%), Northern Africa (43.9%; 95% CI 37.6–50.3%) and Asia (37.5%; 95% CI 23.6–51.9%) were the travel destinations with the highest proportion of MRE detection. These regions were followed by sub-Saharan Africa (21.8%; 95% CI 12.0–32.4%), South and Central America (18.3%; 95% CI 8.8–28.9%) and North America (16.9%; 95% CI 2.6–35%); while the lowest proportion of detection were observed in Europe (10.3%; 95% CI 5.5–16.1%) and Oceania (6.9%; 95% CI 0–21.1%) (Figure 1).

Risk factors. MRE detection was not associated with sex, age or duration of travel, but it was dependent on the travel destination. In terms of past medical history, it was found that inflammatory bowel disease (OR 2.1; 95% CI 1.2–3.8) was a significant risk factor for MRE colonization. While overseas, use of antibiotics (OR 2.4; 95% CI 1.9–3.0) was the strongest risk factor; antimalarial prophylaxis (OR 1.0; 95% CI 0.8–1.4) was not associated with MRE colonization, but it should be noted that commonly used antimalarial medications include doxycycline (a broad spectrum antibiotic) as well as other medications (e.g. mefloquine, atovaquone/proguanil, chloroquine) that would not be expected to have an effect on selection of AMR Enterobacteriales. Other factors associated with MRE colonization were experiencing traveller's diarrhoea (OR 1.7; 95% CI 1.3–2.3) and having contact with the healthcare system either as inpatients or outpatients (OR 1.5; 95% CI 1.1–2.2). With regards to food exposure, having a vegetarian diet (OR 1.4; 95% CI 1.0–2.0) was found to increase the odds of acquiring MRE compared to other diets, and consuming only bottled water/beverages did not have an impact on MRE colonization. Among the interventions that were examined for their potential impact in decreasing the risk of MRE colonization, meticulous (as defined by primary study authors) hand hygiene and probiotics had no effect, while oral cholera vaccine (OR 1.6; 95% CI 1.0–2.5) was found to increase the odds of MRE colonization, possible due to confounding by indication (Discussion). Backpacker travellers had a 50%

Table 1. Characteristics of the included studies

First author, year of publication	Country	Study period	Study population	Mean/median age (in years)	Proportion of females (%)	Sample size	Type of sample	Collection of sample post-travel	Laboratory detection method	Most common travel destinations	Mean/median duration of travel (in days)	Proportion of MRE detection (%)
Angelin 2015 ⁴²	Sweden	2010–14	Healthcare students travelling for clinical courses	25	78	99	Stool sample	1–2 weeks after return	Selective media	89% Southeast Asia and Africa	45	35
Arcilla 2017 ⁴³	The Netherlands	2012–13	Patients attending travel clinics	51	54	1847	Stool sample	Immediately after return	Selective media + microarray genotyping	82% Asia and Africa	20	34
Bernasconi 2016 ⁴³	Switzerland	2015	People living in Switzerland	44	68	38	Stool sample	Within 1 week after return	Selective media + microarray genotyping	India	18	76
Bevan 2018 ¹⁰	UK	NR	Healthy volunteers	30	NR	18	Stool sample	NR	Selective media + PCR genotyping + whole genome sequencing	India and Sri Lanka	21	89
Kantele 2015 ³⁰	Finland	2009–10	Patients attending a travel clinic	40	61	430	Stool sample	First or second stool after return	Selective media + PCR genotyping	83% Asian and Africa	19	21
Kennedy 2010 ³³	Australia	2008–09	Hospital staff and contacts	45	62	102	Rectal swab	Within 2 weeks after return	Selective media + PCR genotyping	77% Southeast Asia and Europe	21	49
Kuenzli 2014 ⁴⁴	Switzerland	2012–13	Patients attending travel clinics	41	56	170	Rectal swab	Directly after return	Selective media + microarray and PCR/DNA sequence genotyping	India, Bhutan, Nepal and Sri Lanka	18	69
Lausch 2013 ⁴⁷	Denmark	2011	Inpatients with travel history in the last 3 months	37	57	88	Rectal swab and stool sample (if patient had diarrhoea)	Up to 3 months after return	Selective media	52% Asia and Africa	NR	13
Lubbert 2015 ⁴⁶	Germany	2013–14	Patients attending a travel clinic	34	57	191	Stool sample	Within 1 week after return	Selective media + PCR genotyping	88% Asia and Africa	21	30

(continued)

Table 1. Continued

First author, year of publication	Country	Study period	Study population	Mean/median age (in years)	Proportion of females (%)	Sample size	Type of sample	Collection of sample post-travel	Laboratory detection method	Most common travel destinations	Mean/median duration of travel (in days)	Proportion of MRE detection (%)
Macaux 2018 ⁴⁵	France	2014–16	Inpatients with travel history in the past 12 months	62	34	138	Rectal swab	Up to 12 months after return	Selective media	83% Asia, sub-Saharan Africa and the Mediterranean	NR	NA
Nakayama 2018 ⁴⁸	Japan	2015–16	Japanese travellers to Vietnam	37	42	19	Stool sample	Within 3 weeks after return	Selective media + PCR genotyping	Vietnam	7	88
Ostholm-Balkhed 2013 ³⁹	Sweden	2008–09	Patients attending vacation clinics	54	59	226	Stool sample	Up to 4 months after return	Selective media + PCR genotyping	57% Sub-Saharan Africa and Asia	16	30
Paltansing 2013 ¹²	The Netherlands	2011	Patients attending a travel clinic	33	63	338	Rectal swabs	Immediately after return	Selective media + microarray genotyping	49% Southeast Asia and Central Africa	21	33
Reuland 2016 ³⁷	The Netherlands	2012–13	Patients attending a vacation clinic	33	58	418	Rectal swab or stool sample	Within 2 weeks after return	Selective media + microarray genotyping	Africa, Asia and Latin America	14	23
Ruppe 2015 ²⁸	France	2012–13	Patients attending international vaccination centres	36	61	574	Stool sample	Within 1 week after return	Selective media + PCR genotyping	Sub-Saharan Africa, Asia and Latin America	20	50
Schaumburg 2019 ³⁸	Germany and The Netherlands	2016–18	Patients attending vaccination centres	32	59	132	Stool sample	Within 1 week after return	Selective media + molecular assay	68% Asia and Africa	18	46
Tangden 2010 ⁴⁰	Sweden	2007–09	Patients attending travel clinics	43	55	100	Stool sample	NR	Selective media + PCR genotyping	65% Africa and Asia	14	24
Vading 2016 ⁴¹	Sweden	2013–15	Patients attending a travel clinic	49	68	175	Rectal swab	NR	Selective media + PCR genotyping	Southeast Asia, India, Northern Africa and Turkey	14	38
von Wintersdorff 2014 ³⁵	The Netherlands	2010–12	Patients attending travel clinics	43	58	122	Stool sample	Immediately after return	Meta-genomic	48% Southeast Asia and India	21	32
Weisenberg 2012 ⁴⁹	USA	2009–10	Patients attending a travel clinic	66	68	28	Stool sample	Within 1 week after return	Selective media + PCR genotyping	54% Sub-Saharan Africa and South Asia	16	25

NR, not reported; NA, not applicable. The study was a matched case-control.




Figure 1. Proportion of acquisition of MRE by region of destination.

(OR 1.5; 95% CI 1.2–1.8) increased odds of acquiring MRE compared to other types of travellers (Table 2).

Aetiology and resistance genes. Of the 2276 MRE isolates, the vast majority was *E. coli* (92.0%; 95% CI 84.9–97.3%) followed by *Klebsiella pneumoniae* (5.9%; 95% CI 1.2–11.8%). The remaining 2% of isolates included *Citrobacter* spp. (*freundii* and *amalonaticus*), *Proteus* spp. (*vulgaris* and *mirabilis*), *Enterobacter cloacae*, *Enterococcus faecium*, *Acinetobacter baumannii*, *Morganella morganii* and a case of non-Typhi *Salmonella enterica*.

*bla*_{CTX-M} (92.3%; 95% CI 44.6–100%) was the most frequent resistance gene found in the MRE isolates. CTX-M group 1 was the most common followed by CTX-M Groups 9 and 4. *bla*_{SHV} and *bla*_{TEM} were found in 3.0% (95% CI 0–2.0%) and 2.0% (95% CI 0–1.8%) of the isolates, respectively. Enterobacterales with carbapenem- (*bla*_{OXA}^{28,37,45} and *bla*_{NDM}^{28,44,45}) and colistin- (*mcr*⁴⁸) resistance genes were also isolated. A total of 50 isolates with carbapenem-resistant genes were identified mainly from travellers returning from Southeast Asia and the Indian subcontinent,^{28,37,44,45} while the 3 isolates with colistin-resistant genes were from travellers returning from Vietnam.⁴⁸

Discussion

In this systematic review and meta-analysis, we compiled data from 20 different studies with over 5000 travellers to update the evidence on the proportion and risk factors for MRE detection in international travellers. The highest proportions of MRE detection were observed from travellers returning from South Asia, Northern Africa, Asia, sub-Saharan Africa and South and Central America; our findings correspond with the global epidemiology of AMR⁴ and the ESBL-producing Enterobacterales colonization prevalence among healthy individuals.⁵⁰ In a meta-analysis by Karanika *et al.*, international travellers were found

to be four times more likely to be colonized by ESBL-producing Enterobacterales than non-travellers.⁵⁰ These findings highlight the importance of international travel to low- and middle-income countries in the global spread of MRE.^{51,52} In recent years, dissemination of emerging pathogens such as NDM-1 carbapenem-resistant *K. pneumoniae* from India⁵³ and *mcr*-1 colistin-resistant *E. coli* from China⁵⁴ were facilitated by international travel.^{55,56} Of concern is that our review revealed that carbapenem-^{28,37,44,45} and colistin⁴⁸-resistant Enterobacterales are being detected in international travellers. Investment in infrastructure to provide access to clean water and adequate sanitation and hygiene may reduce transmission in low- and middle-income countries,⁴ and possibly reduce the global spread of AMR.

Contact with the healthcare system, traveller's diarrhoea and antibiotic use while travelling were strongly associated with MRE detection. The possibility that these factors are associated through reverse causality with MRE cannot be excluded (i.e. MRE infection leading to contact with the healthcare system, traveller's diarrhoea and/or use of antibiotics). Even in the event the observed associations are due to reverse causality, these factors are markers for MRE detection and can be used to identify travellers at higher risk of being colonized by MRE. Among the risk factors identified, one that travellers have control over is antibiotic consumption; thus during pre-travel consultations, healthcare professionals need to emphasize the use of antibiotics only when required.

Acute diarrhoea is the most common condition affecting travellers to South and Central Asia and the second most common among travellers worldwide.⁵⁷ Therefore, priority areas of research have been identified to generate evidence-based recommendations for preventing traveller's diarrhoea.⁵⁸ In line with the research priority areas, inflammatory bowel disease, a vegetarian diet and backpacking were identified as host risk factors for MRE detection. Although it was not possible to examine the exposure to different classes and doses of antibiotics in the

Table 2. Risk factors for acquisition of MRE

	OR (95% CI)	I ² (%)	Number of studies with adjusted estimates/total number of studies
Demographic			
Sex (female)	0.91 (0.80–1.03)	0	2/14
Age (per 10-year increase)	1.01 (0.97–1.06)	0	2/5
Travel-associated data			
Duration of travel (per week increase)	1.02 (0.99–1.06)	0	1/5
Travel destination (reference Europe)*	1	NA	0/10
South Asia	1.52 (1.31–1.76)		0/19
Northern Africa	1.33 (1.25–1.43)		0/11
Asia	1.26 (1.17–1.35)		0/16
Sub-Saharan Africa	1.12 (0.99–1.28)		0/15
Americas	1.11 (1.02–1.20)		0/16
Past medical history			
Inflammatory bowel disease	2.09 (1.16–3.77)	0	1/3
Chronic disease	1.01 (0.65–1.57)	57	2/5
Admitted to a hospital in the previous 3–6 months	1.79 (0.09–36.63)	68	1/2
Antibiotic use in the previous 3–12 months	1.00 (0.76–1.32)	12	3/4
Medical history while overseas			
Antibiotic use	2.38 (1.88–3.00)	0	4/12
Anti-acid use	1.08 (0.59–1.97)	25	0/5
Antimalarial prophylaxis	1.04 (0.75–1.44)	10	0/3
Travellers' diarrhoea	1.69 (1.25–2.30)	46	5/13
Contact with healthcare system	1.53 (1.09–2.15)	20	1/10
Food exposure			
Vegetarian diet	1.41 (1.01–1.96)	0	0/5
Only consume bottled water/beverages	1.29 (0.50–3.34)	77	1/6
Protective factors			
Oral cholera vaccine	1.61 (1.04–2.50)	0	0/2
Meticulous hand hygiene	1.10 (0.81–1.49)	0	1/4
Probiotics	1.06 (0.78–1.45)	0	0/2
Type of traveller			
Backpackers	1.46 (1.20–1.78)	0	2/7
Holiday makers	0.99 (0.70–1.38)	0	0/4
Visiting friends and relatives	0.95 (0.55–1.63)	29	0/5
Business travellers	0.78 (0.46–1.30)	0	0/4

OR, odds ratio; CI, confidence interval; NA, not applicable.

*OR obtained through meta-regression. Americas include North, Central and South America. Oceania was excluded from the analysis because too few studies ⁽³⁾ reported data for this region.

current review, findings by Ruppe *et al.*²⁸ indicate that use of different classes of antibiotics has distinct effects on the risk of MRE selection. They found that antibiotics used for malaria prophylaxis (i.e. doxycycline) did not increase the risk of MRE selection (OR 0.9; 95% CI 0.6–1.6), while beta-lactams were associated with an increase in risk (OR 4.2; 95% CI 1.5–12.1).²⁸

Non-antibiotic approaches to prevent MRE colonization were also examined. Findings from the current review align with a recent randomized controlled trial that found that *Lactobacillus rhamnosus* GG did not reduce MRE colonization in travellers.⁵⁹ The effect of oral cholera vaccine was also assessed, and paradoxically it was found that cholera vaccine was associated with an increase in the risk of MRE colonization. Considering that the oral cholera vaccine provides some cross-protection against enterotoxigenic *E. coli*, one of the most common causes of traveller's diarrhoea, this association may seem contradictory. The most likely explanation for the observation is that unadjusted estimates were analysed and the

result is affected by confounding by indication, i.e. travellers going to regions with higher risk of MRE are more likely to receive oral cholera vaccine. In our previous study of MRE detection in Australian travellers, those travelling to the Indian subcontinent were more likely to receive oral cholera vaccine than travellers going to other destinations; when the analysis was adjusted for travel destination and traveller's comorbidities, the point estimate indicated a protective effect of oral cholera vaccine against MRE colonization.⁶⁰ To prevent confounding by indication, randomized controlled trials are needed to provide robust evidence on the effect of potentially protective interventions.

It is clear that international travellers are a high-risk group; over 50% of travellers returning from South Asia are asymptomatic MRE carriers, and there is increasing evidence of subsequent household transmission of MRE.^{12,13} Cases of ESBL-producing *E. coli* bacteraemia have been reported after prostate biopsies,¹⁷ and international travel has been identified as the

risk factor for fluoroquinolone-resistant and ESBL-producing *E. coli* infections.^{16,61} Therefore, the role of screening and contact precaution of returned travellers from clinical and public health perspectives need to be further examined.⁶² It is unfeasible to screen all returning travellers, but in a healthcare setting, it may be feasible to screen those who are planning to undergo higher-risk medical procedures; although current evidence suggests that ESBL-producing *E. coli* is less likely to spread among patients in hospitals than other Enterobacterales species.⁶³ Current studies on international patient transfers have shown that targeted screening programs of high-risk patients may be a more cost-effective strategy than mass screening,⁶⁴ and contact precaution and environmental cleaning with chlorine-based disinfectant can prevent transmission.⁶⁵ Given that prolonged MRE carriage is uncommon, most travellers (~90%) are culture-negative within 6 months,^{28,33,46} strategies should aim to reduce transmission and avoid elective surgical interventions within this period.

The findings of our systematic review and meta-analysis should be understood in light of some of the limitations. Estimates extracted and synthesized from observational studies included adjusted as well as unadjusted estimates; thus pooled estimates may be affected by confounding variables. This may be the case of potential protective interventions, where paradoxical findings were observed. To examine regional and within-country differences in MRE detection, as well as the risk of different antibiotic classes and doses, more granular data are required. Individual patient data meta-analysis may provide valuable information to answer these questions. The current review examined the risk factors relevant to average travellers from high-income countries; thus findings may not be applicable to special groups of travellers such as military personnel, refugees and asylum seekers or hospital transfers as they may have different risk profiles. The median age of travellers in the included studies ranged from 25 to 66 years, so findings may not be generalizable to children. Some studies collected post-travel samples between 3 and 12 months post travel; thus we cannot rule out the possibility that the participants became colonized after their return.

In conclusion, international travel is an important driver for MRE spread worldwide. South Asia as a travel destination is a major risk factor for MRE acquisition. Additional risk factors for individuals to become colonized include inflammatory bowel disease, use of antibiotics, traveller's diarrhoea, contact with the healthcare system, having a vegetarian diet and backpacking. Future research needs to identify effective interventions to reduce the risk of MRE colonization and design strategies to reduce local transmission on return, including transmission to household and community members, as well as wider dissemination into the environment and animal populations.

Supplementary Data

Supplementary data are available at *JTM* online.

Acknowledgements

The authors acknowledge the assistance provided by Mr Tsheten Tsheten for creating the choropleth map.

Authors' contributions

Conception and design of the study: L.F.K., D.J.M. and C.L.L.; collection and assembly of the data: L.F.K. and J.S.; analysis and interpretation: L.F.K., L.Y., D.J.M. and C.L.L.; manuscript writing: all authors; final approval of the manuscript: all authors.

Funding

L.F.K. and C.L.L. were supported by Australian National Health and Medical Research Council Early Career Fellowships (APP1158469 and APP1109035). M.K. was supported by an Australian National Health and Medical Research Council Career Development Fellowship (APP1145997). J.S. was supported by the Australian National University, Australia Degree by Research Scholarship.

Conflicts of interest: None declared.

References

- de Kraker ME, Davey PG, Grundmann H. Mortality and hospital stay associated with resistant *Staphylococcus aureus* and *Escherichia coli* bacteremia: estimating the burden of antibiotic resistance in Europe. *PLoS Med* 2011; 8:e1001104.
- Stewardson AJ, Allignol A, Beyersmann J *et al.* The health and economic burden of bloodstream infections caused by antimicrobial-susceptible and non-susceptible Enterobacteriaceae and *Staphylococcus aureus* in European hospitals, 2010 and 2011: a multicentre retrospective cohort study. *Euro Surveill* 2016; 21:30319.
- O'Neil J. Antimicrobial resistance: tackling a crisis for the health and wealth of nations. 2014. https://amr-review.org/sites/default/files/AMR%20Review%20Paper%20-%20Tackling%20a%20crisis%20for%20the%20health%20and%20wealth%20of%20nations_1.pdf (28 August 2019, date last accessed).
- Collignon P, Beggs JJ, Walsh TR *et al.* Anthropological and socio-economic factors contributing to global antimicrobial resistance: a univariate and multivariable analysis. *Lancet Planet Health* 2018; 2:e398–405.
- Ventola CL. The antibiotic resistance crisis: part 1: causes and threats. *P T* 2015; 40:277–83.
- Woerther PL, Burdet C, Chachaty E, Andremonet A. Trends in human fecal carriage of extended-spectrum β -lactamases in the community: toward the globalization of CTX-M. *Clin Microbiol Rev* 2013; 26:744–58.
- Klein EY, Tseng KK, Pant S, Laxminarayan R. Tracking global trends in the effectiveness of antibiotic therapy using the drug resistance index. *BMJ Glob Health* 2019; 4:e001315.
- Kumarasamy KK, Toleman MA, Walsh TR *et al.* Emergence of a new antibiotic resistance mechanism in India, Pakistan, and the UK: a molecular, biological, and epidemiological study. *Lancet Infect Dis* 2010; 10:597–602.
- Collignon P. Antibiotic resistance: are we all doomed? *Intern Med J* 2015; 45:1109–15.
- Bevan ER, McNally A, Thomas CM *et al.* Acquisition and loss of CTX-M-producing and non-producing *Escherichia coli* in the fecal microbiome of travelers to South Asia. *MBio* 2018; 9:e02408–18.
- Chen LH, Wilson ME. The globalization of healthcare: implications of medical tourism for the infectious disease clinician. *Clin Infect Dis* 2013; 57:1752–9.
- Paltansing S, Vlot JA, Kraakman ME *et al.* Extended-spectrum β -lactamase-producing Enterobacteriaceae among travelers from the Netherlands. *Emerg Infect Dis* 2013; 19:1206–13.

13. Arcilla MS, van Hattem JM, Haverkate MR *et al.* Import and spread of extended-spectrum beta-lactamase-producing Enterobacteriaceae by international travellers (COMBAT study): a prospective, multi-centre cohort study. *Lancet Infect Dis* 2017; 17:78–85.
14. Wilder-Smith A, Goh KT, Barkham T, Paton NI. Hajj-associated outbreak strain of *Neisseria meningitidis* serogroup W135: estimates of the attack rate in a defined population and the risk of invasive disease developing in carriers. *Clin Infect Dis* 2003; 36:679–83.
15. Fournier S, Lepointeur M, Kassis-Chikhani N *et al.* Link between carbapenemase-producing Enterobacteria carriage and cross-border exchanges: eight-year surveillance in a large French multihospitals institution. *J Travel Med* 2012; 19:320–3.
16. Patel U, Dasgupta P, Amoroso P *et al.* Infection after transrectal ultrasonography-guided prostate biopsy: increased relative risks after recent international travel or antibiotic use. *BJU Int* 2012; 109:1781–5.
17. Williamson DA, Masters J, Freeman J, Roberts S. Travel-associated extended-spectrum beta-lactamase-producing *Escherichia coli* bloodstream infection following transrectal ultrasound-guided prostate biopsy. *BJU Int* 2012; 109:E21–2.
18. Hassing RJ, Alsmas J, Arcilla MS *et al.* International travel and acquisition of multidrug-resistant Enterobacteriaceae: a systematic review. *Euro Surveill* 2015; 20:30074.
19. Moher D, Liberati A, Tetzlaff J *et al.* Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med* 2009; 6:e1000097.
20. Ouzzani M, Hammady H, Fedorowicz Z, Elmagarmid A. Rayyan—a web and mobile app for systematic reviews. *Syst Rev* 2016; 5:210.
21. Ruppé E, Andremont A, Armand-Lefevre L. Digestive tract colonization by multidrug-resistant Enterobacteriaceae in travellers: an update. *Travel Med Infect Dis* 2018; 21:28–35.
22. Armand-Lefevre L, Andremont A, Ruppé E. Travel and acquisition of multidrug-resistant Enterobacteriaceae. *Med Mal Infect* 2018; 48:431–41.
23. Doi SA, Barendregt JJ, Khan S *et al.* Advances in the meta-analysis of heterogeneous clinical trials I: the inverse variance heterogeneity model. *Contemp Clin Trials* 2015; 45:130–8.
24. Furuya-Kanamori L, Barendregt JJ, Doi SAR. A new improved graphical and quantitative method for detecting bias in meta-analysis. *Int J Evid Based Healthc* 2018; 16:195–203.
25. Van Hattem JM, Arcilla MS, Bootsma MCJ *et al.* Prolonged carriage and potential onward transmission of carbapenemase-producing Enterobacteriaceae in Dutch travelers. *Future Microbiol* 2016; 11:857–64.
26. van Hattem JM, Arcilla MS, Grobusch MP *et al.* Travel-related acquisition of diarrhoeagenic bacteria, enteric viruses and parasites in a prospective cohort of 98 Dutch travellers. *Travel Med Infect Dis* 2017; 19:33–6.
27. van Hattem JM, Cabal A, Arcilla MS *et al.* Fenic *Escherichia coli* virulence genes in intercontinental travellers: a prospective, multi-centre study. *Travel Med Infect Dis* 2019.
28. Ruppé E, Armand-Lefevre L, Estellat C *et al.* High rate of acquisition but short duration of carriage of multidrug-resistant Enterobacteriaceae after travel to the tropics. *Clin Infect Dis* 2015; 61: 593–600.
29. Lorme F, Maataoui N, Rondinaud E *et al.* Acquisition of plasmid-mediated cephalosporinase producing Enterobacteriaceae after a travel to the tropics. *PLoS One* 2018; 13:e0206909.
30. Kantele A, Laaveri T, Mero S *et al.* Antimicrobials increase travelers' risk of colonization by extended-spectrum beta-lactamase-producing Enterobacteriaceae. *Clin Infect Dis* 2015; 60:837–46.
31. Kantele A, Mero S, Kirveskari J, Laaveri T. Fluoroquinolone antibiotic users select fluoroquinolone-resistant ESBL-producing Enterobacteriaceae (ESBL-PE) - data of a prospective traveller study. *Travel Med Infect Dis* 2017; 16:23–30.
32. Lääveri T, Vilkinen K, Pakkanen S *et al.* Despite antibiotic treatment of travellers' diarrhoea, pathogens are found in stools from half of travellers at return. *Travel Med Infect Dis* 2018; 23:49–55.
33. Kennedy K, Collignon P. Colonisation with *Escherichia coli* resistant to "critically important" antibiotics: a high risk for international travellers. *Eur J Clin Microbiol Infect Dis* 2010; 29:1501–6.
34. Rogers BA, Kennedy KJ, Sidjabat HE *et al.* Prolonged carriage of resistant *E. coli* by returned travellers: clonality, risk factors and bacterial characteristics. *Eur J Clin Microbiol Infect Dis* 2012; 31:2413–20.
35. von Wintersdorff CJ, Penders J, Stobberingh EE *et al.* High rates of antimicrobial drug resistance gene acquisition after international travel, The Netherlands. *Emerg Infect Dis* 2014; 20:649–57.
36. von Wintersdorff CJH, Wolfs PFG, van Niekerk JM *et al.* Detection of the plasmid-mediated colistin-resistance gene *mcr-1* in faecal metagenomes of Dutch travellers. *J Antimicrob Chemother* 2016; 71:3416–9.
37. Reuland EA, Sonder GJ, Stolte I *et al.* Travel to Asia and traveller's diarrhoea with antibiotic treatment are independent risk factors for acquiring ciprofloxacin-resistant and extended spectrum beta-lactamase-producing Enterobacteriaceae—a prospective cohort study. *Clin Microbiol Infect* 2016; 22:731.e1–7.
38. Schaumburg F, Sertic SM, Correa-Martinez C *et al.* Acquisition and colonization dynamics of antimicrobial-resistant bacteria during international travel: a prospective cohort study. *Clin Microbiol Infect* 2019; 25:1287.e1–7.
39. Östholm-Balkhed A, Tärnberg M, Nilsson M *et al.* Travel-associated faecal colonization with *esbl*-producing enterobacteriaceae: incidence and risk factors. *J Antimicrob Chemother* 2013; 68:2144–53.
40. Tängdén T, Cars O, Melhus Å, Löwdin E. Foreign travel is a major risk factor for colonization with *Escherichia coli* producing CTX-M-type extended-spectrum β -lactamases: a prospective study with Swedish volunteers. *Antimicrob Agents Chemother* 2010; 54:3564–8.
41. Vading M, Kabir MH, Kalin M *et al.* Frequent acquisition of low-virulence strains of ESBL-producing *Escherichia coli* in travellers. *J Antimicrob Chemother* 2016; 71:3548–55.
42. Angelin M, Forsell J, Granlund M *et al.* Risk factors for colonization with extended-spectrum beta-lactamase producing Enterobacteriaceae in healthcare students on clinical assignment abroad: a prospective study. *Travel Med Infect Dis* 2015; 13:223–9.
43. Bernasconi OJ, Kuenzli E, Pires J *et al.* Travelers can import colistin-resistant Enterobacteriaceae, including those possessing the plasmid-mediated *mcr-1* gene. *Antimicrob Agents Chemother* 2016; 60:5080–4.
44. Kuenzli E, Jaeger VK, Frei R *et al.* High colonization rates of extended-spectrum beta-lactamase (ESBL)-producing *Escherichia coli* in Swiss travellers to South Asia—a prospective observational multicentre cohort study looking at epidemiology, microbiology and risk factors. *BMC Infect Dis* 2014; 14:528.
45. Macaux L, Ndoye O, Cordel H *et al.* Extensively-drug-resistant bacteria carriers among overseas travellers: one-third had not been hospitalized previously. *Int J Antimicrob Agents* 2018; 52:385–9.
46. Lübbert C, Straube L, Stein C *et al.* Colonization with extended-spectrum beta-lactamase-producing and carbapenemase-producing Enterobacteriaceae in international travelers returning to Germany. *Int J Med Microbiol* 2015; 305:148–56.
47. Lausch KR, Fuursted K, Larsen CS, Storgaard M. Colonisation with multi-resistant Enterobacteriaceae in hospitalised Danish patients with a history of recent travel: a cross-sectional study. *Travel Med Infect Dis* 2013; 11:320–3.

48. Nakayama T, Kumeda Y, Kawahara R *et al.* Carriage of colistin-resistant, extended-spectrum β -lactamase-producing *Escherichia coli* harboring the mcr-1 resistance gene after short-term international travel to Vietnam. *Infect Drug Resist* 2018; 11: 391–5.
49. Weisenberg SA, Mediavilla JR, Chen L *et al.* Extended spectrum beta-lactamase-producing Enterobacteriaceae in international travelers and non-travelers in New York city. *PLoS One* 2012; 7: e45141.
50. Karanika S, Karantanos T, Arvanitis M *et al.* Fecal colonization with extended-spectrum beta-lactamase-producing Enterobacteriaceae and risk factors among healthy individuals: a systematic review and Metaanalysis. *Clin Infect Dis* 2016; 63:310–8.
51. Frost I, Van Boeckel TP, Pires J *et al.* Global geographic trends in antimicrobial resistance: the role of international travel. *J Travel Med* 2019; taz036.
52. Vila J. Multidrug-resistant bacteria without borders: role of international trips in the spread of multidrug-resistant bacteria. *J Travel Med* 2015; 22:289–91.
53. Yong D, Toleman MA, Giske CG *et al.* Characterization of a new metallo-beta-lactamase gene, bla(NDM-1), and a novel erythromycin esterase gene carried on a unique genetic structure in *Klebsiella pneumoniae* sequence type 14 from India. *Antimicrob Agents Chemother* 2009; 53:5046–54.
54. Liu YY, Wang Y, Walsh TR *et al.* Emergence of plasmid-mediated colistin resistance mechanism MCR-1 in animals and human beings in China: a microbiological and molecular biological study. *Lancet Infect Dis* 2016; 16:161–8.
55. Berrazeg M, Diene S, Medjahed L *et al.* New Delhi Metallo-beta-lactamase around the world: an eReview using Google Maps. *Euro Surveill* 2014; 19:20809.
56. Schwarz S, Johnson AP. Transferable resistance to colistin: a new but old threat. *J Antimicrob Chemother* 2016; 71:2066–70.
57. Freedman DO, Weld LH, Kozarsky PE *et al.* Spectrum of disease and relation to place of exposure among ill returned travelers. *N Engl J Med* 2006; 354:119–30.
58. DuPont HL, Steffen R. Use of antimicrobial agents for treatment and prevention of travellers' diarrhoea in the face of enhanced risk of transient fecal carriage of multi-drug resistant enterobacteriaceae: setting the stage for consensus recommendations. *J Travel Med* 2017; 24:S57–s62.
59. Dall LB, Lausch KR, Gedebjerg A *et al.* Do probiotics prevent colonization with multi-resistant Enterobacteriaceae during travel? A randomized controlled trial. *Travel Med Infect Dis* 2019; 27:81–6.
60. Furuya-Kanamori L, Mills DJ, Robson J, Lau CL. Acquisition rate and risk factors for multidrug resistance Enterobacteriaceae among Australian travellers. Poster presentation at the 16th Conference of the ISTM Washington DC, USA: International Society of Travel Medicine, 2019.
61. Knaapila J, Kallio H, Hakanen AJ *et al.* Antibiotic susceptibility of intestinal *Escherichia coli* in men undergoing transrectal prostate biopsies: a prospective, registered, multicentre study. *BJU Int* 2018; 122:203–10.
62. Petersen E, Mohsin J. Should travelers be screened for multi-drug resistant (MDR) bacteria after visiting high risk areas such as India? *Travel Med Infect Dis* 2016; 14:591–4.
63. Duval A, Obadia T, Boëlle P-Y *et al.* Close proximity interactions support transmission of ESBL-K. pneumoniae but not ESBL-E. coli in healthcare settings. *PLoS Comput Biol* 2019; 15:e1006496-e.
64. Kaspar T, Schweiger A, Droz S, Marschall J. Colonization with resistant microorganisms in patients transferred from abroad: who needs to be screened? *Antimicrob Resist Infect Control* 2015; 4:31. doi: 10.1186/s13756-015-0071-6.
65. Chua KY, Grayson ML, Burgess AN *et al.* The growing burden of multidrug-resistant infections among returned Australian travellers. *Med J Aust* 2014; 200:116–8.