

This is a pre-copyedited, author-produced PDF of an article accepted for publication in Quarterly Journal of Mathematics, following peer review. The version of record
Michael K. Murray and Paul Norbury

JNR monopolies

Quarterly Journal of Mathematics, 2021; 72(1-2):387-405, is available online at:
<http://dx.doi.org/10.1093/qmath/haaa033>

© The Author(s) 2020. Published by Oxford University Press. All rights reserved.

PERMISSIONS

https://academic.oup.com/journals/pages/self_archiving_policy_b

Accepted Manuscript

The accepted manuscript (AM) is the final draft author manuscript, as accepted for publication by a journal, including modifications based on referees' suggestions, before it has undergone copyediting, typesetting and proof correction. This is sometimes referred to as the post-print version.

Immediately upon publication authors may:

- Immediately upload their AM to their own personal webpage (excluding commercial websites and repositories)
- Immediately upload their AM to their institutional or other non-commercial subject based repositories on the proviso that it is not made publicly available until after the specified embargo period

After the embargo period authors may:

Upload their AM to institutional repository or other non-commercial repositories and make it publicly available. Accepted Manuscripts may not be uploaded or shared on commercial websites or repositories, unless the website or repository has signed a licensing agreement with OUP permitting such uploading or sharing.

Embargo periods

Embargo periods may vary between journals. For details of a journal's specific embargo period, please see the information for each individual title on our [Accepted Manuscript embargo](#) page.

When uploading an accepted manuscript to a repository, authors should include the following acknowledgment as well as a link to the version of record. This will connect the published version to the AM version in the repository and help ensure that the article is cited correctly.

This is a pre-copyedited, author-produced version of an article accepted for publication in [insert journal title] following peer review. The version of record [insert complete citation information here] is available online at: xxxxxx [insert URL and DOI of the article on the OUP website].

17 July 2024

<http://hdl.handle.net/2440/132678>

JNR MONOPOLES

MICHAEL K. MURRAY AND PAUL NORBURY

ABSTRACT. We review the theory of JNR, mass $\frac{1}{2}$ hyperbolic monopoles in particular their spectral curves and rational maps. These are used to establish conditions for a spectral curve to be the spectral curve of a JNR monopole and to show that the rational map of a JNR monopole arises by scattering using results of Atiyah. We show that for JNR monopoles the holomorphic sphere has a remarkably simple form and show that this can be used to give a formula for the energy density at infinity. In conclusion we illustrate some examples of the energy density at infinity of JNR monopoles.

Dedicated to the memory of Sir Michael Atiyah

CONTENTS

1. Introduction	2
2. Hyperbolic monopoles	4
3. JNR monopoles	5
3.1. JNR spectral curve	5
3.2. Scattering	7
3.3. JNR rational map	7
4. A criterion to be a JNR spectral curve	8
4.1. (N, N) spectral curves	8
4.2. Grids	9
4.3. JNR monopoles and grids	9
5. Holomorphic spheres	10

2010 *Mathematics Subject Classification.* 81T13, 53C07, 14D21.

MKM acknowledges support under the Australian Research Council's *Discovery Projects* funding scheme (project number DP180100383). PN acknowledges support under the Australian Research Council's *Discovery Projects* funding scheme (project numbers DP170102028 and DP180103891).

5.1. $SO(3)$ action on JNR monopoles	12
6. Discrete Nahm equations	13
7. Projection to rational maps	15
8. The energy density at infinity	16
References	19

1. INTRODUCTION

The first author was introduced to monopoles by Sir Michael Atiyah in 1980 when Atiyah proposed that generalising Nigel Hitchin's then recent article [14] from $SU(2)$ to an arbitrary compact Lie group would make a good DPhil project. After some years of expert supervision by Atiyah this proved to be the case.

In 1989 Atiyah came to the Australian National University to attend a Miniconference on Geometry and Physics [3]. He talked to Rodney Baxter about his work with Perk and Au-Yang on the chiral Potts model. His interest was particularly in the fact that their new solutions of the chiral Potts model [9] involved algebraic curves of genus greater than 1. This led to him making what he later told the second author was a 'flight of fancy' that this class of curves might be related to the spectral curves of monopoles, another interesting class of curves and that the chiral Potts integrable model solutions might also be related to monopoles and instantons and the whole question of self-duality and integrability [17].

In 1990 the first author visited Atiyah at Oxford and we resolved the first part of this conjecture identifying the Baxter, Perk, Au-Yang curves with particular spectral curves of zero mass hyperbolic monopoles [4, 7]. Hyperbolic monopoles had been introduced earlier by Atiyah in [2]. However finding a relationship between monopoles and the actual solutions of the chiral Potts model has remained elusive. We discussed it a number of times over the years in various locations including the Master's Lodge at Trinity College and at various times in Edinburgh. Part of the difficulty would seem to be understanding the significance of the mass zero hyperbolic monopoles which are probably better thought of as limits of positive mass hyperbolic monopoles¹.

Trying to resolve Atiyah's conjecture has motivated a number of interesting explorations of hyperbolic monopoles by the current two authors and their colleagues [20, 19, 22, 21]. While it would be particularly nice for this volume to say we were about to present the resolution of this conjecture that is sadly not the case. We present instead a discussion of some interesting results involving mass $\frac{1}{2}$ hyperbolic monopoles. It seemed appropriate however in the current circumstances to explain the deeper underlying motivation for our interest in hyperbolic monopoles.

¹See for example [19] for further discussion of this point.

Monopoles are solutions of the Bogomolny equations which arise as the dimensional reduction of time-invariant instantons on \mathbb{R}^4 . In [2] Atiyah noticed that by exploiting the conformal invariance of the self-duality equations one can pass from instantons on $S^4 - S^2$ to instantons on $\mathbb{H}^3 \times S^1$ with metric the product of the hyperbolic metric and the circle metric and then by demanding rotation invariance in the S^1 direction to solutions of the Bogomolny equations on hyperbolic space \mathbb{H}^3 . As in the Euclidean case appropriate boundary conditions can be imposed resulting in the definition of the mass and charge of the monopole. One finds that the mass is in $\frac{1}{2}\mathbb{Z}$ if and only if under the conformal transformation above the instanton on $S^4 - S^2$ extends to an instanton on S^4 . In [16] Manton and Sutcliffe showed how to construct particular mass $\frac{1}{2}$ monopoles known as JNR monopoles in this way using an ansatz for instantons due to Jackiw, Nohl and Rebbi [15], the formula for which also occurred in [12]. These instantons were early examples of what later became known as the Atiyah, Drinfeld, Hitchin, Manin (ADHM) construction [5]. In [16] Manton and Sutcliffe construct JNR monopoles with various symmetries including: spherical, axial, tetrahedral, octahedral and icosahedral. Following on this work [10, 11] give formulae for the spectral curve and the rational map of the general JNR monopole and for these particular symmetric cases. What is quite remarkable about JNR monopoles is how simple and explicit the formulae are for the spectral curves and rational maps, not just in the symmetric cases but also for the general JNR monopole.

Using these results in Section 4 we make a number of additions to this theory. We give a general geometric constraint that characterises the spectral curves of JNR monopoles. This is the existence of a so-called grid. We show that this can be used to define the non-vanishing section of the line bundle L^{2p+N} the existence of which is one of the important properties of a spectral curve of a mass p charge N monopole. In turn we show that this section can be used as scattering data to define the rational map as in the Euclidean case.

We represent the monopole in terms of holomorphic, or twistor, data rather than writing down the explicit fields, for most of the paper. The energy density of a monopole is the L^2 norm of its curvature which gives a type of location of the fields of the monopole. Moreover, a hyperbolic monopole is uniquely determined by the limit at infinity of its energy density. In Section 8 we determine the energy density of the JNR monopole at infinity. To achieve this we use the holomorphic sphere of the monopole, described in Section 5, which was constructed for hyperbolic monopoles of any mass in work of the current authors with Singer [19]. This is a holomorphic map $q: \mathbb{P}_1 \rightarrow \mathbb{P}_N$, between complex projective spaces, determined up to the action of $U(N+1)$. One satisfying aspect of this paper is a simple formula for the holomorphic sphere in terms of the JNR data given by Proposition 5.2, which leads to a formula for the energy density of the monopole at infinity. In Section 6 we relate JNR data to the discrete Nahm equations. In Section 7 we discuss the relationship between the holomorphic sphere and the different rational maps obtained by scattering from different points at infinity.

Our thanks to an anonymous referee for helpful remarks.

2. HYPERBOLIC MONOPOLES

A monopole in Euclidean, respectively hyperbolic, space is a pair (A, Φ) consisting of a connection A with L^2 curvature F_A defined on a trivial bundle E over \mathbb{R}^3 , respectively \mathbb{H}^3 , with structure group $SU(2)$, and a Higgs field $\Phi : \mathbb{R}^3 (\mathbb{H}^3) \rightarrow \mathfrak{su}(2)$ that solves the Bogomolny equation

$$(2.1) \quad d_A \Phi = *F_A$$

and satisfies a boundary condition which we do not need to consider in detail but which has two important consequences:

- There is a limit $\lim_{r \rightarrow \infty} \|\Phi\| = p$, known as the *mass* of the monopole.
- There is a well-defined map on the two-sphere at infinity $\Phi_\infty : S^2 \rightarrow S^2$ whose topological degree is called the *charge* of the monopole.

The Euclidean and hyperbolic metrics appear in the Bogomolny equation (2.1) via the Hodge star operator $*$. Many features of Euclidean and hyperbolic monopoles behave in a similar fashion. However the mass is quite different. In the Euclidean case we can transform any monopole of a given non-zero mass to a monopole of a different non-zero mass by a simple rescaling of space and the fields. In the hyperbolic case we can perform the same procedure but after rescaling hyperbolic space by dilation the curvature of the metric changes. So if we fix a standard copy of \mathbb{H}^3 of scalar curvature -1 the monopoles of different masses must be considered as different.

We will be concerned primarily with the twistor picture of hyperbolic monopoles for which we need the mini-twistor space [14] of all oriented geodesics of \mathbb{H}^3 . Recall that a geodesic in \mathbb{H}^3 is determined by the two points where it meets the sphere at infinity. Thus the mini-twistor space is the complex manifold

$$Q = \mathbb{P}_1 \times \mathbb{P}_1 - \bar{\Delta}$$

where the point $(\eta, \zeta) \in \mathbb{P}_1 \times \mathbb{P}_1$ represents the geodesic that runs from $\hat{\eta} = -1/\bar{\eta}$, the antipodal point of η , to ζ considered as points on the sphere at infinity. The antidiagonal $\bar{\Delta}$ has been removed as it represents a geodesic from z to itself.

The monopole (A, Φ) (up to gauge transformation) is determined by an algebraic curve $S \subset Q$ called the *spectral curve* of the monopole. See [14] for the Euclidean case and Atiyah's foundational work [2] for the hyperbolic case of half-integral mass. The spectral curve is the collection of all oriented geodesics γ along which the equation

$$(\nabla_\gamma - i\Phi)s = 0$$

has L^2 solutions. The spectral curve of a hyperbolic monopole of charge N and mass p satisfies a number of conditions:

- It is compact of bi-degree (N, N) . Hence determined by a holomorphic section of the bundle $\mathcal{O}(N, N)$.
- It is *real*, that is fixed by the anti-holomorphic involution τ which reverses the orientation of the geodesic.

- There is a non-vanishing holomorphic section of L^{2p+N} over S where $L = \mathcal{O}(1, -1)$.

The moduli space of charge N monopoles (A, Φ) modulo gauge transformations is known to be a manifold of dimension $4N - 1$ independent of the mass of the monopole. The structure of this moduli space was proven by Atiyah [1] to be diffeomorphic to the space of based rational maps from \mathbb{P}_1 to \mathbb{P}_1 . Atiyah proved this in [1] as the S^1 invariant case of the more general theorem that instantons with structure group G on S^4 are in one-to-one correspondence with based holomorphic maps from \mathbb{P}_1 to the loop group ΩG . Circle invariant instantons, equivalently half-integer mass hyperbolic monopoles, correspond to maps from \mathbb{P}_1 to conjugacy classes of homomorphisms from S^1 to the G inside ΩG which are homogeneous spaces G/P of G . For $SU(2)$ instantons, the homogeneous space is $SU(2)/U(1) = \mathbb{P}_1$, and the moduli space of charge N , mass $p \in \frac{1}{2}\mathbb{Z}$ hyperbolic monopoles modulo gauge transformations that are the identity at ∞ is homeomorphic to degree N based maps from \mathbb{P}_1 to \mathbb{P}_1 . This space has dimension $4N$, and when we allow all gauge transformations, the rational map is well-defined up to rescaling by a factor in $U(1)$ and the dimension is $4N - 1$. In both the Euclidean and hyperbolic cases it is difficult to write down monopoles (A, Φ) or their spectral data or rational map except in highly symmetric cases.

3. JNR MONOPOLES

Definition 3.1. *JNR data* consists of a collection of $N + 1$ positive numbers $\lambda_i > 0$ called *weights* and $N + 1$ distinct complex numbers γ_i called *poles*.

As explained in [16] each collection of JNR data determines a hyperbolic monopole of mass $\frac{1}{2}$ and charge N . This can be seen from the degree (N, N) spectral curve given in Section 3.1 which admits a trivialisation of the line bundle $\mathcal{O}(N + 2p, -N - 2p)$ for $p = \frac{1}{2}$ proving that it corresponds to a charge N hyperbolic monopole of mass $\frac{1}{2}$. It can also be seen by writing down the fields directly as in [16], which is related to the ADHM constructions of instantons. The ADHM construction is discussed in Section 6 in terms of the discrete Nahm equations.

If we rescale the weights in Definition 3.1 we obtain a gauge equivalent monopole. We might expect the JNR monopoles to be a submanifold of dimension $3N + 2$ inside the $4N - 1$ dimensional moduli space of all monopoles of charge N . As discussed in [10, Section 2] this is true except for some low N with the result that for $N = 1, 2, 3$ all monopoles are obtained as JNR monopoles. There seems to be no clear geometric picture of which monopoles are JNR monopoles for $N > 3$.

3.1. JNR spectral curve. It is shown in [10] that the spectral curve of a JNR monopole is given in terms of its JNR data $\{\lambda_i, \gamma_i \mid i = 0, \dots, N\}$ by the equation

$$(3.1) \quad 0 = p(\eta, \zeta) = \sum_{i=0}^N \lambda_i^2 \prod_{\substack{j=0 \\ j \neq i}}^N (\zeta - \gamma_j)(1 + \eta \bar{\gamma}_j).$$

Notice that, as expected from the remarks above, rescaling the weights leaves the spectral curve unchanged.

It is simple to check the basic conditions satisfied by a JNR spectral curve. First notice that

$$(3.2) \quad p(\eta, \zeta) = (-1)^N \eta^N \zeta^N \overline{p\left(\frac{-1}{\bar{\zeta}}, \frac{-1}{\bar{\eta}}\right)}$$

which shows that the spectral curve is real. Moreover

$$\begin{aligned} p\left(\frac{-1}{\bar{\zeta}}, \zeta\right) &= \sum_{i=0}^N \lambda_i^2 \prod_{\substack{j=0 \\ j \neq i}}^N (\zeta - \gamma_j) \left(1 + \frac{-1}{\bar{\zeta}} \bar{\gamma}_j\right) \\ &= \frac{1}{\bar{\zeta}^N} \sum_{i=0}^N \lambda_i^2 \prod_{\substack{j=0 \\ j \neq i}}^N |\zeta - \gamma_j|^2 \\ &\neq 0 \end{aligned}$$

for any ζ so the spectral curve does not intersect the anti-diagonal. Hence it is a compact subset of Q .

If S is a spectral curve of a mass $p = \frac{1}{2}$ monopole it must have a non-vanishing section s of

$$L^{2p+N} = L^{1+N} = \mathcal{O}(N+1, -N-1).$$

We construct this by restricting the global meromorphic section

$$(3.3) \quad s(\eta, \zeta) = \frac{\prod_{i=0}^N (\zeta - \gamma_i)}{\prod_{i=0}^N (1 + \eta \bar{\gamma}_i)}$$

of $\mathcal{O}(N+1, -N-1)$.

Proposition 3.2. *The section s defined in equation (3.3) is non-vanishing on the spectral curve arising from the JNR data.*

Proof. The only points (η, ζ) at which this might not be true are those points where either of $\zeta = \gamma_i$ or $\eta = \frac{-1}{\bar{\gamma}_j}$ holds. The only points on the spectral curve like that are $(\eta, \zeta) = (\frac{-1}{\bar{\gamma}_j}, \gamma_i)$ with $i \neq j$. Fix such an i and j . Then on a small enough open set around this point we have

$$s(\eta, \zeta) = \sigma(\eta, \zeta) \frac{(\zeta - \gamma_i)}{(1 + \eta \bar{\gamma}_j)}$$

with σ non-vanishing. Likewise we have

$$p(\eta, \zeta) = \alpha(\eta, \zeta)(\zeta - \gamma_i) + \beta(\eta, \zeta)(1 + \eta \bar{\gamma}_j) + \gamma(\eta, \zeta)(\zeta - \gamma_i)(1 + \eta \bar{\gamma}_j)$$

with α, β, γ non-vanishing. Hence

$$s(\zeta, \eta) = \frac{-\sigma}{\alpha} (\beta + \gamma(\zeta - \gamma_i)) + p \left(\frac{1}{\alpha(1 + \eta \bar{\gamma}_j)} \right)$$

and

$$\frac{-\sigma}{\alpha} (\beta + \gamma(\zeta - \gamma_i))$$

is non-vanishing in a neighbourhood of $(\frac{-1}{\gamma_j}, \gamma_i)$. \square

3.2. Scattering. Each point in \mathbb{H}^3 defines a two-sphere of oriented geodesics through it. As that point approaches a point γ at infinity this two-sphere of oriented geodesics approaches a pair of generators in Q corresponding to the geodesics beginning at $-1/\bar{\eta} = \gamma$ or ending at $\zeta = \gamma$ or the solutions to

$$0 = (\zeta - \gamma)(1 + \eta\bar{\gamma}).$$

Notice that we can normalise $\sum_{i=0}^N \lambda_i^2 = 1$. Then if we let one value say $\lambda_k \rightarrow 1$ and all the others go to zero the spectral curve (3.1) approaches the solutions of

$$0 = \prod_{\substack{j=0 \\ j \neq k}}^N (\zeta - \gamma_j)(1 + \eta\bar{\gamma}_j).$$

that correspond to the set of geodesics through the points $\gamma_0, \dots, \gamma_{k-1}, \gamma_{k+1}, \gamma_N$ at infinity in hyperbolic space. We can think of the monopole as approaching a monopole located at $N - 1$ points near the $\gamma_0, \dots, \gamma_{k-1}, \gamma_{k+1}, \gamma_N$. We comment again about this in Section 8.

3.3. JNR rational map. A remarkable result about both Euclidean and hyperbolic monopoles is that the *framed* moduli spaces of charge N monopoles are diffeomorphic to the space of all based rational (holomorphic) maps $\mathcal{R}: \mathbb{P}_1 \rightarrow \mathbb{P}_1$. Here based means that $\mathcal{R}(\infty) = 0$. The framed moduli space uses gauge transformations that are the identity at a chosen point on the sphere S_∞^2 of directions at infinity in \mathbb{R}^2 or \mathbb{H}^3 . The usual moduli space is a quotient of the framed moduli space by S^1 . In Cockburn's PhD thesis [11] (see also [10]) the rational map of a JNR monopole is calculated explicitly in terms of JNR data as

$$(3.4) \quad \mathcal{R}(z) = \frac{\sum_{i=0}^N \sum_{j=i+1}^N \lambda_i^2 \lambda_j^2 (\gamma_i - \gamma_j)^2 \prod_{\substack{k=0 \\ k \neq i, j}}^N (z - \gamma_k)}{\left(\sum_{i=0}^N \lambda_i^2 \right) \left(\sum_{j=0}^N \lambda_j^2 \prod_{\substack{k=0 \\ k \neq j}}^N (z - \gamma_k) \right)}$$

Again we note that this is unchanged if we rescale the weights.

We show that (3.4) can also be calculated by using the scattering construction of Atiyah [2, page 30] (given in more detail in the Euclidean case in [6, page 127]).

Proposition 3.3. *The rational map (3.4) for JNR monopoles agrees with the rational map Atiyah defines for hyperbolic monopoles via scattering.*

Proof. To simplify the calculation let $\sum_{i=0}^N \lambda_i^2 = 1$. Then following Atiyah, the denominator of the rational map is

$$Q(z) = p(0, z) = \sum_{i=0}^N \lambda_i^2 \prod_{\substack{j=0 \\ j \neq i}}^N (z - \gamma_j).$$

The numerator $P(z)$ is the polynomial of degree $N - 1$ satisfying

$$(3.5) \quad P(z) = s(0, z) + (az + b)Q(z)$$

where s is the section of L^{N+1} defined in (3.3). The scattering rational map is $P(z)/Q(z)$. We want to show that this agrees with $\mathcal{R}(z)$. Notice that $Q(z)$ is the denominator of (3.4).

First we calculate that

$$P(z) = \prod_{i=0}^N (z - \gamma_i)^N + (az + b) \sum_{i=0}^N \lambda_i^2 \prod_{\substack{j=0 \\ j \neq i}}^N (z - \gamma_j)$$

and choosing a and b to remove the terms of order z^{N+1} and z^N we find that $a = -1$ and $b = \sum_{i=0}^N \lambda_i^2 \gamma_i$. Substituting this back in we obtain

$$\begin{aligned} P(z) &= \prod_{i=0}^N (z - \gamma_i) + (-z + \sum_{k=0}^N \lambda_k^2 \gamma_k) \sum_{i=0}^N \lambda_i^2 \prod_{\substack{j=0 \\ j \neq i}}^N (z - \gamma_j) \\ &= \sum_{i,k=0}^N \lambda_i^2 \lambda_k^2 \left(\prod_{j=0}^N (z - \gamma_j) - (z - \lambda_k) \prod_{\substack{j=0 \\ j \neq i}}^N (z - \gamma_j) \right) \\ &= \sum_{i,k=0}^N \lambda_i^2 \lambda_k^2 \left(1 - \frac{(z - \lambda_k)}{(z - \gamma_i)} \right) \prod_{j=0}^N (z - \gamma_j) \\ &= \sum_{i,k=0}^N \lambda_i^2 \lambda_k^2 \left(\frac{(\lambda_k - \lambda_i)}{(z - \gamma_i)} \right) \prod_{j=0}^N (z - \gamma_j) \\ &= \sum_{i < k=0}^N \lambda_i^2 \lambda_k^2 \left(\frac{(\lambda_k - \lambda_i)}{(z - \gamma_i)} + \frac{(\lambda_i - \lambda_k)}{(z - \gamma_k)} \right) \prod_{j=0}^N (z - \gamma_j) \\ &= \sum_{i < k=0}^N \lambda_i^2 \lambda_k^2 \left(\frac{(\lambda_k - \lambda_i)^2}{(z - \gamma_i)(z - \gamma_k)} \right) \prod_{j=0}^N (z - \gamma_j) \end{aligned}$$

Hence $P(z)$ is the numerator of (3.4) and thus $\mathcal{R}(z) = P(z)/Q(z)$. \square

4. A CRITERION TO BE A JNR SPECTRAL CURVE

In this section we give a criterion for when a spectral curve must be the JNR curve of some JNR data.

4.1. (N, N) spectral curves. We need some normalisations for (N, N) spectral curves. Let S be a real (N, N) curve not intersecting the anti-diagonal defined by

$p(\eta, \zeta) = 0$. Then there is a $\lambda \neq 0$ such that

$$p(\eta, \zeta) = \lambda \eta^N \zeta^N \overline{p(\frac{-1}{\bar{\zeta}}, \frac{-1}{\bar{\eta}})}$$

Moreover if we replace $p(\eta, \zeta)$ by $\mu p(\eta, \zeta)$ for some complex number μ then λ is replaced by $\lambda \frac{\mu}{\bar{\mu}}$. We choose μ so that

$$p(\eta, \zeta) = (-1)^N \eta^N \zeta^N \overline{p(\frac{-1}{\bar{\zeta}}, \frac{-1}{\bar{\eta}})}.$$

It follows from this normalisation and the fact that S does not intersect the anti-diagonal that

$$p(\frac{-1}{\bar{\zeta}}, \zeta) \bar{\zeta}^N$$

is either negative or positive and we can, without loss of generality assume that it is positive. So in summary we normalise our spectral curve defining polynomials so that

$$(4.1) \quad p(\eta, \zeta) = (-1)^N \eta^N \zeta^N \overline{p(\frac{-1}{\bar{\zeta}}, \frac{-1}{\bar{\eta}})} \quad \text{and} \quad p(\frac{-1}{\bar{\zeta}}, \zeta) \bar{\zeta}^N > 0.$$

It is straightforward to check that the JNR polynomial defined in equation (3.1) is normalised this way.

4.2. Grids.

Definition 4.1. Let us define a *grid* in $Q \subset \mathbb{P}_1 \times \mathbb{P}_1$ to be a subset of the form

$$\mathcal{G} = \{(\frac{-1}{\bar{\gamma}_i}, \gamma_j) \mid 0 \leq i \neq j \leq N\}$$

for γ_i a collection of $N + 1$ distinct points in \mathbb{P}_1 .

Notice that a grid is a collection of $N(N + 1)$ points which is real and does not intersect the anti-diagonal. Sometimes we need to include the points we have excluded to avoid the anti-diagonal so we define $\bar{\mathcal{G}}$ to be

$$\bar{\mathcal{G}} = \{(\frac{-1}{\bar{\gamma}_i}, \gamma_j) \mid 0 \leq i, j \leq N\}$$

which is $(N + 1)^2$ points. We say that a curve S in $\mathbb{P}_1 \times \mathbb{P}_1$ *admits a grid* if there is a grid \mathcal{G} with $\mathcal{G} \subset S$.

4.3. JNR monopoles and grids. Clearly a JNR spectral curve admits the grid determined by the poles in the JNR data. In fact the converse is true as shown by the following proposition.

Proposition 4.2. *Let S be a real curve of degree (N, N) not intersecting the anti-diagonal and admitting a grid \mathcal{G} determined by points $\gamma_0, \dots, \gamma_N$. Then there exist positive real numbers $\lambda_0, \dots, \lambda_N$ such that S is the spectral curve determined by the JNR data with poles $\gamma_0, \dots, \gamma_N$ and weights $\lambda_0, \dots, \lambda_N$.*

Proof. Let \tilde{p} define S and normalise it as above. Define

$$\lambda_i^2 = \bar{\gamma}_i^N \frac{\tilde{p}(\frac{-1}{\bar{\gamma}_i}, \gamma_i)}{\prod_{\substack{j=0 \\ j \neq i}}^N |\gamma_i - \gamma_j|^2}$$

and use it to define p by equation (3.1). It follows that \tilde{p} and p agree on $\bar{\mathcal{G}}$. So from the exact sequence in sheaf cohomology

$$0 \rightarrow \mathcal{O}_Q(-1, -1) \rightarrow \mathcal{O}_Q(N, N) \rightarrow \mathcal{O}_{\bar{\mathcal{G}}}(N, N) \rightarrow 0$$

we deduce that $p = \tilde{p}$. \square

Hence we have:

Corollary 4.3. *A spectral curve comes from JNR data if and only if it admits a grid.*

Consider the formal dimension count for grids and JNR data. Notice that the vector space $H^0(Q, \mathcal{O}(N, N))$ has dimension $(N+1)^2$ so the space of curves has dimension $(N+1)^2 - 1 = N^2 + 2N$ real dimensions. The closed grid has $(N+1)^2 = N^2 + 2N + 1$ points. If we remove the intersection with the anti-diagonal that leaves the grid which has $N^2 + N$ points in real pairs. If we require the real curve to go through the grid that is $\frac{1}{2}(N^2 + N)$ constraints each contributing two real dimensions. So we might reasonably expect the space of spectral curves admitting grids to be determined by $N^2 + 2N - (N^2 + N) = N$ parameters. This fits with $N+1$ weights less the overall scaling factor of 1.

5. HOLOMORPHIC SPHERES

In [19] with Singer we constructed a new invariant, particular to hyperbolic monopoles called the holomorphic sphere. A hyperbolic monopole has a well-defined limit at infinity given by a reducible connection A_∞ over a two-sphere. Denote by F_{A_∞} the curvature of the reducible connection. We showed that:

Theorem 5.1. [19, Theorem 1] *An $SU(2)$ hyperbolic monopole (A, Φ) of charge N is determined by a degree N holomorphic embedding $q : \mathbb{P}_1 \rightarrow \mathbb{P}_N$ with the properties:*

- (i) $\Sigma = \{(\eta, \zeta) \in \mathbb{P}_1 \times \mathbb{P}_1 \mid \langle q(\hat{\eta}), q(\hat{\zeta}) \rangle = 0\}$ is the spectral curve of (A, Φ) ;
- (ii) q is uniquely defined up to the action of $U(N+1)$ on its image;
- (iii) $F_{A_\infty} = q^*\omega$, for ω the Kahler form on \mathbb{P}_N .

Here we have changed the notation using (η, ζ) for (w, z) in [19] and $\hat{\eta} = \frac{-1}{\bar{\eta}}$. Also $\langle \cdot, \cdot \rangle$ is the Hermitian inner product, conjugate in the first factor.

Property (iii) of Theorem 5.1 can be used to prove that F_{A_∞} uniquely determines the monopole (A, Φ) up to gauge transformation. It potentially gives one a way to study monopoles from a two-dimensional perspective much like the AdS-CFT correspondence which proposes a relationship between string theory on anti-de Sitter spacetime and conformal field theory on the boundary. In general a degree N holomorphic embedding $q : \mathbb{P}_1 \rightarrow \mathbb{P}_N$ does not correspond to a hyperbolic monopole. The first challenge then, to implement such ideas, is to understand the space of holomorphic spheres, or fields at infinity F_{A_∞} . This is achieved for JNR monopoles in the proposition below.

Proposition 5.2. *The holomorphic sphere for a JNR monopole is given by*

$$(5.1) \quad q(z) = \left[\frac{\lambda_0}{z - \gamma_0}, \frac{\lambda_1}{z - \gamma_1}, \dots, \frac{\lambda_N}{z - \gamma_N} \right].$$

Proof. We need to connect with the construction of the holomorphic sphere in [19]. Comparing the normalisations in [19, Lemma 2.1] with those in equation (4.1) we see that ψ in [19] is related to p by

$$\psi(\eta, \zeta) = \frac{(-1)^N}{\eta^N} p(\eta, \zeta).$$

Then we need to use [19, Theorem 4] which shows that q is determined by ψ and satisfies

$$(5.2) \quad \psi(\eta, \zeta) = \langle q(\frac{-1}{\bar{\eta}}), q(\zeta) \rangle.$$

In fact from the discussion there it follows that if $q : \mathbb{P}_1 \rightarrow \mathbb{P}_N$ is of degree N and full, that is does not have its image in a proper subspace, then equation (5.2) determines q up to the action of $U(N+1)$. So we have

$$\begin{aligned} \psi(\eta, \zeta) &= \left(\frac{-1}{\eta}\right)^N \sum_{i=0}^N \lambda_i^2 \prod_{\substack{j=0 \\ j \neq i}}^N (\zeta - \gamma_j)(1 + \eta \bar{\gamma}_j) \\ &= \sum_{i=0}^N \lambda_i^2 \prod_{\substack{j=0 \\ j \neq i}}^N (\zeta - \gamma_j) \left(\frac{-1}{\eta} - \bar{\gamma}_j\right) \\ &= \sum_{i=0}^N \left(\frac{\lambda_i}{\frac{-1}{\eta} - \bar{\gamma}_i} \prod_{\substack{j=0 \\ j \neq i}}^N \left(\frac{-1}{\eta} - \bar{\gamma}_j\right) \right) \left(\frac{\lambda_i}{\zeta - \gamma_i} \prod_{j=0}^N (\zeta - \gamma_j) \right) \\ &= \sum_{i=0}^N \left(\frac{\lambda_i}{\frac{-1}{\bar{\eta}} - \gamma_i} \prod_{\substack{j=0 \\ j \neq i}}^N \left(\frac{-1}{\bar{\eta}} - \gamma_j\right) \right) \left(\frac{\lambda_i}{\zeta - \gamma_i} \prod_{j=0}^N (\zeta - \gamma_j) \right) \\ &= \langle q(\frac{-1}{\bar{\eta}}), q(\zeta) \rangle \end{aligned}$$

so that

$$(5.3) \quad q(z) = \left[\frac{\lambda_0}{z - \gamma_0} \left(\prod_{j=0}^N (z - \gamma_j) \right), \dots, \frac{\lambda_N}{z - \gamma_N} \left(\prod_{j=0}^N (z - \gamma_j) \right) \right] \\ = \left[\frac{\lambda_0}{z - \gamma_0}, \dots, \frac{\lambda_N}{z - \gamma_N} \right].$$

has degree N and satisfies (5.2). If q is not full then there is some $(\alpha_0, \dots, \alpha_N)$ such that

$$\sum_{k=0}^N \frac{\alpha_k \lambda_k}{(z - \gamma_k)} = 0$$

which is not possible as the γ_k are distinct and the $\lambda_k > 0$. Hence the result follows. \square

We shall see in Section 8 that along with $F_{A_\infty} = q^* \omega$ this can be used to give a formula for the energy density of a JNR monopole at infinity.

5.1. $SO(3)$ action on JNR monopoles. A consequence of Corollary 4.3 and Proposition 5.2 is that the action by isometries of $SO(3)$ on \mathbb{H}^3 preserves JNR monopoles. To see this, note that the spectral curve is rotated by the ambient $SO(3)$ action on $\mathbb{P}_1 \times \mathbb{P}_1$. Hence the action of $SO(3)$ on the holomorphic sphere is given by $q(z) \mapsto q(g^{-1} \cdot z)$. (Note that this only works for $SO(3)$ and fails for $PSL(2, \mathbb{C})$.) Here $g^{-1} \cdot z = \frac{\bar{a}z - b}{bz + a}$ for $g \in SU(2)/\pm I \cong SO(3)$. Hence the i th term of q becomes

$$\frac{\lambda_i}{z - \gamma_i} \mapsto \frac{\lambda_i}{\frac{\bar{a}z - b}{bz + a} - \gamma_i} = \frac{\lambda_i(\bar{b}z + a)}{(\bar{a} - \bar{b}\gamma_i)z - (a\gamma_i + b)} = \frac{\frac{\lambda_i}{\bar{a} - b\gamma_i}(\bar{b}z + a)}{z - \frac{a\gamma_i + b}{\bar{a} - b\gamma_i}}.$$

After dividing by the common factor of $\bar{b}z + a$ and replacing $\bar{a} - \bar{b}\gamma_i$ by $|\bar{a} - \bar{b}\gamma_i|$ via a diagonal $U(N+1)$ action on \mathbb{P}_N we get

$$q(z) = \left[\frac{\lambda_0}{z - \gamma_0}, \frac{\lambda_1}{z - \gamma_1}, \dots, \frac{\lambda_N}{z - \gamma_N} \right] \mapsto \left[\frac{\frac{\lambda_0}{|\bar{a} - b\gamma_0|}}{z - \frac{a\gamma_0 + b}{\bar{a} - b\gamma_0}}, \frac{\frac{\lambda_1}{|\bar{a} - b\gamma_1|}}{z - \frac{a\gamma_1 + b}{\bar{a} - b\gamma_1}}, \dots, \frac{\frac{\lambda_N}{|\bar{a} - b\gamma_N|}}{z - \frac{a\gamma_N + b}{\bar{a} - b\gamma_N}} \right].$$

Since the holomorphic sphere uniquely determines the monopole, we conclude that the $SO(3)$ action on JNR data is given by

$$\begin{pmatrix} a & b \\ -\bar{b} & \bar{a} \end{pmatrix} \cdot \{\lambda_j, \gamma_j\} = \left\{ \frac{\lambda_j}{|\bar{a} - \bar{b}\gamma_j|}, \frac{a\gamma_j + b}{\bar{a} - \bar{b}\gamma_j} \right\}.$$

In particular the $\frac{1}{2}(N+1)(N+2)$ functions

$$\frac{\lambda_i^2 \lambda_j^2}{|1 + \bar{\gamma}_i \gamma_j|^2}$$

are invariant functions. The orbit of $SO(3)$ is generically 3-dimensional for $N > 1$ hence there are at least $\frac{1}{2}(N+1)(N+2) + 3 - (3N+3) = \frac{1}{2}(N-1)(N-2)$ relations among the invariant functions

6. DISCRETE NAHM EQUATIONS

A hyperbolic monopole of charge N and mass p corresponds to an S^1 -invariant instanton on S^4 of Chern class $c_2 = 2pN$, proven by Atiyah in [2]. Austin and Braam proved in [8] that to an $SU(2)$ half integral mass charge N hyperbolic monopole one can associate a solution of the discrete Nahm equations which is essentially a decomposition into S^1 weight spaces of the ADHM data for the instanton on S^4 . This gives a discrete analogue of the Nahm (ordinary differential) equations used to produce Euclidean monopoles. The discrete Nahm data is given in the case $p \in \mathbb{Z} + \frac{1}{2}$ by:

$$\mathcal{M} = \{(\beta_{2i}, \gamma_{2i-1}, v) \in \mathfrak{gl}(N, \mathbb{C}) \times \mathfrak{gl}(N, \mathbb{C}) \times \mathbb{C}^N \mid i \in [-p + \frac{1}{2}, p - \frac{1}{2}] \cap \mathbb{Z} \text{ satisfying (6.1), (6.2), (6.3), (6.4), (6.5)}\} / \mathcal{G}$$

$$\begin{aligned} (6.1) \quad \gamma_j &= \gamma_{-j}^T & -2p+2 \leq j \leq 2p-2, \quad j \text{ odd} \\ (6.2) \quad \beta_j &= \beta_{-j}^T & -2p+1 \leq j \leq 2p-1, \quad j \text{ even} \\ (6.3) \quad \beta_{j-1}\gamma_j - \gamma_j\beta_{j+1} &= 0 & -2p+2 \leq j \leq 2p-2, \quad j \text{ odd} \\ (6.4) \quad [\bar{\beta}_j^T, \beta_j] + \bar{\gamma}_{j-1}^T\gamma_{j-1} - \gamma_{j+1}\bar{\gamma}_{j+1}^T &= 0 & -2p+3 \leq j \leq 2p-3, \quad j \text{ even} \\ (6.5) \quad [\bar{\beta}_{2p-1}^T, \beta_{2p-1}] + v\bar{v}^T - \bar{\gamma}_{2p-2}^T\gamma_{2p-2} &= 0 \end{aligned}$$

which admits an action of a gauge group

$$\mathcal{G} = \{g_j \in U(N) \mid j = -2p+1, -2p+3, \dots, 0, \dots, 2p-3, 2p-1, \quad g_j = \bar{g}_{-j}\}$$

by

$$\begin{aligned} \beta_j &\mapsto g_j \beta_j g_j^{-1} \\ \gamma_j &\mapsto g_{j-1} \gamma_j g_{j+1}^{-1} \\ v &\mapsto g_{-2p+1} v. \end{aligned}$$

These are known as the discrete Nahm equations. One can equivalently take the following description with complex gauge group

$$\mathcal{M} = \{(\beta_{2i}, \gamma_{2i-1}, v) \in \mathfrak{gl}(N, \mathbb{C}) \times \mathfrak{gl}(N, \mathbb{C}) \times \mathbb{C}^N \mid i \in [0, p - \frac{1}{2}] \cap \mathbb{Z}, \beta_0 = \beta_0^T \text{ satisfying (6.3) + stability}\} / \mathcal{G}^{\mathbb{C}}$$

where

$$\mathcal{G}^{\mathbb{C}} = \{g_j \in GL(N, \mathbb{C}) \mid j = 0, \dots, 2p-3, 2p-1, \quad g_0 \in O(N, \mathbb{C})\}$$

and the stability condition is determined explicitly in [8] and given here in (6.8) for the $p = \frac{1}{2}$ case. The two descriptions of \mathcal{M} are isomorphic since equations (6.4) and (6.5) can be interpreted as the moment map of \mathcal{G} .

The pair (β_{-2p+1}, v) determines the full solution of the discrete Nahm equations. It was shown in [8] that the map

$$(6.6) \quad \begin{pmatrix} \beta_{-2p+1} - z \\ v^T \end{pmatrix} : \mathbb{C}^N \rightarrow \mathbb{C}^{N+1}$$

is a monad on S^2 which determines the boundary value of the hyperbolic monopole. The monad can be interpreted as a degree N holomorphic map $q : \mathbb{P}_1 \rightarrow \mathbb{P}_N$ given explicitly by

$$(6.7) \quad q(z) = \begin{pmatrix} \det(\beta_{-2p+1} - z) \\ -\det(\beta_{-2p+1} - z) \cdot (\beta_{-2p+1}^T - z)^{-1}v \end{pmatrix}.$$

The map is well-defined up to the $U(N)$ action on the first N coordinates, since β_{-2p+1} admits a $U(N)$ action. It gives the holomorphic sphere defined in Section 5.

When $p = \frac{1}{2}$, the complex discrete Nahm data becomes

$$\mathcal{M}_N = \{(\beta, v) \in \mathfrak{gl}(N, \mathbb{C}) \times \mathbb{C}^N \mid \beta = \beta^T \text{ satisfying (6.8)}\} / O(N, \mathbb{C})$$

where $\beta = \beta_0$, the discrete Nahm equations are trivial and the stability condition is

$$(6.8) \quad \{\beta^j v \mid j = 0, \dots, N-1\} \text{ span } \mathbb{C}^N.$$

Note that the dimension is given by

$$\dim_{\mathbb{C}} \mathcal{M}_N = \frac{1}{2}N(N+1) + N - \frac{1}{2}N(N-1) = 2N$$

as expected.

JNR monopoles of charge N give a submanifold of \mathcal{M}_N . The map from JNR data $\{\lambda_i, \gamma_i \mid i = 0, \dots, N\}$ to the discrete Nahm data \mathcal{M}_N can be deduced from the map from JNR data to ADHM data in Cockburn's PhD thesis [11]. It is rather complicated but we will show here that the map from JNR data to discrete Nahm data is rather simple on a limit of JNR data.

Suppose that β is orthogonally diagonalisable. Then using $O(N, \mathbb{C})$ we may assume $\beta = \text{diag}(\gamma_1, \gamma_2, \dots, \gamma_N)$ and we will assume $v = (\lambda_1, \dots, \lambda_N) \in \mathbb{R}^N$ although more generally $v \in \mathbb{C}^N$. Hence (β, v) is stable if and only if the γ_i are distinct. By (6.7)

$$q(z) = \left[1, \frac{\lambda_1}{z - \gamma_1}, \frac{\lambda_2}{z - \gamma_2}, \dots, \frac{\lambda_N}{z - \gamma_N} \right].$$

This gives the $t \rightarrow \infty$ limit of the holomorphic sphere (5.1) for $\lambda_0 = t = -\gamma_0$. The limit is still a hyperbolic monopole. Its grid is $\mathcal{G} = \{(\frac{-1}{\gamma_i}, \gamma_j) \mid 0 \leq i \neq j \leq N\}$ where $\gamma_0 = \infty$ so it contains the points $(0, \gamma_j)$ and $(\frac{-1}{\gamma_j}, \infty)$ for $j = 1, \dots, N$. This is natural since in general $\mathcal{G} \subset \mathbb{P}_1 \times \mathbb{P}_1$. One can send any $\lambda_i = t = -\gamma_i$ to infinity, and it also corresponds neatly to the discrete Nahm data by a permutation.

The spectral curve and rational map each behave well under this limit. The spectral curve is given by

$$0 = p(\eta, \zeta) = \prod_{j=1}^N (\zeta - \gamma_j)(1 + \eta \bar{\gamma}_j) - \eta \sum_{i=1}^N |\lambda_i|^2 \prod_{\substack{j=1 \\ j \neq i}}^N (\zeta - \gamma_j)(1 + \eta \bar{\gamma}_j)$$

The rational map is of the form used by Donaldson in [13].

$$\mathcal{R}_0(z) = \sum_{i=1}^N \frac{\lambda_i^2}{z - \gamma_i}.$$

It is obtained from $\mathcal{R}_0(z) = v^T(\beta - z)^{-1}v$ and agrees with the limit of the rational map (3.4) $\mathcal{R}_0(z) = \lim_{t \rightarrow \infty} \mathcal{R}(z)$ for $\lambda_0 = t = -\gamma_0$.

The behaviour of the grid, spectral curve, rational map and holomorphic sphere under $\zeta_i \rightarrow \infty$ suggests that JNR data can naturally be generalised to allow $\zeta_i \in \mathbb{P}_1$.

7. PROJECTION TO RATIONAL MAPS

The holomorphic sphere $q : \mathbb{P}_1 \rightarrow \mathbb{P}_N$ uniquely determines the monopole (A, Φ) up to gauge transformation. The rational map is defined by choosing a point on the two-sphere at infinity $p \in S_\infty^2$, restricting gauge transformations to limit to the identity at p , i.e. framed gauge transformations, then scattering from p . As described in Section 3.3, the full group of gauge transformations reduces to $U(1)$ in the limit at the point p , so if we do not restrict to based gauge transformations at p , the rational map is well-defined up to rescaling by $e^{i\theta}$.

In [19] we show in Theorem 3 that, for each monopole, there is a linear map $\mathbb{P}_N \rightarrow L_p = \mathbb{P}_1 \subset \mathbb{P}_N$ which composed with the holomorphic sphere q gives the rational map up to rescaling. The point $p \in S_\infty^2$ is the scattering direction which is chosen to be ∞ in Section 3.3, and the rational map takes the form (3.4).

Lemma 7.1. *There exists a unique $a = (a_0, a_1, \dots, a_N) \in \mathbb{C}^{N+1}$ such that the rational map is given by*

$$\mathcal{R}(z) = \frac{\frac{a_0 \lambda_0}{z - \gamma_0} + \frac{a_1 \lambda_1}{z - \gamma_1} + \dots + \frac{a_N \lambda_N}{z - \gamma_N}}{\frac{\lambda_0^2}{z - \gamma_0} + \frac{\lambda_1^2}{z - \gamma_1} + \dots + \frac{\lambda_N^2}{z - \gamma_N}} = \frac{\langle \bar{a}, q(z) \rangle}{\langle q(\infty), q(z) \rangle}$$

Proof. To simplify the calculation assume that $\sum_{i=0}^N \lambda_i^2 = 1$. This just changes the definition of the a_i . If we divide through by $\prod_{k=0}^N (z - \gamma_k)$ we get the correct denominator and the numerator becomes

$$\begin{aligned} \sum_{i=0}^N \sum_{j=i+1}^N \lambda_i^2 \lambda_j^2 \frac{(\gamma_i - \gamma_j)^2}{(z - \gamma_i)(z - \gamma_j)} &= \frac{1}{2} \sum_{i,j=0}^N \lambda_i^2 \lambda_j^2 \frac{(\gamma_i - \gamma_j)^2}{(z - \gamma_i)(z - \gamma_j)} \\ &= \frac{1}{2} \sum_{i,j=0}^N \lambda_i^2 \lambda_j^2 \left(\frac{\gamma_i - \gamma_j}{z - \gamma_i} + \frac{\gamma_j - \gamma_i}{z - \gamma_j} \right) \\ &= \sum_{i,j=0}^N \lambda_i^2 \lambda_j^2 \frac{\gamma_i - \gamma_j}{z - \gamma_i} \\ &= \sum_{i=0}^N \left(\sum_{j=0}^N \lambda_i \lambda_j^2 (\gamma_i - \gamma_j) \right) \frac{\lambda_i}{z - \gamma_i}. \end{aligned}$$

This gives the required result with

$$a_i = \sum_{j=0}^N \lambda_i \lambda_j^2 (\gamma_i - \gamma_j).$$

Note that $a_0 \lambda_0 + a_1 \lambda_1 + \dots + a_N \lambda_N = 0$ so that $\mathcal{R}(\infty) = 0$. \square

Hence

$$\mathcal{R}(z) = \frac{\frac{\gamma_0 \lambda_0^2}{z-\gamma_0} + \frac{\gamma_1 \lambda_1^2}{z-\gamma_1} + \dots + \frac{\gamma_N \lambda_N^2}{z-\gamma_N}}{\frac{\lambda_0^2}{z-\gamma_0} + \frac{\lambda_1^2}{z-\gamma_1} + \dots + \frac{\lambda_N^2}{z-\gamma_N}} - \langle \gamma \rangle$$

where

$$\langle \gamma \rangle = \frac{\sum_{j=0}^N \lambda_j^2 \gamma_j}{\sum_{j=0}^N \lambda_j^2}.$$

Thus $\mathcal{R}(z)$ is achieved up to scale by projecting $q(z)$ onto the line $L \subset \mathbb{P}_N$ spanned by

$$[\lambda_0, \dots, \lambda_N], \quad [\bar{\gamma}_0 \lambda_0, \dots, \bar{\gamma}_N \lambda_N]$$

which uses Gram-Schmidt as follows. Set $v_1 = (\lambda_0, \dots, \lambda_N)$. Then

$$v_2 = (\bar{\gamma}_0 \lambda_0, \dots, \bar{\gamma}_N \lambda_N) - \langle \bar{\gamma} \rangle v_1$$

is orthogonal to v_1 and projection is given by

$$Pq(z) = \frac{\langle v_1, q(z) \rangle v_1}{\langle v_1, v_1 \rangle} + \frac{\langle v_2, q(z) \rangle v_2}{\langle v_2, v_2 \rangle}$$

which is essentially $\mathcal{R}(z)$ except for a scale factor of

$$\frac{\langle v_2, v_2 \rangle}{\langle v_1, v_1 \rangle}.$$

Instead of scattering from $\infty \in S_\infty^2$, as one might expect, the rational map obtained by scattering from γ_i is simpler. It is of the form:

$$\mathcal{R}_{\gamma_i}(z) = (z - \gamma_i) \sum_{j \neq i} \frac{a_{ij} \lambda_i}{z - \gamma_j}$$

for some $a_{ij} \in \mathbb{C}$ which is obtained by projection of q onto $L_\zeta = \mathbb{P}_1 \subset \mathbb{P}_N$.

8. THE ENERGY DENSITY AT INFINITY

From [19] we know that the energy density of the monopole at infinity is given by $q^* \omega$ where ω is the Kähler form on \mathbb{P}_N and $q: \mathbb{P}_1 \rightarrow \mathbb{P}_N$ the holomorphic sphere. There are various ways to calculate this and we will use

$$(8.1) \quad E(q) = \frac{|q \wedge \partial q|^2}{|q|^4} (1 + |z|^2)^2.$$

Here we have lifted q to a map into \mathbb{C}^{N+1} and we regard $q \wedge \partial q$ as an element of $\bigwedge^2 \mathbb{C}^{N+1}$ with the inner product induced from \mathbb{C}^{N+1} .




FIGURE 8.1. Graph of the energy density at infinity for $N = 3$ JNR monopole

If we let e^0, \dots, e^N be the standard basis of \mathbb{C}^{N+1} then from (5.3) we can take

$$q(z) = \sum_{i=0}^N \frac{\lambda_i}{z - \gamma_i} e^i.$$

so that

$$\partial q(z) = \sum_{i=0}^N \frac{-\lambda_i}{(z - \gamma_i)^2} e^i.$$

and thus

$$q \wedge \partial q = \sum_{0 \leq i < j \leq N} \left(\frac{\lambda_i}{(z - \gamma_i)^2} \frac{\lambda_j}{(z - \gamma_j)} - \frac{\lambda_j}{(z - \gamma_j)^2} \frac{\lambda_i}{(z - \gamma_i)} \right) e^i \wedge e^j$$

Substituting into the formula for the energy density (8.1) we obtain

$$(8.2) \quad E(q)(z) = \frac{\sum_{0 \leq i < j \leq N} \left| \frac{\lambda_i}{(z - \gamma_i)^2} \frac{\lambda_j}{(z - \gamma_j)} - \frac{\lambda_j}{(z - \gamma_j)^2} \frac{\lambda_i}{(z - \gamma_i)} \right|^2}{\left| \sum_{i=0}^N \left| \frac{\lambda_i}{(z - \gamma_i)} \right|^2 \right|^2} (1 + |z|^2)^2$$




FIGURE 8.2. Contours of the energy density at infinity for the $N = 3$ tetrahedrally symmetric JNR monopole

A calculation shows that

$$(8.3) \quad E(q)(\gamma_k) = \sum_{0 \leq j \neq k \leq N} \left(\frac{\lambda_j}{\lambda_k} \right)^2 \frac{(1 + |\gamma_k|^2)^2}{|\gamma_k - \gamma_j|^2}.$$

Example 8.1. Take $N = 3$, $\lambda_0 = 1$, $\lambda_1 = 0.8$, $\lambda_2 = 0.6$, $\lambda_3 = 0.6$, $\gamma_0 = 1$, $\gamma_1 = i$, $\gamma_2 = -1$ and $\gamma_3 = -i$. Then we obtain the energy density in 8.1.

Example 8.2. If we take the example of the $N = 3$ tetrahedrally symmetric monopole from [10] we obtain the energy density in 8.2.

If the parameters are varied by lowering all the values of λ_i except say the k th one uniformly towards 0 then as discussed in Section 3.2 we expect that the monopole (at least its spectral curve) approaches a monopole located near the points $\gamma_0, \dots, \gamma_{k-1}, \gamma_{k+1}, \gamma_N$ at infinity. Similarly we see here that $E(q)(z)$ approaches a sum of delta functions at $\gamma_0, \dots, \gamma_{k-1}, \gamma_{k+1}, \gamma_N$.

In principle it should be possible to calculate the information metric (see for example [18]) induced from the energy density at infinity in terms of the JNR data.

REFERENCES

- [1] M.F. Atiyah, Instantons in two and four dimensions, *Commun. Math. Phys.*, **93**(4), 437–451, 1984.
- [2] M.F. Atiyah, Magnetic monopoles in hyperbolic space, *Proc. Bombay Colloq. on: Vector Bundles on Algebraic Varieties*, 1–34 (1987).
- [3] M.F. Atiyah, *The geometry and physics of knots*, In Miniconference on Geometry and Physics. Michael N. Barber and Michael K. Murray, eds. Proceedings of the Centre for Mathematical Analysis, v. 22. (Centre for Mathematics and its Applications, Mathematical Sciences Institute, The Australian National University, 1989), 1–17. <https://projecteuclid.org/euclid.pcma/1416406250>
- [4] M.F. Atiyah, Magnetic monopoles and the Yang-Baxter equations, *Internat. J. Modern Phys. A*, **6**(16), 2761–2774, 1991. Topological methods in quantum field theory (Trieste, 1990).
- [5] M.F. Atiyah, N.J. Hitchin, V.G. Drinfeld and Yu.I. Manin, Construction of instantons, *Phys. Lett. A*, **65**(3), 185–187, 1978.
- [6] M.F. Atiyah and N. Hitchin. *The Geometry and Dynamics of Magnetic Monopoles*. M. B. Porter Lectures. Princeton University Press, Princeton, NJ, 1988.
- [7] Monopoles and Yang-Baxter equations, M.F. Atiyah and M.K. Murray, in *Further Advances in Twistor Theory* Volume 2, Pitman, London, 1995.
- [8] D.M. Austin and P.J. Braam. Boundary values of hyperbolic monopoles. *Nonlinearity*, **3**, 809–823 (1990).
- [9] R.J. Baxter, J.H.H. Perk and H. Au-Yang, New solutions of the star-triangle relations for the Chiral-Potts model, *Phys. Lett. A*, **128**(3,4), 138–142, 1988.
- [10] S. Bolognesi, A. Cockburn and P. Sutcliffe, Hyperbolic monopoles, JNR data and spectral curves, *Nonlinearity*, **28**(1), 211–235, 2015.
- [11] A.H. Cockburn, *Aspects of Vortices and Hyperbolic Monopoles*, Durham theses, Durham University. Available at Durham E-Theses Online: <http://etheses.dur.ac.uk/1107>, 2015.
- [12] E. Corrigan and D.B. Fairlie, Scalar field theory and exact solutions to a classical $SU(2)$ gauge theory, *Phys. Lett.*, **67B**(1), 69–71, 1977.
- [13] S.K. Donaldson, *Nahm's equations and the classification of monopoles*, *Commun. Math. Phys.*, **96**, 387–407, 1984.
- [14] N.J. Hitchin, Monopoles and geodesics, *Commun. Math. Phys.*, **83**(4), 579–602, 1982.
- [15] R. Jackiw, C. Nohl and C. Rebbi, Conformal properties of pseudoparticle configurations. *Phys. Rev. D* **15**, 1642–1646, 1977.
- [16] N.S. Manton and P.M. Sutcliffe, Platonic Hyperbolic Monopoles, *Commun. Math. Phys.*, **325**, 821–845, 2014.
- [17] *Integrability, Self-duality, and Twistor Theory*, L.J. Mason and N.M.J. Woodhouse, Oxford University Press, Oxford, 1996.
- [18] M.K. Murray. The information metric on rational maps. *Experiment. Math.*, **2**(4), 271–279, 1993.
- [19] M.K. Murray, P. Norbury and M.A. Singer. Hyperbolic monopoles and holomorphic spheres. *Ann. Global Anal. Geom.*, **23**(2), 101–128, 2003.
- [20] M. Murray and M. Singer. Spectral curves of non-integral hyperbolic monopoles. *Nonlinearity*, **9**(4), 973–997, 1996.
- [21] M.K. Murray and M.A. Singer. On the complete integrability of the discrete Nahm equations. *Commun. Math. Phys.*, **210**(2), 497–519, 2000.
- [22] P. Norbury and N.M. Romão, Spectral curves and the mass of hyperbolic monopoles, *Commun. Math. Phys.* **270**(2), 295–333, 2007.

(Michael K. Murray) SCHOOL OF MATHEMATICAL SCIENCES, UNIVERSITY OF ADELAIDE, ADELAIDE, SA 5005, AUSTRALIA

Email address, Michael K. Murray: `michael.murray@adelaide.edu.au`

(Paul Norbury) SCHOOL OF MATHEMATICS AND STATISTICS, UNIVERSITY OF MELBOURNE, VIC 3010, AUSTRALIA

Email address, Paul Norbury: `norbury@unimelb.edu.au`