



Contents available at [ScienceDirect](https://www.sciencedirect.com)

Diabetes Research
and Clinical Practice

journal homepage: www.elsevier.com/locate/diabres



International
Diabetes
Federation



Women with type 1 diabetes exhibit a progressive increase in gut *Saccharomyces cerevisiae* in pregnancy associated with evidence of gut inflammation



Esther Bandala-Sanchez^{a,b,1}, Alexandra J. Roth-Schulze^{a,b,1}, Helena Oakey^c, Megan A.S. Penno^c, Naiara G. Bediaga^{a,b}, Gaetano Naselli^{a,b}, Katrina M. Ngui^{a,b}, Alannah D. Smith^{a,b}, Dexing Huang^{a,b}, Enrique Zozaya-Valdes^{a,b}, Rebecca L. Thomson^c, James D. Brown^c, Peter J. Vuillermin^d, Simon C. Barry^c, Maria E. Craig^{e,f}, William D. Rawlinson^{g,h}, Elizabeth A. Davisⁱ, Mark Harris^{j,k}, Georgia Soldatos^l, Peter G. Colman^m, John M. Wentworth^{a,b,m}, Aveni Haynesⁱ, Grant Morahanⁿ, Richard O. Sinnott^o, Anthony T. Papenfuss^{p,q,r,s}, Jennifer J. Couper^{c,t}, Leonard C. Harrison^{a,b,*}, on behalf of the ENDIA Study Group

^a Walter and Eliza Hall Institute of Medical Research, Melbourne, Australia

^b Department of Medical Biology, University of Melbourne, Melbourne, VIC, Australia

^c The University of Adelaide, Robinson Research Institute, Adelaide Medical School, University of Adelaide, Adelaide, SA, Australia

^d Faculty of School of Medicine, Deakin University and Child Health Research Unit, Barwon Health, Geelong, VIC, Australia

^e School of Women's and Children's Health, Faculty of Medicine, University of New South Wales, Sydney, NSW, Australia

^f Institute of Endocrinology and Diabetes, The Children's Hospital at Westmead, Sydney, NSW, Australia

^g Virology Research Laboratory, Serology and Virology Division, South Eastern Area Laboratory Services Microbiology, NSW Health Pathology, Sydney, NSW, Australia

^h School of Medical Sciences, Biotechnology and Biomolecular Sciences, Women's and Children's Health, University of New South Wales, Sydney, NSW, Australia

ⁱ Telethon Institute for Child Health Research, Centre for Child Health Research, University of Western Australia, Perth, WA, Australia

^j The University of Queensland Diamantina Institute, Faculty of Medicine, University of Queensland, Translational Research Institute, Woolloongabba, QLD, Australia

^k Queensland Children's Hospital, South Brisbane, QLD, Australia

^l Monash Centre for Health Research and Implementation, School of Public Health and Preventive Medicine, Monash University, Melbourne and Diabetes and Vascular Medicine Unit, Monash Health, Melbourne, VIC, Australia

^m Department of Diabetes and Endocrinology, Royal Melbourne Hospital, Melbourne, VIC, Australia

Abbreviations: ASCA, Anti-*Saccharomyces cerevisiae* antibodies; ASV, Amplicon sequence variants; bp, Base pair; CPM, Counts per million; DQESv2, Dietary Questionnaire for Epidemiological Studies version 2; ENDIA, Environmental determinants of islet autoimmunity; FDR, False discovery rate; GEE, Generalized estimating equations; HbA1c, Haemoglobin A1c; HLA, Human leukocyte antigen; HREC, Human research ethics committee; I-FABP, Intestinal fatty-acid binding protein; IgA, Immunoglobulin A; IgG, Immunoglobulin G; ITS1, Internal transcribed spacer 1; LCBD, Local contribution to beta diversity; LogFC, Log2 fold-change; mg, milligram; OTU, Operational taxonomic unit; PCoA, Principal components analysis; RMA-PERMANOVARE, Repeated measure-aware permutation analysis of variance; SCFA, Short-chain fatty acid; SD, Standard deviation; T1D, Type 1 diabetes; WEHI, Walter and Eliza Hall Institute of Medical Research; WMS, Whole metagenomic sequencing; Zinb, Zero-inflated negative binomial regression; 1,5-AG, 1,5-Anhydroglucitol

* Corresponding author.

E-mail address: harrison@wehi.edu.au (L.C. Harrison).

¹ These authors contributed equally to this work and share first authorship.

<https://doi.org/10.1016/j.diabres.2022.109189>

0168-8227/© 2022 The Authors. Published by Elsevier B.V.

This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

ⁿ Centre for Diabetes Research, Harry Perkins Institute of Medical Research, The University of Western Australia, Perth, WA, Australia

^o Melbourne eResearch Group, School of Computing and Information Services, University of Melbourne, Melbourne, VIC, Australia

^p Walter and Eliza Hall Institute of Medical Research, Melbourne, VIC, Australia

^q Department of Medical Biology and School of Mathematics and Statistics, University of Melbourne, Melbourne, VIC, Australia

^r Bioinformatics and Cancer Genomics Laboratory, Peter MacCallum Cancer Centre, Melbourne, VIC, Australia

^s Sir Peter MacCallum Department of Oncology, University of Melbourne, Parkville, VIC, Australia

^t Women's and Children's Hospital, Adelaide, SA, Australia

ARTICLE INFO

Article history:

Received 15 September 2021

Received in revised form

21 December 2021

Accepted 24 December 2021

Available online 18 January 2022

Keywords:

Gut

Microbiome

Fungi

Mycobiome

Saccharomyces

Faecalibacterium prausnitzii

Pregnancy

Type 1 diabetes

ITS1

Inflammation

ABSTRACT

Aims: Studies of the gut microbiome have focused on its bacterial composition. We aimed to characterize the gut fungal microbiome (mycobiome) across pregnancy in women with and without type 1 diabetes.

Methods: Faecal samples (n = 162) were collected from 70 pregnant women (45 with and 25 without type 1 diabetes) across all trimesters. Fungi were analysed by internal transcribed spacer 1 amplicon sequencing. Markers of intestinal inflammation (faecal calprotectin) and intestinal epithelial integrity (serum intestinal fatty acid binding protein; I-FABP), and serum antibodies to *Saccharomyces cerevisiae* (ASCA) were measured.

Results: Women with type 1 diabetes had decreased fungal alpha diversity by the third trimester, associated with an increased abundance of *Saccharomyces cerevisiae* that was inversely related to the abundance of the anti-inflammatory butyrate-producing bacterium *Faecalibacterium prausnitzii*. Women with type 1 diabetes had higher concentrations of calprotectin, I-FABP and ASCA.

Conclusions: Women with type 1 diabetes exhibit a shift in the gut mycobiome across pregnancy associated with evidence of gut inflammation and impaired intestinal barrier function. The relevance of these findings to the higher rate of pregnancy complications in type 1 diabetes warrants further study.

© 2022 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

The autoimmune T-cell response that destroys insulin-producing beta cells in the pancreatic islets in type 1 diabetes (T1D) is thought to be promoted by environmental factors on a background of genetic susceptibility. A role for the gut bacterial microbiome, at the interface with the external environment, in the development of T1D is supported by animal models [1,2,3]. In humans, a decrease in compositional diversity and the relative abundance of potentially beneficial short-chain fatty acid-producing bacteria has been documented before and after the clinical presentation of T1D [4–11]. In addition, we recently reported that women with T1D exhibit a shift towards a more pro-inflammatory gut bacterial microbiome during pregnancy [12]. However, the gut microbiome is a complex ecosystem that comprises not only bacteria, but fungi, viruses, archaea and protozoa [13–16]. Fungi are ubiquitous in the environment and may interact with and modify other components of the microbiome, thus contributing directly and indirectly to the role of the microbiome in health and disease. Fungal species from the genera *Candida*, *Saccharomyces*, *Penicillium*, *Aspergillus*, *Cryptococcus*, *Malassezia*, *Cladosporium*, *Galactomyces*, *Debaryomyces* and *Trichosporon*

have been identified in the healthy human gut, representing 0.1%–1.0% of the gut microbiota [13,15,17]. Based on culture methods, the outgrowth of *Candida* genera in the gastrointestinal tract has been linked to poor glycaemic control in T1D [18,19,20], but the gut mycobiome has not been characterized in this disease or in pregnancy. In the present study, we analysed the gut mycobiome in each trimester of pregnancy in a cohort of women with and without T1D in the Australia-wide Environmental Determinants of Islet Autoimmunity (ENDIA) pregnancy-birth cohort study [21].

2. Methods

2.1. Participants

The study comprised 70 pregnancies in 70 women, 45 women with established T1D on daily insulin treatment and 25 women with no history of T1D, being a representative sample of those in the ENDIA pregnancy-birth cohort study. One hundred and sixty-two faecal samples were collected across all trimesters (34 in trimester 1, 65 in trimester 2 and 63 in trimester 3; [Supplementary Fig. 1](#)). These samples were also included in a recent study of the bacterial microbiome in

pregnancy [12]. The main criterion for participation in ENDIA was a child with a first-degree relative with T1D. Table 1 summarizes and compares characteristics of the T1D and non-T1D participants.

Women who provided written informed consent were enrolled in the study from 2013 to 2016 at eight clinical sites. Each had up to three study visits during pregnancy, ideally one in each trimester. The study was approved by a Human Research Ethics Committee (HREC) at each clinical site, with the Women's and Children's Hospital HREC in Adelaide acting as the lead under the Australian National Mutual Acceptance Scheme (reference number HREC/16/WCHN/066). ENDIA is registered on the Australia New Zealand Clinical Trials Registry (ACTRN1261300794707).

2.2. Collection-processing of faecal samples

ENDIA participants self-collected faecal samples and placed them into a sterile 70 mL collection jar, which was stored in home refrigerator for 6–24 h before delivery to the laboratory in an insulated container. Sample aliquots were transferred into 6x sterile 5 mL screw cap tubes and stored at -80°C until further processing as described [22].

2.3. DNA extraction

DNA was extracted from approximately 100 mg of faeces with the Zymobiomics kit (Zymo Research) as per manufacturer's instructions, quantified using a Qubit dsDNA BR (Broad Range) assay kit (Thermo Fisher Scientific) and stored at -80°C until further processing.

2.4. Fungal ITS1 amplification and sequencing

For fungal analysis, the internal transcribed spacer 1 (ITS1) region was first amplified from faecal DNA. ITS1 primers were ITS1F (CTTGGTCATTTAGAGGAAGTAA) and ITS1R (GCTCGCTTCTTCATCGATGC) [17]. Two PCRs were performed. Cycling conditions for the first PCR were: initial denaturation at 95°C for 3 min, 20 amplification cycles at 94°C for 30 s 56°C for 90 s and 72°C for 3 min, followed by a final extension step at 72°C for 10 min. Conditions for the second PCR to add unique index barcodes to each sample were: initial denaturation at 95°C for 3 min, 25 amplification cycles at 94°C for 45 s, 57°C for 60 s and 72°C for 90 s, followed by a final extension step at 72°C for 10 min. PCR products were visualized by agarose gel electrophoresis and quantified with Tape Station (Agilent 2100 Bioanalyzer). Equimolar amounts of purified amplicons were pooled and sequenced on the Illumina MiSeq platform using the Illumina MiSeq v3 600-cycle (2×300 bp) kit.

2.5. Fungal sequence analysis

ITS1 amplicon sequence data were processed with QIIME2 (version 2017.12) [23]. Briefly, Q2-dada2 denoise-paired [24] was used to construct features (amplicon sequence variants; ASV). Due to the variable size of the ITS1 region (300–900 bp) to avoid loss of information paired-end reads were not merged and posterior analysis was performed only on

single-end read mode using read 1. Representative sequences from read 1 were taxonomically classified using a naive Bayes classifier pre-trained with the UNITE (version 7-99 01.12.2017) eukaryotic nuclear ribosomal ITS region database [25], and a phylogenetic tree was built within qiime2 using first the mafft function to align the sequences and the fasttree function to infer the approximately-maximum-likelihood phylogenetic tree. Only sequences classified as fungi were analysed. The resulting taxonomic profile, i.e., features and their counts per sample, was imported into the phyloseq [26] package in R [27] and agglomerated into operational taxonomic units (OTUs) based on a phylogenetic tree (a cophenetic distance of 0.02 was used as proxy for the difference between fungal species) with the phyloseq function tip_glom. OTUs with a total abundance across samples $< 0.01\%$ of the total and unclassified OTUs at the Kingdom taxonomic level were removed. The resulting OTUs were further agglomerated based on species classification when available (i.e., where two or more OTUs were classified to the same species they were merged into one OTU). Due to the large difference in library sizes, samples were normalized, i.e., subsampled without replacement to the size of the smallest library (i. e. 1000 sequences) per sample using the phyloseq function rarefy_even_depth. Normalized data was used for all statistical analyses.

2.6. Bacterial sequence analysis

Bacterial sequences for a subset of 84 samples were obtained from metagenomic shotgun sequencing that had been performed in a separate study [12].

2.7. Estimation of fibre intake

Maternal diet during pregnancy was measured at the third trimester visit using a validated 74 item food frequency questionnaire, Dietary Questionnaire for Epidemiological Studies version 2 (DQESv2) [28]. Even though this was administered only in the third trimester, evidence for stability of dietary intake over the course of the pregnancy was obtained from a separate, purpose-built ENDIA Pregnancy Lifestyle Questionnaire administered before each of the three study visits during pregnancy. This assessed consumption of milk (dairy and non-dairy), caffeinated and decaffeinated tea and coffee, caffeine-containing soft drinks, dairy products, soy, gluten containing cereals (wheat, barley, and rye) and non-gluten containing cereals (rice, corn, and oats). Analysis across the study visits revealed that on 86% of occasions respondents reported either the same unit or within one-unit difference between visits 1–2, visits 1–3, and visits 2–3. Magnitude changes of four or five units were reported on $< 2\%$ of occasions. This supports the DQESv2 as being reflective of the whole pregnancy period.

2.8. Measurement of serum 1,5-anhydroglucitol (1,5-AG)

Serum 1,5-anhydroglucitol (1,5-AG), an index of glucose control in pregnancy [29], was measured by GlycoMark (Nippon Kayaku Co. Ltd., New York, NY, US).

Table 1 – Summary of characteristics of non-T1D and T1D pregnancies.

General	Non-T1D	T1D	P-value **
Overall number of samples: n (%)	59 (36)	103 (64)	
Trimester 1	15 (18.8)	23 (18.5)	
Trimester 2	21 (43.5)	40 (47.1)	
Trimester 3	23 (37.7)	40 (34.5)	
All three trimesters (% pregnancies)	12 (46)	17 (38)	
All three trimesters (% samples)	12 (20)	17 (17)	
Gestational age in days at faecal sample: mean (SD)			
Trimester 1	85.4 (18.7)	80.2 (14.7)	0.34
Trimester 2	164 (20.1)	159 (19.5)	0.31
Trimester 3	246 (16.9)	232 (15.0)	0.0011
Maternal			
Overall number of pregnancies	26	45	
Age in years at conception: mean (SD)	32.6 (3.24)	31.5 (4.28)	0.29
Assisted conception: n (%)	2 (8)	4 (9)	0.86
Twin pregnancy: n (%)	0 (0.0)	0 (0.0)	
Nulliparous: n (%)	7 (26.9)	26 (57.8)	0.014
Pre-eclampsia: n (%)	0 (0)	8 (18)	0.0050
Group B Streptococcus positive: n (%)	6 (23)	5 (11)	0.19
Genito-urinary infections: n (%)	2 (8)	4 (9)	0.86
Pre-pregnancy BMI: mean (SD)	26.1 (5.59)	26.0 (5.00)	0.998
Underweight (<18.5): n (%)	0 (0)	1 (2)	
Normal weight (18.5–24.9): n (%)	14 (54)	18 (40)	
Overweight weight (25–29.9): n (%)	5 (19)	15 (33)	
Obese (>30): n (%)	7 (27)	11 (24)	
Gestational weight gain (kg): Mean (SD)	12.6 (3.99)	11.9 (5.02)	0.51
Paternal			
Age in years at conception: mean (SD)	34.9 (6.32)	32.9 (3.85)	0.090
Pre-pregnancy BMI: mean (SD)	27.7 (4.15)	27.6 (4.14)	0.89
Underweight (BMI < 18.5): n (%)	0 (0)	0 (0)	
Normal weight (BMI 18.5–24.9): n (%)	4 (15)	8 (18)	
Overweight (BMI 25–29.9): n (%)	10 (38)	9 (20)	
Obese (>30): n (%)	4 (15)	11 (24)	
Missing: n (%)	8 (31)	17 (38)	
Maternal demographics			
Born in Australia: n (%)			
Yes	21 (81)	39 (87)	0.51
Education beyond High School: n (%)			
Yes	21 (81)	37 (82)	0.88
Lives in a metro area: n (%)	24 (92)	41 (91)	0.86

General	Non-T1D	T1D	P-value **
Socio-Economic Indexes for Areas (SEIFA) Index of Relative Socio-Economic Disadvantage (IRSD)			
Quintile 1n (%)	2 (8)	2 (4)	
Quintile 2n (%)	2 (8)	5 (11)	
Quintile 3n (%)	6 (23)	9 (20)	
Quintile 4n (%)	5 (19)	10 (22)	
Quintile 5n (%)	11 (42)	19 (42)	0.90
Smoking during pregnancy: n (%)	1 (4)	1 (2)	0.67
Household smoking during pregnancy: n (%)	3 (12)	5 (11)	0.91
Adults in house during pregnancy: n (%)			
One	0 (0)	3 (7)	
Two	24 (92)	39 (87)	
More than two	1 (4)	3 (7)	0.62
Children in house during pregnancy: n (%)			
None	7 (27)	24 (53)	
One	7 (27)	13 (29)	
Two	7 (27)	6 (13)	
More than two	4 (15)	1 (2)	0.0068
Furred pet ownership during pregnancy: n (%)	14 (54)	27 (60)	0.66
Diet and physical activity in pregnancy			
Diet: mean (SD)			
Energy/day (kJ)	7770 (1710)	6950 (2180)	0.14
Fat (g)	82.2 (20.3)	78.7 (27.7)	0.61
Protein (g)	88.5 (17.9)	86.5 (28.3)	0.77
Carbohydrate (g)	193 (47.6)	152 (52.4)	0.0042
Fiber (g)	20.6 (4.18)	19.9 (5.83)	0.61
Diet: Missing: n (%)	4 (15)	7 (16)	
Alcohol consumed: n (%)			
Yes	6 (23)	6 (13)	
Unknown	1 (4)	1 (2)	0.28
Total level of physical activity (MET) (h/week): mean (SD) [65]	258 (97.1)	278 (110)	0.57
Biological data			
HbA1c (%)			
Trimester 1: median (IQR)	-	6.5 (1.1)	
Trimester 2: median (IQR)	-	6.2 (0.95)	
Trimester 3: median (IQR)	-	6.4 (0.75)	
Trimester 1: missing	-	4 (17)	
Trimester 2: missing	-	5 (13)	
Trimester 3: missing	-	14 (35)	
Serum 1,5-anhydroglucitol (AG) (ug/mL)			
Trimester 1: median (IQR)	14.1 (11.1)	3.2 (1.8)	
Trimester 2: median (IQR)	10.8 (6.2)	2.4 (2.3)	
Trimester 3: median (IQR)	9.2 (5.4)	2.4 (2.1)	
Trimester 1: mean (SD)	14.3 (5.3)	3.92 (2.5)	<0.001
Trimester 2: mean (SD)	11.0 (3.9)	2.66 (1.9)	<0.001
Trimester 3: mean (SD)	9.34 (3.7)	2.79 (2.0)	<0.001

General	Non-T1D	T1D	P-value **
Trimester 1: missing n (%)	0 (0)	0 (0)	
Trimester 2: missing n (%)	0 (0)	3 (8)	
Trimester 3: missing n (%)	0 (0)	9 (23)	
Serum vitamin D (nmol/L): mean (SD)			
Trimester 1	83.5 (23.8)	81.0 (22.6)	0.74
Trimester 2	95.1 (24.0)	89.3 (26.7)	0.34
Trimester 3	88.8 (31.3)	94.5 (31.6)	0.52
Trimester 1: missing n (%)	2 (13)	1 (4)	
Trimester 2: missing n (%)	1 (5)	3 (8)	
Trimester 3: missing n (%)	2 (9)	6 (15)	
Maternal HLA: n (%)			
DR34	14 (54)	22 (49)	
DR3 or DR4	4 (15)	18 (40)	
DRXX	8 (31)	5 (11)	0.031
Known supplements pre-pregnancy and pregnancy			
Antibiotics: n (%)	NA	NA	NA
	7 (27)	14 (31)	0.71
Anticoagulants: n (%)	0 (0)	1 (2)	0.998
	3 (12)	16 (36)	0.021
Antihypertensive agents: n (%)	NA	NA	NA
	0 (0)	9 (20)	0.0028
Known other supplements pre-pregnancy and pregnancy			
Biotin: n (%)	10 (39)	9 (20)	0.095
	23 (89)	41 (91)	0.72
Calcium: n (%)	11 (42)	10 (22)	0.078
	23 (89)	43 (96)	0.28
Iron	12 (46)	9 (20)	0.023
	25 (96)	44 (98)	0.69
Magnesium: n (%)	12 (46)	9 (20)	0.023
	24 (92)	42 (93)	0.87
Selenium: n (%)	10 (39)	9 (20)	0.095
	23 (89)	41 (91)	0.72
Vitamin B1: n (%)	12 (46)	9 (20)	0.023
	24 (92)	42 (93)	0.87
Vitamin B2: n (%)	12 (46)	9 (20)	0.023
	24 (92)	42 (93)	0.87
Vitamin B3: n (%)	12 (46)	9 (20)	0.023
	24 (92)	42 (93)	0.87
Vitamin B5: n (%)	6 (23)	7 (16)	0.43
	12 (46)	29 (64)	0.14
Vitamin B6: n (%)	12 (46)	9 (20)	0.023
	24 (92)	42 (93)	0.87
Vitamin B9 (folate): n (%)	11 (42)	14 (31)	0.34
	25 (96)	45 (100)	0.997
Vitamin B12: n (%)	12 (46)	9 (20)	0.023
	24 (92)	41 (91)	0.86

General	Non-T1D	T1D	P-value **
Vitamin D: n (%)	12 (46)	10 (22)	0.039
	25 (96)	43 (96)	0.90
Vitamin E: n (%)	11 (42)	9 (20)	0.048
	21 (81)	38 (84)	0.69
OTHER			
Vaccine: n (%)			
Yes (Flu only)	1 (4)	4 (9)	
Yes (Pertussis only)	3 (12)	7 (16)	
Yes (Flu and Pertussis)	9 (35)	14 (31)	0.80
Mode of delivery: n (%)			
Vaginal	20 (77)	15 (33)	
Caesarean (with labour)	0 (0)	8 (18)	
Caesarean (without labour)	6 (23)	22 (49)	<0.001
Log transformation was used for Maternal BMI, Metabolic equivalent of task, 1,5-anhydroglucitol in trimesters 1 and 2 and for Vitamin D in trimesters 1 and 3.			
Square root transformation was used for 1,5-anhydroglucitol in trimester 3 and Vitamin D in trimester 2.			
Nine questionnaires had to be excluded due to unrealistic dietary intake (energy < 4500 kJ/day or >20,000 kJ/day), as previously reported in other pregnancy studies (http://dx.doi.org/10.1111/j.1740-8709.2007.00104.x).			
Hb1Ac, serum 1,5-AG ($\mu\text{g/mL}$) and serum vitamin D are based on samples rather than pregnancies			

2.9. Measurement of faecal calprotectin

Calprotectin, an index of intestinal inflammation [30], was measured in faecal samples from trimester three by enzyme-linked immunoassay (CALPRO™ Oslo, Norway) according to the manufacturer's instructions. The protein extracts were prepared using 50 mg of the same faecal sample used for the mycobiome analysis. The concentration of faecal calprotectin was expressed as milligrams per kg, the normal range being 5–50 mg/kg [31,32].

2.10. Measurement of serum intestinal fatty-acid binding protein (I-FABP)

Serum intestinal fatty acid-binding protein (I-FABP), a marker of intestinal epithelial damage [33], was measured across the three trimesters by enzyme-linked immunoassay (Hycult Biotech, The Netherlands) according to the manufacturer's instructions. The concentration of I-FABP was expressed as pg/mL, the normal range being 20–485 pg/mL [34].

2.11. Measurement of serum IgA/IgG antibodies to *Saccharomyces cerevisiae*

Serum IgA and IgG antibodies specific for *Saccharomyces* (ASCA) were measured across the three trimesters of pregnancy by enzyme-linked immunoassay (QUANTA Lite ASCA, Inova Diagnostics, San Diego, CA, US) according to the manufacturer's instructions. Results were expressed as arbitrary units based on the values of the positive and negative controls, with a cut-off for positivity of 10 U/mL.

2.12. Statistical analyses

Alpha diversity (diversity within microbial communities) was obtained from the number of observed OTUs and InvSimpson index using the function `estimate_richness` from the R package `phyloseq`. For testing differences in alpha diversity between women with and without T1D, Generalized Estimating Equations (GEEs) [35] were applied using the R function `geeglm` from package `geepack` v1.2-1 [36] to account for possible correlation of multiple measurements within a participant over time. Parameter family was set to default "Gaussian". The default empirical (robust or 'sandwich') estimator was used to ensure that estimates were robust to misspecification of the correlation structure. The model used for the regression included T1D status and trimester as well as their interaction term (T1D × trimester) to test if differences in alpha diversity between T1D and non-T1D women changed across trimesters and was adjusted for the human leukocyte antigen (HLA) category. Beta diversity (diversity between microbial communities) was determined with `phyloseq` (function `distance`, `method="bray"`) on proportional log transformed data. This function calculates Bray-Curtis coefficients, which mea-

sure the distance between communities based on the taxa that they contain and their abundances. Differences in beta diversity were evaluated with a modified version of the Adonis function from the `vegan` package [37], which performs a repeated measure aware (RMA) PERMANOVA test [38]. This statistical model included T1D status and time with their interaction adjusted for HLA category as in the alpha diversity model. In addition, interactions between time and other factors were also tested as shown in the Results. Due to the small number of taxa (at each taxonomic level from OTU to phylum) with a prevalence greater than 50% in women with and without T1D, differential abundance analysis was performed in a univariate manner, i.e., a test was applied to each individual taxon. Given that zero inflation and over dispersion were present in the data a zero inflated negative binomial generalized linear mixed model, using the NB1 parametrization was the preferred model after considering other models in this analysis. This was fit with the function `'glmmTMB'` from the `glmmTMB` R package [39]. The model included the T1Dstatus * trimester interaction term and was adjusted for HLA category and a random factor for mothers that accounts for the correlation between counts from the same mother at different trimesters. Because none of the predictors influenced the excess zero logistic part of the model, the `'ziformula'` parameter was set to 1. If the interaction term was statistically significant, the package `'emmeans'` [40] from R was used for post-hoc comparisons between T1D and non-T1D women at each trimester and for comparisons between trimesters within T1D or non-T1D women and for estimating marginal means and their 95% confidence intervals (CI), and as this is a complex model calculated by simulation. The latter was performed averaging over the HLA, considering the differing effects of each category of HLA (Table 1). Predicted counts, i.e. estimated marginal or adjusted means and standard errors generated based on the zero inflated negative binomial linear model for T1D and non-T1D participants within each trimester averaged over HLA were plotted. A P-value ≤ 0.05 was considered significant. Zero inflated negative binomial linear (zinb) mixed models were fit with the function `zeroinfl` (R package `pscl`) [41] to determine associations of the relative abundance of *S. cerevisiae* with dietary intake (fibre, fat, carbohydrate, and specific carbohydrates [resistant starch, soluble and insoluble fibre]) in average grams consumed daily, and with the concentrations of serum 1,5-AG, faecal calprotectin, serum I-FABP and serum ASCA IgA and IgG as well as with the abundance of bacteria. Models were fitted with two terms, one with six levels for the combination of T1D status and trimester and the other for the covariate/variable of interest. For this, a hierarchical approach to testing significance (i.e., backward elimination) was used in which the interaction was tested before the main effects in both parts of the model (i.e., negative binomial and zero inflated) using the likelihood ratio test, function `lrtest` from R package `lmtest` [42]. When the interaction was not significant the model was further

reduced to investigate the significance of the main effects, which determined if the association with *S. cerevisiae* was significant. Since the “mother id” variance was essentially zero, this random factor was not included in the models.

3. Results

3.1. Study population and sequencing output

Forty-five pregnant women with T1D and 25 without T1D each provided up to three faecal samples across pregnancy (162 samples in total; [Supplementary Fig. 1](#)) for analysis by ITS-amplicon sequencing using the ITS1 sub-region. After quality filtering, $13,241 \pm 8394$ (mean \pm SD) reads per sample were obtained with the Illumina MiSeq sequencing machine. Overall, a total of 234 fungal OTUs was identified with ranges per sample in women with T1D of 2–9 in trimester 1, 1–12 in trimester 2 and 1–14 in trimester 3 and in women without T1D of 2–8 in trimester 1, 1–9 in trimester 2 and 2–11 in trimester 3 (see [Fig. 1](#) for more details).

3.2. Characterization of gut fungi in women with and without T1D in pregnancy

Out of 194 OTUs that were annotated up to the phylum level, 62 contained on average 63% of the total sequences per sample; the rest were classified only as Fungi (i.e., kingdom

taxonomic level; [Supplementary Fig. 2](#)). As observed previously [17], most of fungal OTUs were specific to each individual. In general, samples were dominated by the phylum Ascomycota within which two OTUs were dominant, representing on average $\sim 50\%$ of the total OTUs. One was an unclassified fungus and the other was classified to the *S. cerevisiae* species ([Supplementary Fig. 2](#)). *S. cerevisiae* was present in 71.8% and 64.4% of women with and without T1D, at an average abundance of 42.3% and 34.9%, respectively. *Candida albicans* species, in the phylum Ascomycota, was present in 10.8% and 7.2% of women with and without T1D, at an average abundance of 56% and 30%, respectively.

In the alpha diversity analysis, a significant interaction ($P = 0.023$) was found between trimester and T1D status for fungal richness. This did not differ between women with and without T1D in trimester 1 or 2 ([Fig. 1A](#), and [1B](#); [Table 2](#)) but decreased significantly in trimester 3 in women with T1D ([Fig. 1C](#); [Table 2](#)). No interactions or significant differences in the InvSimpson index of alpha diversity were detected ([Supplementary Fig. 3](#); [Table 2](#)). Beta diversity between women with and without T1D was not significantly different at any taxonomic level ([Table 3](#)), as expected by the low OTU frequency across samples in women in either group ([Supplementary Fig. 4](#)). However, there was a significant difference between trimesters at the class and phylum level ([Table 3](#)). Differential abundance analysis was performed in a univariate manner for taxa with a prevalence $> 50\%$ in




Fig. 1 – Alpha diversity (richness) by T1D status of women across trimesters. Number of fungal OTUs (richness) observed in samples from women with (T1D; red) and without T1D (non-T1D; blue) in **A**) the first (T1), **B**) second (T2), and **C**) third (T3) trimesters during pregnancy. Boxes show the inter-quartile range, and the line within the box indicates the median. The * denotes a significant difference (P -value < 0.05) between T1D and non-T1D.

Table 2 – Alpha diversity comparison (P-values).

	Interaction (T1D status:Time)	T1D STATUS (MAIN EFFECT)	TRIMESTERS (MAIN EFFECT)
Richness	0.034	–	–
InvSimpson	0.058	0.48	0.79
INTERACTION TABLE: T1D vs. non-T1D within each trimester			
	T1	T2	T3
Richness	0.37	0.14	0.018
INTERACTION TABLE: Time (trimester) within each of T1D and non-T1D			
	Time in T1D	Time in non-T1D	
Richness	0.15	0.19	

Note: In bold are significant P-values (<=0.05).

INTERACTION TABLE: T1D vs. non-T1D within each trimester: P-values obtained after testing T1D vs. non-T1D in each trimester, after the interaction between trimester and T1D status was significant.

INTERACTION TABLE: Time (trimester) within each of T1D and non-T1D: P-values obtained after testing differences between trimesters in women with or without T1D after the interaction between trimester and T1D status was significant.

Interaction (T1D status:Time): P-value from testing the interaction between terms T1D status and Time/Trimester. In the statistical model, the interaction between variables T1D status and Trimester were included to test if the possible differences in alpha diversity between T1D and non-T1D depend or are conditioned to trimester or T1D status (e.g. if there are differences between T1D and non-T1D only in one trimester or if there are differences between trimesters only in women with T1D).

T1D STATUS (MAIN EFFECT): P-value from testing differences between T1D and non-T1D women across the three trimesters, once the interaction between T1D status and trimester was tested and found not to be significant.

TRIMESTERS (MAIN EFFECT): P-value from testing differences between trimesters in women with and without T1D, once the interaction between T1D status and trimester was tested and was not significant.

Table 3 – Beta diversity comparison (P-values).

Beta diversity comparison between women with and without T1D - significant p-values (<0.05) in bold.			
	Interaction (T1D status:Time)	T1D STATUS (MAIN EFFECT)	TRIMESTERS (MAIN EFFECT)
Species	0.058	0.92	0.43
Genus	0.17	0.76	0.07
Family	0.18	0.77	0.075
Order	0.088	0.45	0.051
Class	0.087	0.66	0.024
Phylum	0.15	0.37	0.014

Interaction (T1D status:Time): P-value from testing the interaction between terms T1D status and Time/Trimester. In the statistical model, the interaction between variables T1D status and Trimester were included to test if the possible differences in alpha diversity between T1D and non-T1D depend or are conditioned to trimester or T1D status (e.g. if there are differences between T1D and non-T1D only in one trimester or if there are differences between trimesters only in women with T1D).

T1D STATUS (MAIN EFFECT): P-value from testing differences between T1D and non-T1D women across the three trimesters, once the interaction between T1D status and trimester was tested and found not to be significant.

TRIMESTERS (MAIN EFFECT): P-value from testing differences between trimesters in women with and without T1D, once the interaction between T1D status and trimester was tested and was not significant.

women with or without T1D. The predicted counts, i.e. estimated marginal or adjusted means and standard errors generated based on the zero inflated negative binomial linear model of *S. cerevisiae* were significantly decreased in women with T1D in trimester 1 and then increased across pregnancy in women with T1D and decreased in women without T1D (Fig. 2; Table 4), being significantly higher in women with T1D in trimester 3. Similar patterns were observed for the genus *Saccharomyces*, the family *Saccharomycetaceae*, the order *Saccharomycetales*, the class *Saccharomycetes* and the phylum

Ascomycota. This very similar pattern observed at different taxonomic levels is most likely explained by the percentage of *Ascomycota* reads classified as *S. cerevisiae* (Fig. 2; Table 4).

3.3. Relationship between *Saccharomyces* and blood glucose control and diet in pregnancy

To determine if the abundance of *Saccharomyces* was related to blood glucose control or diet, zero inflated negative binomial linear mixed models were fit with serum 1,5-AG, and with




Fig. 2 – Differentially abundant taxa (mean \pm SEM of predicted counts) detected by ITS1 amplicon sequencing between T1D (red) and non-T1D (blue) women and between trimesters. Predicted counts: estimated marginal or adjusted count means and standard errors generated based on the zero inflated negative binomial linear model. Predicted count means are shown as a point in each trimester for differentially abundant taxa in women with (red) and without (blue) T1D. An * between points denotes a significant difference (P-value < 0.05) between the groups in that trimester. A + sign denotes a significant difference in abundance between trimesters 1 and 3 and the colour denotes if the difference was observed in T1D (red) or non-T1D (blue) women. A) genus and B) Phylum.

dietary components (fat, carbohydrate, resistant starch, and total soluble and insoluble fibre). No significant associations were found (Table 5).

3.4. Intestinal inflammation and anti-*Saccharomyces cerevisiae* antibodies in women with T1D

Faecal calprotectin, released from neutrophils and monocytes, is a marker of intestinal inflammation that has been associated with impaired integrity and increased permeability of the epithelial barrier [43]. Serum intestinal fatty acid-binding protein (I-FABP) is a marker of intestinal epithelial damage [33]. We measured faecal calprotectin in trimester three and serum I-FABP in each trimester. Both markers were increased in women with T1D (Fig. 3; Supplementary Table 2 and Supplementary Table 3) and were correlated with each other (Pearson $R^2 = 0.58$; $P = 0.001$). Because gut inflammation with impaired epithelial barrier function could promote exposure and immunity to *Saccharomyces* we measured serum IgG and IgA ASCA. Both were higher in women with T1D across pregnancy (Fig. 4; Supplementary Table 1) but did not correlate with the abundance of *S. cerevisiae*.

3.5. Relationship between *Saccharomyces cerevisiae* and gut bacteria

We sought to address inter-microbial kingdom relationships and determine if the increase in *S. cerevisiae* in women with T1D was associated with a decrease in bacteria that produce anti-inflammatory short chain fatty acids (SCFAs), as we had recently observed [12]. A “zinb” linear mixed model approach was used to search for associations between *S. cerevisiae* and bacterial taxa identified as SCFA producers by metagenomic sequencing [12]. The species and genera of SCFA producers are listed in Table 6. The higher abundance

of *S. cerevisiae* was associated with a significantly lower abundance of *Faecalibacterium prausnitzii* (Table 6; Fig. 5).

4. Discussion

As the fungal compartment is only 0.1–1.0% of the total microbiome [13,44,15,17] and is not well represented in whole genome metagenome sequencing [17], we employed fungal-specific ITS1 amplicon sequencing of faecal DNA to analyse the gut mycobiota in pregnancy and T1D. Similar to what we observed, a high proportion of the Ascomycota phylum and, to a lesser extent, the Basidiomycota phylum, has been reported in the gut [17,45,46]. We found that Ascomycota was the dominant phylum in pregnant women and observed a decrease in fungal alpha diversity across pregnancy, mainly in trimester 3 in women with T1D. At this time, *Saccharomyces*, the most predominant genus within the Ascomycota phylum, increased in women with T1D. Changes in the gut mycobiome across pregnancy were distinct between women with and without T1D.

The species *S. cerevisiae* is recognised to colonize the human gut [47,48] and in some human populations is more abundant than *Candida* spp [49]. *S. cerevisiae* is present in most diets and its abundance in the gut may depend on nutrient availability. However, a study that reported associations between nutrients and fungal populations in faecal samples did not find a relationship between the nutrient composition of the diet and *Saccharomyces*, either recent or long-term [50]. We also found no association between diet and the abundance of *Saccharomyces*, but our dietary assessment tool did not specifically define yeast intake.

Poor glycaemic control in diabetes is associated with overgrowth of some fungi, especially *Candida albicans*, which is more prevalent in the faeces of individuals with T1D or T2D [51,52]. In our study, *C. albicans* was present in faecal samples from a minority of women with and without T1D and, in con-

Table 4 – Differential abundance analysis (P-values).

Differential abundant taxa between T1D and non-T1D women				Differential abundant taxa between trimesters 1 and 3			
Taxonomic level	Trimester	Classification	P.value	Taxonomic level	T1D status	Classification	P.value
OTU	T1	Saccharomyces_cerevisiae	0.0006	OTU	T1D	Saccharomyces_cerevisiae	0.001
OTU	T2	Saccharomyces_cerevisiae	0.28	OTU	non-T1D	Saccharomyces_cerevisiae	0.004
OTU	T3	Saccharomyces_cerevisiae	0.015	Genus	T1D	Saccharomyces	0.001
Genus	T1	Saccharomyces	0.0006	Genus	non-T1D	Saccharomyces	0.004
Genus	T2	Saccharomyces	0.28	Family	T1D	Saccharomycetaceae	0.002
Genus	T3	Saccharomyces	0.015	Family	non-T1D	Saccharomycetaceae	0.005
Family	T1	Saccharomycetaceae	0.001	Order	T1D	Saccharomycetales	0.0003
Family	T2	Saccharomycetaceae	0.23	Order	non-T1D	Saccharomycetales	0.02
Family	T3	Saccharomycetaceae	0.017	Class	T1D	Saccharomycetes	0.0003
Order	T1	Saccharomycetales	0.005	Class	non-T1D	Saccharomycetes	0.02
Order	T2	Saccharomycetales	0.43	Phylum	T1D	Ascomycota	0.01
Order	T3	Saccharomycetales	0.004	Phylum	non-T1D	Ascomycota	0.48
Class	T1	Saccharomycetes	0.005				
Class	T2	Saccharomycetes	0.43				
Class	T3	Saccharomycetes	0.004				
Phylum	T1	Ascomycota	0.22				
Phylum	T2	Ascomycota	0.17				
Phylum	T3	Ascomycota	0.049				

Note: In bold are significant P-values (i.e. ≤ 0.05)

Table 5 – Results (p-values) from likelihood ratio test comparing zero inflated negative binomial models to test associations with the abundances of *S. cerevisiae*.

Counts (Negative binomial part of the model)			
	Interaction	Variable	Trimester_T1D
Carbohydrates (g)	0.315	0.68	0.22
Resistant starch (g)	0.243	0.422	0.21
Fibre (g)	0.447	0.693	0.241
Soluble fibre (g)	0.791	0.62	0.228
Insoluble fibre (g)	0.3	0.96	0.225
Fat (g)	0.376	0.932	0.234
1,5-anhydroglucitol (AG) (ug/mL)	0.814	0.944	0.208
HbA1c [only T1D women]	0.279	0.516	0.048
I-FABP (pg/mL)	0.71	0.411	0.189
Calprotectin (mg/kg) [only trimester 3]	0.603	0.76	0.65
ASCA IgA (U/mL)	0.069	0.719	0.064
ASCA IgG (U/mL)	0.009**	0.45	0.047

For one unit increase in ASCA IgG the odds of having *S. cerevisiae* decrease by $1 - \exp(-9.034) = 100\%$ in trimester 1 in women with T1D

Binomial (zero in flation part of the model)			
	Interaction	Variable	Trimester_T1D
Carbohydrates (g)	0.535	0.477	0.21
Resistant starch (g)	0.36	0.766	0.193
Fibre (g)	0.293	0.599	0.186
Soluble fibre (g)	0.627	0.514	0.188
Insoluble fibre (g)	0.005***	0.133	0.21
Fat (g)	0.106	41	0.169
1,5-anhydroglucitol (AG) (ug/mL)	0.622	0.82	0.613
HbA1c [only T1D women]	0.866	0.426	0.29
I-FABP (pg/mL)	0.462	0.705	0.527
Calprotectin (mg/kg) [only trimester 3]	0.251	0.01***	0.079

Interaction not significant in the full results below. So there is no association

ASCA IgA (U/mL)	0.561	0.056	0.484
ASCA IgG (U/mL)	0.123	0.481	0.53

For one unit increase in Calprotectin the odds of having *S. cerevisiae* decrease by $1 - \exp(-0.017) = 2\%$

***Complete results from zinb test for ASCA IgG association with *S. cerevisiae*
Call:
zeroinfl(formula = sac ~ ASCA IgG* t1dtri | 1, data = TTMetadataF, dist = "negbin")

Count model coefficients (negbin with log link):

	Estimate	Std. Error	z value	Pr(> z)
(Intercept)	6.193	0.790	7.837	4.61E-15
ASCA IgG	0.729	2.628	0.277	0.781
t1dtrino_T2	-1.448	1.075	-1.346	0.178
t1dtrino_T3	-1.484	0.992	-1.495	0.135
t1dtriyes_T1	0.911	1.073	0.849	0.396
t1dtriyes_T2	0.442	0.922	0.479	0.632
t1dtriyes_T3	0.472	0.970	0.487	0.626
FacI:t1dtrino_T2	4.146	4.236	0.979	0.328
FacI:t1dtrino_T3	2.239	3.445	0.65	0.516
FacI:t1dtriyes_T1	-9.034	3.298	-2.74	0.00615

For one unit increase in ASCA IgG the odds of having *S. cerevisiae* decrease by $1 - \exp(-9.034) = 100\%$ in trimester 1 in women with T1D

Table 5 – (continued)

FacI:t1dtriyes_T2	-1.672	2.758	-0.606	0.544	
FacI:t1dtriyes_T3	-1.693	3.050	-0.555	0.579	
Log(theta)	-0.134	0.156	-0.86	0.390	
Zero-inflation model coefficients (binomial with logit link):					
	Estimate	Std. Error	z value	Pr(> z)	
(Intercept)	-0.714	0.199	-3.58	0.000338	
***Complete results from zinb test for Calprotectin association with <i>S. cerevisiae</i>					
Call: zeroinfl(formula = sac ~ Calprotectin t1dtri + Calprotectin, data = TTMetadataF, dist = "negbin")					
Count model coefficients (negbin with log link):					
	Estimate	Std. Error	z value	Pr(> z)	
(Intercept)	5.533	0.294	18.85	<2e-16	
Calprotectin	0.001	0.002	0.665	0.506	
Log(theta)	0.008	0.279	0.027	0.978	
Count model coefficients (negbin with log link):					
	Estimate	Std. Error	z value	Pr(> z)	
(Intercept)	-0.396	0.616	-0.642	0.521	
t1dtriyes_T3	1.398	0.828	1.687	0.092	
Calprotectin	-0.017	0.011	-1.564	0.118	For one unit increase in Calprotectin the odds of having <i>S. cerevisiae</i> decrease by $1 - \exp(-0.017) = 2\%$
***Complete results from zinb test for insoluble fiber association with <i>S. cerevisiae</i>					
Call: zeroinfl(formula = sac ~ Calprotectin t1dtri + Insoluble fibre, data = TTMetadataF, dist = "negbin")					
Count model coefficients (negbin with log link):					
	Estimate	Std. Error	z value	Pr(> z)	
(Intercept)	5.982	0.313	19.099	<2e-16	
Insoluble fibre	0.003	0.033	0.095	0.924	
Log(theta)	-0.364	0.135	-2.698	0.007	
Count model coefficients (negbin with log link):					
	Estimate	Std. Error	z value	Pr(> z)	
(Intercept)	1.288	1.556	0.828	0.408	
Insoluble fibre	-0.095	0.134	-0.703	0.482	
t1dtrino_T2	-3.080	2.067	-1.49	0.136	
t1dtrino_T3	-6.964	3.257	-2.138	0.033	
t1dtriyes_T1	1.718	2.854	0.602	0.547	
t1dtriyes_T2	-2.280	1.832	-1.245	0.213	
t1dtriyes_T3	-3.794	1.875	-2.024	0.043	
FacI:t1dtrino_T2	0.178	0.182	0.975	0.330	
FacI:t1dtrino_T3	0.554	0.284	1.955	0.051	Interaction not significant

Table 5 – (continued)

FacI:t1dtriyes_T1	-0.641	0.438	-1.463	0.144
FacI:t1dtriyes_T2	0.110	0.171	0.644	0.520
FacI:t1dtriyes_T3	0.286	0.171	1.669	0.095

Note Trimester_T1D status was fitted as a single term with 6 levels corresponding to the combination of the 2 levels of T1D status and the 3 trimesters.

Sac: *Saccharomyces cerevisiae* abundance

Variable: This refers to the variable being tested, as named in each row (e.g. Carbohydrates [g], Resistant starch [g], etc)

Interaction: This refers to the p-value obtained from testing if there is as significant interaction between the “variable of interest” and the term “Trimester_T1D status”. If the interaction is significant, it means that the possible differences in alpha or beta diversity between T1D and non-T1D depend or are conditioned to trimester and/or T1D status (e.g. if there are differences between T1D and non-T1D only in one trimester or if there are differences between trimesters only in women with T1D).




Fig. 3 – Faecal calprotectin and serum intestinal fatty-acid binding protein (iFABP) in pregnant women. Boxplots show the distribution of concentrations of faecal calprotectin in trimester 3 and serum I-FABP across trimesters in women with (red) and without (blue) type 1 diabetes. The * denotes a significant difference between T1D and non-T1D.




Fig. 4 – Anti-*Saccharomyces cerevisiae* antibodies in pregnant women. Serum from pregnant women with and without T1D across pregnancy was assayed for ASCA antibodies by ELISA. A) IgA ASCAs are higher in women with T1D compared to women without T1D across pregnancy. B) IgG ASCAs are higher in women with T1D compared to women without T1D across pregnancy. Results are expressed as absorbance (OD450). Boxplots showing the distribution for log transformed ASCA IgA and IgG measurements during pregnancy in women with (red) and without (blue) type 1 diabetes. The * denotes a significant difference between T1D and non-T1D women across pregnancy.

trast to *S. cerevisiae*, the difference in abundance between the groups was not significant. A possible explanation is that despite the difference in serum 1,5-AG between T1D and non-T1D women (Table 1), diabetes control in women with T1D was carefully monitored in pregnancy to minimise hyperglycaemia and prevent glycosuria. This was reflected by their median HbA1C across pregnancy of 6.33%, which is in the recommended range. In addition, we found no correlation between the concentration of serum 1,5-AG and the abundance of *S. cerevisiae*.

Sakly *et al* [53] reported higher concentrations of serum ASCA in individuals with T1D without celiac disease and suggested that this was due to increased intestinal permeability. Increased concentrations of serum ASCA have been noted in other autoimmune diseases, including antiphospholipid syn-

drome (APS) [54], systemic lupus erythematosus [55] and rheumatoid arthritis [56] in which impaired intestinal barrier function has been reported [57]. In women with T1D, intestinal inflammation was reflected by higher concentrations of faecal calprotectin and impaired epithelial integrity by higher concentrations of serum I-FABP. These are likely to be secondary to the lower abundance of *F. prausnitzii* and other producers of butyrate [58], as butyrate prevents gut inflammation and maintains the intestinal epithelial barrier [59,60]. We [11,12] and others [61] have previously reported impaired epithelial barrier function and gut permeability in T1D.

The ability of *S. cerevisiae* to produce a range of antimicrobial factors [62,63,64] may account for its reciprocal association with the abundance of *F. prausnitzii*, although why this bacterium is specifically affected is unclear and

Table 6 – Results (p-values) from likelihood ratio test comparing zero inflated negative binomial models.

Counts (Negative binomial part of the model)		Interaction	Variable	Trimester_T1D	Interpretation
Species	Ruminococcus_torques	0.817	0.761	0.878	
	Eubacterium_rectale	0.083	0.604	0.882	
	Roseburia_intestinalis	0.369	0.756	0.913	
	Anaerostipes_hadrus	0.27	0.397	0.918	
	Lachnospiraceae_bacterium_5_1_63FAA	0.851	0.795	0.869	
	Roseburia_inulinivorans	0.988	0.963	0.877	
	Faecalibacterium_prausnitzii	0.625	0.005**	0.972	For one unit increase in <i>Faecalibacterium_prausnitzii</i> the odds of having <i>S. cerevisiae</i> decrease by $1-\exp(-0.085) = 8\%$
Genus	Ruminococcus	0.588	0.17	0.697	
	Eubacterium	0.152	0.453	0.88	
	Roseburia	0.876	0.399	0.911	
	Anaerostipes	0.245	0.301	0.928	
	Faecalibacterium	0.634	0.005**	0.972	For one unit increase in <i>Faecalibacterium_prausnitzii</i> the odds of having <i>Saccharomyces</i> genus decrease by $1-\exp(-0.085) = 8\%$
Binomial (zero inflation part of the model)					
		Interaction	Variable	Trimester_T1D	
Species	Ruminococcus_torques	0.419	0.121	0.064	
	Eubacterium_rectale	0.554	0.323	0.036	
	Roseburia_intestinalis	0.321	0.377	0.037	
	Anaerostipes_hadrus	0.35	0.088	0.036	
	Lachnospiraceae_bacterium_5_1_63FAA	0.937	0.746	0.042	
	Roseburia_inulinivorans	0.696	0.599	0.049	
	Faecalibacterium_prausnitzii	0.457	0.894	0.043	
Genus	Ruminococcus	0.961	0.427	0.036	
	Eubacterium	0.433	0.279	0.062	
	Roseburia	0.353	0.087	0.033	
	Anaerostipes	0.287	0.114	0.037	
	Faecalibacterium	0.444	0.899	0.043	
***Complete results from zinb test for significant bacterial taxa associations with <i>S. cerevisiae</i>					
Call:					
zeroinfl(formula = sac ~ Faecalibacterium_prausnitzii t1dtri, data = TTMetadata, dist = "negbin")					
Count model coefficients (negbin with log link):					
		Estimate	Std. Error	z value	Pr(> z)
(Intercept		6.320	0.203	31.081	<2e-16
Faecalibacterium_prausnitzii		-0.085	0.024	-3.616	0.0003
					For one unit increase in <i>Faecalibacterium_prausnitzii</i> the odds of having <i>Saccharomyces</i> decrease by $1-\exp(-0.085) = 8\%$
Log(theta)		-0.116	0.178	-0.655	0.512
Zero-inflation model coefficients (binomial with logit link):					
		Estimate	Std. Error	z value	Pr(> z)
(Intercept)		3.35E-01	7.96E-01	0.421	0.6739
Genus	t1dtrino_T2	-1.711	0.928	-1.843	0.065

Table 6 – (continued)

t1dtrino_T3	-1.094	0.825	-1.325	0.1852	
t1dtriyes_T1	-17.970	2316.000	-0.008	0.994	
t1dtriyes_T2	-1.708	0.937	-1.823	0.068	
t1dtriyes_T3	-0.798	0.807	-0.989	0.323	
Call:					
zeroinfl(formula = sac ~ Faecalibacterium t1dtri, data = TTMetadata, dist = "negbin")					
Count model coefficients (negbin with log link):					
	Estimate	Std. Error	z value	Pr(> z)	
(Intercept)	6.319	0.203	31.066	<2e-16	
Faecalibacterium	-0.085	0.024	-3.603	0.0003	For one unit increase in <i>Faecalibacterium prausnitzii</i> the odds of having <i>Saccharomyces</i> genus decrease by $1 - \exp(-0.085) = 8\%$
Log(theta)	-0.117	0.178	-0.66	0.509	
Zero-inflation model coefficients (binomial with logit link):					
	Estimate	Std. Error	z value	Pr(> z)	
(Intercept)	0.338	0.796	0.424	0.671	
t1dtrino_T2	-1.711	0.928	-1.844	0.065	
t1dtrino_T3	-1.094	0.825	-1.326	0.185	
t1dtriyes_T1	-17.969	2315.989	-0.008	0.994	
t1dtriyes_T2	-1.709	0.937	-1.824	0.068	
t1dtriyes_T3	-0.799	0.807	-0.99	0.322	

Note Trimester_T1D status was fitted as a single term with 6 levels corresponding to the combination of the 2 levels of T1D status and the 3 trimesters.

Sac: *Saccharomyces cerevisiae* abundance

t1dtri: Refers to composed factor of T1D status and trimester

Variable: This refers to the variable being tested, as named in each row (e.g. Carbohydrates [g], Resistant starch [g], etc)

Interaction: This refers to the p-value obtained from testing if there is as significant interaction between the "variable of interest" and the term "Trimester_T1D status". If the interaction is significant, it means that the possible differences in alpha or beta diversity between T1D and non-T1D depend or are conditioned to trimester and/or T1D status (e.g. if there are differences between T1D and non-T1D only in one trimester or if there are differences between trimesters only in women with T1D).




Fig. 5 – Predicted counts of the bacterium *Faecalibacterium prausnitzii* in relation to the predicted counts of *Saccharomyces cerevisiae* in relation to non-T1D and T1D status, and trimester variables. Predicted counts: estimated marginal or adjusted means and standard errors generated based on the zero inflated negative binomial linear model used to fit the data.

highlights the need to understand the complex community interactions within the gut microbiome. The increased abundance of *Saccharomyces* in later pregnancy in women with T1D, associated with a decrease in beneficial bacteria such as *F. prausnitzii*, could be detrimental to the health of the mother and child.

In summary, the gut mycobiome changes across pregnancy in both women with and without T1D. Women with T1D exhibit a progressive increase in the abundance of *S. cerevisiae*, associated with a decrease in the abundance of the bacterium *F. prausnitzii*, together with evidence of gut inflammation and impaired epithelial integrity. The relevance of these findings to the higher rate of pregnancy complications in women with T1D and the potential to ameliorate these complications by pro-biotic or anti-fungal interventions may be worthy of further study.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgements

The authors thank Stephen A. Wilcox (Genomics Hub, WEHI) for advice on DNA sequencing. The ENDIA Study Group (May 2021) is composed of the following authors: Investigators: Simon C Barry, Maria E Craig, Peter G Colman, Jennifer J Couper, Elizabeth A Davis, Mark Harris, Leonard C Harrison, Aveni Haynes, Grant Morahan, Helena Oakey, Megan A S Penno, William D Rawlinson, Richard O Sinnott, Georgia Soldatos, Rebecca L Thomson, Peter J Vuillermin, John M Wentworth. Associate investigators: Emma Hamilton-Williams, Ki Wook Kim. Project and Data Managers: Amanda J Anderson, Pat Ashwood, James Brown, Kelly J McGorm, William Hu. Coordinators: Sarah Beresford, Samantha Bertram, Debra Bezuidenhout, Susan Brandrick, Carlie Butterworth, Jacki Catteau,

Nakita Clements, Kyana Gartrell, Alison Gwiazdzinski, Candice Hall, Gail Harper, Amanda Hulley, Mikayla Hoffman, Renee Kludas, Christine Monagle, Belinda Moore, Benjamin Ramoso, Alison Roberts, Georgina Thompson, Alexandra Tully, Isabelle Vicary, Rosemary Wood. Dietitians: Rachel Battersby, Stephanie Savio. Post-doctoral Fellows: Esther Bandala Sanchez, Naiara Bediaga, Chris Hope, Tim Sadlon, Alexandra Roth Schulze. Research Officers: Sabrina Binkowski, Bek Brittain, Minh Bui, Dylan Foskett, Dexing Huang, Dao Huynh, Stuti Kapadia, Asma Minhaj, Gaetano Naselli, Katrina Ngui, Trung Nguyen, Yeon Park, Kelly Watson, Cynthia Yau. Administrator: Leanne Cavenett.

The ENDIA Study Group would like to thank the following institutions and individuals for their contribution to ENDIA recruitment and follow-up:

Lead Clinical Recruitment/Follow-up Sites: The Women's and Children's Hospital, SA (Jennifer Couper). Royal Melbourne Hospital, Vic (Peter Colman, John Wentworth, Leonard Harrison). Barwon Health, Vic (Peter Vuillermin). Monash Health, Vic (Georgia Soldatos). Children's Hospital at Westmead, NSW (Maria Craig). Royal Hospital for Women, NSW (Maria Craig). St George Hospital, NSW (Maria Craig). Princess Margaret Hospital/Perth Children's Hospital, WA (Elizabeth Davis, Aveni Haynes). Mater Mother's Hospital/Queensland Children's Hospital, QLD (Mark Harris, Andrew Cotterill)

Lead Academic Sites: The University of Adelaide/Robinson Research Institute, SA (Jennifer Couper, Megan Penno, Rebecca Thomson, Kelly McGorm, Helena Oakey, Simon Barry). WEHI, Vic (Leonard Harrison, John Wentworth). University of New South Wales, NSW (Maria Craig, William Rawlinson). University of Sydney, NSW (Maria Craig). University of Western Australia/Telethon Kids Institute/Harry Perkins Institute, WA (Elizabeth Davis, Aveni Haynes, Grant Morahan), University of Melbourne, Vic (Richard Sinnott). University of Queensland, QLD (Mark Harris)

Satellite Recruitment/Birthing Sites (SA): Country Health SA (Jennifer Couper), Flinders Medical Centre (Brian Coppin),

Lyell McEwin Hospital (Jennifer Couper), Ashford Hospital, Flinders Private Hospital, North Eastern Community Hospital.

Satellite Recruitment/Birthing Sites (VIC): Royal Women's Hospital (Alison Nankervis), Ballarat Base Hospital (David Song), Bendigo Health (Mark Savage), Epworth Geelong Hospital (Peter Vuillermin), Mercy Hospital for Women (Christine Houlihan, Peter Colman), St. John of God Geelong (Peter Vuillermin), Sunshine Hospital/ Joan Kirner Women's and Children's Hospital (Balasubramanian Krishnamurthy), Werribee Mercy Hospital (Sheetal Tipnis)

Satellite Recruitment/Birthing Sites (NSW): Hunter Diabetes Centre (Claire Morbey), John Hunter Hospital (Maria Craig), John Hunter Children's Hospital (Maria Craig), Sydney Children's Hospital (Maria Craig), The Sutherland Hospital (Maria Craig), Westmead Hospital (Maria Craig), North Shore Private Hospital (Maria Craig)

Satellite Recruitment/Birthing Sites (WA, all under Elizabeth Davis): Armadale Hospital, Bentley Hospital, Joondalup Health Campus, King Edward Memorial Hospital, Rockingham General Hospital, St John of God Mt. Lawley, St John of God Murdoch, St John of God Subiaco.

Satellite Recruitment/Birthing Sites (QLD): Royal Brisbane and Women's Hospital (Michael d'Emden), Wesley Hospital (Stephen Cook, Andrew Cotterill)

Satellite Recruitment/Birthing Sites (NT): Royal Darwin Hospital (Louise Maple-Brown)

Referring Physicians in Private: Natalie Harrison/Geelong Diabetes & Endocrinology (Vic), Dorothy Graham (WA), Linda McKendrick (SA), Amanda Terry (SA).

Other Birthing Hospitals: Albury-Wodonga Hospital, Angliss Hospital, Auburn Public Hospital, Bankstown Hospital, Bathurst Base Hospital, Beaudesert Hospital, Beijing United Family Hospital (China), Belmont Hospital, Berri Hospital, Blacktown Hospital, Box Hill Hospital, Broome Hospital, Buderim Private Hospital, Bunbury Hospital, Burnside War Memorial Hospital, Caboolture Hospital, Cabrini Hospital Malvern, Cairns Hospital, Cairns Private Hospital, Calvary Hospital, Calvary Hospital Bruce, Calvary Hospital Lenah Valley, Calvary Hospital Wagga, Calvary John James Hospital, Campbelltown Hospital, Canberra Hospital, Casey Hospital, Centenary Hospital for Women and Children, Coffs Harbour Base Hospital, Dandenong Hospital, Darwin Birth Centre, Darwin Private Hospital, Dubbo Base Hospital, Echuca Regional Health, Emerald Hospital, Epworth Freemasons Hospital, Fiona Stanley Hospital, Frances Perry House, Frankston Hospital, Gawler Hospital, Gippsland Health, Glengarry Private Hospital, Gold Coast Private Hospital, Gold Coast University Hospital, Gosford Hospital, Gosford Private Hospital, Goulburn Valley Health, Grafton Base Hospital, Greenslopes Private Hospital, Griffith Base Hospital, Gunnedah Hospital, Hawkes Bay Hospital (New Zealand), Hervey Bay Hospital, Hobart Private Hospital, Hornsby Hospital, Hurstville Private Hospital, Ipswich Hospital, Jessie McPherson Private Hospital, John Flynn Private Hospital, Kalgoorlie Hospital, Kapunda Hospital, Kareena Private Hospital, Katoomba Hospital, Launceston General Hospital, Lismore Base Hospital, Liverpool Hospital, Logan Hospital, Mackay Base Hospital, Maitland Hospital, Manly Hospital, Mater

Hospital Sydney, Mater Mothers' Hospital, Mater Mothers' Private Hospital, Mater Private Hospital Mackay, Mater Private Hospital Redland, Mater Women's and Children's Hospital Townsville, Mitcham Private Hospital, Moruya District Hospital, Mount Barker Hospital, Narrabri Hospital, Nepean Hospital, Nepean Private Hospital, Newcastle Private Hospital, North Gosford Private Hospital, North West Hospital, North West Private Hospital, Northeast Health Wangaratta, Northern Beaches Hospital, Northern Hospital, Northpark Private Hospital, Norwest Private Hospital, Orange Base Hospital, Osborne Park Hospital, Peel Health Campus, Peninsula Private Hospital, Pindara Private Hospital, Port Augusta Hospital, Port Macquarie Hospital, Prince of Wales Hospital, Prince of Wales Private Hospital, Queanbeyan District Hospital, Queen Victoria Hospital, Redcliffe Hospital, Redland Hospital, Riverland General Hospital, Rockhampton Base Hospital, Royal Hobart Hospital, Royal Hospital for Women, Royal North Shore Hospital, Royal Prince Alfred Hospital, Sandringham Hospital, St George Private Hospital, St John of God Ballarat Hospital, St John of God Bendigo Hospital, St John of God Berwick Hospital, St John of God Bunbury Hospital, St John of God Midland Hospital, St Vincent's Hospital, St Vincent's Private Hospital, St Vincent's Private Hospital Toowoomba, Stanthorpe Hospital, Sunnybank Private Hospital, Sunshine Coast University Hospital, Swan District Hospital, Sydney Adventist Hospital, Tamworth Hospital, Tanunda Hospital, The Canberra Hospital, The Tweed Hospital, Toowoomba Base Hospital, Toowoomba Private Hospital, Townsville University Hospital, Wallaroo Hospital, Wangaratta Hospital, Waverley Private Hospital, Westmead Private Hospital, Wodonga Hospital, Wollongong Hospital, Wollongong Private Hospital.

We also gratefully acknowledge the participants and their families who contribute to the ENDIA Study.

Funding

This research was supported by JDRF Australia, the recipient of the Commonwealth of Australia grants for Accelerated Research under the Medical Research Future Fund, and with funding from the Leona M. and Harry B. Helmsley Charitable Trust (grant key 3-SRA-2020-966-M-N). In addition, support was provided by The National Health and Medical Research Council of Australia (NHMRC) Centre of Research Excellence for the Protection of Pancreatic Beta Cells (1078106), NHMRC Program Grant (LCH 1150425), and JDRF International (1-SRA-2018-543-S-B). Additional support was provided by the Victorian State Government Operational Infrastructure Support scheme, Diabetes South Australia, and the NHMRC Research Institute Infrastructure Support Scheme. MEC was supported by a NHMRC Practitioner Fellowship (1136735), and LCH by an Investigator Fellowship (1173945).

Data availability

Submission of sequencing data.

BioProject Accession ID: PRJNA756382.

<http://www.ncbi.nlm.nih.gov/bioproject/756382>.

Appendix A. Supplementary material

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.diabres.2022.109189>.

REFERENCES

- [1] Bach JF. The effect of infections on susceptibility to autoimmune and allergic diseases. *N Engl J Med* 2002;347(12):911–20. <https://doi.org/10.1056/NEJMra020100>.
- [2] King C, Sarvetnick N, von Herrath M. The incidence of type-1 diabetes in NOD mice is modulated by restricted flora not germ-free conditions. *PLoS ONE* 2011;6(2):e17049. <https://doi.org/10.1371/journal.pone.0017049>.
- [3] Mariño E, Richards JL, McLeod KH, et al. Erratum: Gut microbial metabolites limit the frequency of autoimmune T cells and protect against type 1 diabetes. *Nat Immunol* 2017;18(11):1271. <https://doi.org/10.1038/ni1117-1271c>.
- [4] Gulden E, Wong FS, Wen L. The gut microbiota and type 1 diabetes. *Clin Immunol* 2015;159(2):143–53. <https://doi.org/10.1016/j.clim.2015.05.013>.
- [5] Vatanen T, Kostic AD, d’Hennezel E, Siljander H, Franzosa EA, Yassour M, et al. Variation in Microbiome LPS Immunogenicity Contributes to Autoimmunity in Humans. *Cell* 2016;165(6):1551. <https://doi.org/10.1016/j.cell.2016.05.056>.
- [6] de Groot PF, Belzer C, Aydin Ö, et al. Distinct faecal and oral microbiota composition in human type 1 diabetes, an observational study. *PLoS One* 2017;12(12). <https://doi.org/10.1371/journal.pone.0188475>.
- [7] Stewart CJ, Ajami NJ, O’Brien JL, Hutchinson DS, Smith DP, Wong MC, et al. Temporal development of the gut microbiome in early childhood from the TEDDY study. *Nature* 2018;562(7728):583–8.
- [8] Vatanen T, Franzosa EA, Schwager R, Tripathi S, Arthur TD, Vehik K, et al. The human gut microbiome in early-onset type 1 diabetes from the TEDDY study. *Nature* 2018;562(7728):589–94. <https://doi.org/10.1038/s41586-018-0620-2>.
- [9] Higuchi BS, Rodrigues N, Gonzaga MI, et al. Intestinal dysbiosis in autoimmune diabetes is correlated with poor glycaemic control and increased interleukin-6: A Pilot Study. *Front Immunol* 2018;9:1689. <https://doi.org/10.3389/fimmu.2018.01689>.
- [10] Gavin PG, Mullaney JA, Loo D, Cao K-A, Gottlieb PA, Hill MM, et al. Intestinal metaproteomics reveals host-microbiota interactions in subjects at risk for type 1 Diabetes. *Diabetes Care* 2018;41(10):2178–86. <https://doi.org/10.2337/dc18-0777>.
- [11] Harbison JE, Roth-Schulze AJ, Giles LC, Tran CD, Ngui KM, Penno MA, et al. Gut microbiome dysbiosis and increased intestinal permeability in children with islet autoimmunity and type 1 diabetes: A prospective cohort study. *Pediatr Diabetes* 2019. <https://doi.org/10.1111/pedi.12865>.
- [12] Roth-Schulze AJ, Penno MAS, Ngui KM, Oakey H, Bandala-Sanchez E, Smith AD, et al. Type 1 diabetes in pregnancy is associated with distinct changes in the composition and function of the gut microbiome. *Microbiome* 2021;9(1). <https://doi.org/10.1186/s40168-021-01104-y10.21203/rs.3.rs-967264/v1>.
- [13] Qin J, Li R, Raes J, Arumugam M, Burgdorf KS, Manichanh C, et al. A human gut microbial gene catalogue established by metagenomic sequencing. *Nature* 2010;464(7285):59–65. <https://doi.org/10.1038/nature08821>.
- [14] Huseyin CE, O’Toole PW, Cotter PD, Scanlan PD. Forgotten fungi—the gut mycobiome in human health and disease. *FEMS Microbiol Rev* 2017;41(4):479–511. <https://doi.org/10.1093/femsre/fuw047>.
- [15] Hallen-Adams HE, Suhr MJ. Fungi in the healthy human gastrointestinal tract. *Virulence* 2017;8(3):352–8. <https://doi.org/10.1080/21505594.2016.1247140>.
- [16] Richard ML, Sokol H. The gut mycobiota: insights into analysis, environmental interactions, and role in gastrointestinal diseases. *Nat Rev Gastroenterol Hepatol* 2019;16(6):331–45. <https://doi.org/10.1038/s41575-019-0121-2>.
- [17] Nash AK, Auchtung TA, Wong MC, Smith DP, Gesell JR, Ross MC, et al. The gut mycobiome of the Human Microbiome Project healthy cohort. *Microbiome* 2017;5(1). <https://doi.org/10.1186/s40168-017-0373-4>.
- [18] Gosiewski T, Salamon D, Szopa M, Sroka A, Malecki MT, Bulanda M. Quantitative evaluation of fungi of the genus *Candida* in the faeces of adult patients with type 1 and 2 diabetes - a pilot study. *Gut Pathog* 2014;6(1):43. <https://doi.org/10.1186/s13099-014-0043>.
- [19] Soyucen E, Gulcan A, Aktuglu-Zeybek AC, Onal H, Kiykim E, Aydin A. Differences in the gut microbiota of healthy children and those with type 1 diabetes. *Pediatr Int* 2014;56(3):336–43. <https://doi.org/10.1111/ped.12243>.
- [20] Kowalewska B, Zorena K, Szmigiero-Kawko M, Waż P, Myśliwiec M. Higher diversity in fungal species discriminates children with type 1 diabetes mellitus from healthy control. *Patient Prefer Adherence* 2016;21(10):591–9. <https://doi.org/10.2147/PPA.S97852>.
- [21] Penno MAS, Couper JJ, Craig ME, Colman PG, Rawlinson WD, Cotterill AM, et al. Environmental determinants of islet autoimmunity (ENDIA): a pregnancy to early life cohort study in children at-risk of type 1 diabetes. *BMC Pediatr* 2013;13(1). <https://doi.org/10.1186/1471-2431-13-124>.
- [22] Penington JS, Penno MAS, Ngui KM, et al. Influence of faecal collection conditions and 16S rRNA gene sequencing at two centers on human gut microbiota analysis. *Sci Rep* 2018;8(1):4386. <https://doi.org/10.1038/s41598-018-22491-7>.
- [23] Bolyen E, Rideout JR, Dillon MR, et al. Reproducible, interactive, scalable, and extensible microbiome data science using QIIME 2 [published correction appears in *Nat Biotechnol*. 2019 Sep; 37(9): 1091]. *Nat Biotechnol* 2019;37(8):852–7. <https://doi.org/10.1038/s41587-019-0252-6>.
- [24] Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJ, Holmes SP. DADA2: High-resolution sample inference from Illumina amplicon data. *Nat Methods* 2016;13(7):581–3. <https://doi.org/10.1038/nmeth.3869>.
- [25] Nilsson RH, Larsson KH, Taylor AFS, et al. The UNITE database for molecular identification of fungi: handling dark taxa and parallel taxonomic classifications. *Nucleic Acids Res* 2019;47(D1):D259–64. <https://doi.org/10.1093/nar/gky1022>.
- [26] McMurdie PJ, Holmes S. phyloseq: An R package for reproducible interactive analysis and graphics of microbiome census data. *PLoS One* 2013;8(4). <https://doi.org/10.1371/journal.pone.0061217>.
- [27] Team RC. R: A language and environment for statistical computing; 2015. 2018.
- [28] Hodge A, Patterson AJ, Brown WJ, Ireland P, Giles G. The Anti-Cancer Council of Victoria FFQ: relative validity of nutrient intakes compared with weighed food records in young to middle-aged women in a study of iron supplementation [published correction appears in *Aust N Z J Public Health*. 2003; 27(4): 468]. *Aust N Z J Public Health* 2000;24(6):576–83. <https://doi.org/10.1111/j.1467-842x.2000.tb00520.x>.
- [29] Nowak N, Skupien J, Cyganek K, Matejko B, Malecki MT. 1,5-Anhydroglucitol as a marker of maternal glycaemic control and predictor of neonatal birthweight in pregnancies complicated by type 1 diabetes mellitus. *Diabetologia* 2013;56(4):709–13. <https://doi.org/10.1007/s00125-013-2830-3>.

- [30] Tibble JA, Bjarnason I. Faecal calprotectin as an index of intestinal inflammation. *Drugs Today (Barc)* 2001;37(2):85–96. <https://doi.org/10.1358/dot.2001.37.2.614846>.
- [31] Røseth AG, Fagerhol MK, Aadland E, Schjønby H. Assessment of the neutrophil dominating protein calprotectin in faeces. A methodologic study. *Scand J Gastroenterol* 1992;27(9):793–8. <https://doi.org/10.3109/00365529209011186>.
- [32] John E, Kronborg O, Tøn HI, Kristinsson J, Fuglerud P. A new faecal calprotectin test for colorectal neoplasia. *Clinical results and comparison with previous method. Scand J Gastroenterol* 2001;36(3):291–6.
- [33] Thuijls G, Derikx JPM, van Wijck K, Zimmermann LJI, Degraeuwe PL, Mulder TL, et al. Non-invasive markers for early diagnosis and determination of the severity of necrotizing enterocolitis. *Ann Surg* 2010;251(6):1174–80. <https://doi.org/10.1097/SLA.0b013e3181d778c4>.
- [34] Vreugdenhil AC, Wolters VM, Adriaanse MP, Van den Neucker AM, van Bijnen AA, Houwen R, et al. Additional value of serum I-FABP levels for evaluating celiac disease activity in children. *Scand J Gastroenterol* 2011;46(12):1435–41. <https://doi.org/10.3109/00365521.2011.627447>.
- [35] Hardin JW. Generalized estimating equations (GEE). *Encycl Stat Behav Sci: American Cancer Society*; 2005.
- [36] Halekoh U, Højsgaard S, Yan J. The R package geepack for generalized estimating equations. *J Stat Softw* 2006;15:1–11.
- [37] Oksanen J, Blanchet FG, Friendly M, Kindt R, Legendre P, McGlenn D, et al. vegan: community ecology package. *R package version 2.5-2*; 2018.
- [38] Lloyd-Price J, Arze C, Ananthakrishnan AN, Schirmer M, Avila-Pacheco J, Poon TW, et al. Multiomics of the gut microbial ecosystem in inflammatory bowel diseases. *Nature* 2019;569(7758):655–62. <https://doi.org/10.1038/s41586-019-1237-9>.
- [39] Brooks M, Kristensen K, Benthem K, Magnusson A, Berg C, Nielsen A, et al. glmmTMB balances speed and flexibility among packages for zero-inflated generalized linear mixed modeling. *R J* 2017;9(2):378. <https://doi.org/10.32614/RJ-2017-066>.
- [40] Lenth R. Emmeans package: Estimated marginal means, aka least-squares means. *R package version 2019;1(3):5.1*.
- [41] Zeileis A, Kleiber C, Jackman S. Regression Models for Count Data in R. *J Stat Softw* 2008;27(8).
- [42] Zeileis A, Hothorn T. Diagnostic Checking in Regression Relationships. *R News* 2002;2(3):7–10.
- [43] Berstad A, Arslan G, Folvik G. Relationship between intestinal permeability and calprotectin concentration in gut lavage fluid. *Scand J Gastroenterol* 2000;35(1):64–9. <https://doi.org/10.1080/003655200750024551>.
- [44] Arumugam M, Raes J, Pelletier E, Le Paslier D, Yamada T, Mende DR, et al. Enterotypes of the human gut microbiome. *Nature* 2011;473(7346):174–80.
- [45] Bhute SS, Suryavanshi MV, Joshi SM, Yajnik CS, Shouche YS, Ghaskadbi SS. Gut microbial diversity assessment of indian type-2-diabetics reveals alterations in eubacteria, archaea, and eukaryotes. *Front Microbiol* 2017;8:214. <https://doi.org/10.3389/fmicb.2017.00214>.
- [46] Hamad I, Ranque S, Azhar EI, Yasir M, Jiman-Fatani AA, Tissot-Dupont H, et al. Culturomics and amplicon-based metagenomic approaches for the study of fungal population in human gut microbiota. *Sci Rep* 2017;7(1). <https://doi.org/10.1038/s41598-017-17132-4>.
- [47] Xu J, Boyd CM, Livingston E, Meyer W, Madden JF, Mitchell TG. Species and genotypic diversities and similarities of pathogenic yeasts colonizing women. *J Clin Microbiol* 1999;37(12):3835–43. <https://doi.org/10.1128/JCM.37.12.3835-3843.1999>.
- [48] Ramazzotti M, Stefanini I, Di Paola M, De Filippo C, Rizzetto L, Berná L, et al. Population genomics reveals evolution and variation of *Saccharomyces cerevisiae* in the human and insects' gut. *Environ Microbiol* 2019;21(1):50–71. <https://doi.org/10.1111/1462-2920.14422>.
- [49] Angebault C, Djossou F, Abélanet S, et al. *Candida albicans* is not always the preferential yeast colonizing humans: a study in Wayampi Amerindians. *J Infect Dis* 2013;208(10):1705–16. <https://doi.org/10.1093/infdis/jit389>.
- [50] Hoffmann C, Dollive S, Grunberg S, et al. Archaea and fungi of the human gut microbiome: correlations with diet and bacterial residents. *PLoS One* 2013;8(6). <https://doi.org/10.1371/journal.pone.0066019>.
- [51] Nowakowska D, Kurnatowska A, Stray-Pedersen B, Wilczyński J. Species distribution and influence of glycaemic control on fungal infections in pregnant women with diabetes. *J Infect* 2004;48(4):339–46. <https://doi.org/10.1016/j.jinf.2004.01.017>.
- [52] Rodrigues CF, Rodrigues ME, Henriques M. *Candida* sp. infections in patients with diabetes mellitus. *J Clin Med* 2019;8(1):76. <https://doi.org/10.3390/jcm8010076>.
- [53] Sakly W, Mankai A, Sakly N, Thabet Y, Achour A, Ghedira-Besbes L, et al. Anti-*Saccharomyces cerevisiae* antibodies are frequent in type 1 diabetes. *Endocr Pathol* 2010;21(2):108–14. <https://doi.org/10.1007/s12022-010-9118-7>.
- [54] Krause I, Blank M, Cervera R, Font J, Matthias T, Pfeiffer S, et al. Cross-reactive epitopes on beta2-glycoprotein-I and *Saccharomyces cerevisiae* in patients with the antiphospholipid syndrome. *Ann N Y Acad Sci* 2007;1108(1):481–8. <https://doi.org/10.1196/annals.1422.051>.
- [55] Dai H, Li Z, Zhang Y, Lv P, Gao XM. Elevated levels of serum antibodies against *Saccharomyces cerevisiae* mannan in patients with systemic lupus erythematosus. *Lupus* 2009;18(12):1087–90. <https://doi.org/10.1177/0961203309105131>.
- [56] Dai H, Li Z, Zhang Y, Lv P, Gao XM. Elevated levels of serum IgA against *Saccharomyces cerevisiae* mannan in patients with rheumatoid arthritis. *Cell Mol Immunol* 2009;6(5):361–6. <https://doi.org/10.1038/cmi.2009.47>.
- [57] Mu Q, Kirby J, Reilly CM, Luo XM. Leaky gut as a danger signal for autoimmune diseases. *Front Immunol* 2017;8:598. <https://doi.org/10.3389/fimmu.2017.00598>.
- [58] Louis P, Flint HJ. Diversity, metabolism, and microbial ecology of butyrate-producing bacteria from the human large intestine. *FEMS Microbiol Lett* 2009;294(1):1–8. <https://doi.org/10.1111/j.1574-6968.2009.01514.x>.
- [59] Koh A, De Vadder F, Kovatcheva-Datchary P, Bäckhed F. From dietary fibre to host physiology: short-chain fatty acids as key bacterial metabolites. *Cell* 2016;165(6):1332–45. <https://doi.org/10.1016/j.cell.2016.05.041>.
- [60] Zheng L, Kelly CJ, Battista KD, Schaefer R, Lanis JM, Alexeev EE, et al. Microbial-derived butyrate promotes epithelial barrier function through IL-10 receptor-dependent repression of claudin-2. *J Immunol* 2017;199(8):2976–84. <https://doi.org/10.4049/jimmunol.1700105>.
- [61] Leiva-Gea I, Sánchez-Alcoholado L, Martín-Tejedor B, Castellano-Castillo D, Moreno-Indias I, Urda-Cardona A, et al. Gut microbiota differs in composition and functionality between children with type 1 diabetes and MODY2 and healthy control subjects: a case-control study. *Diabetes Care* 2018;41(11):2385–95. <https://doi.org/10.2337/dc18-0253>.
- [62] Magliani W, Conti S, Gerloni M, Bertolotti D, Polonelli L. Yeast killer systems. *Clin Microbiol Rev* 1997;10(3):369–400. <https://doi.org/10.1128/CMR.10.3.369>.
- [63] Melvydas V, Bružauskaitė I, Gedminienė G, Šiekštelė R. A novel *Saccharomyces cerevisiae* killer strain secreting the X factor related to killer activity and inhibition of *S. cerevisiae* K1, K2 and K28 killer toxins. *Indian. J Microbiol* 2016;56(3):335–43. <https://doi.org/10.1007/s12088-016-0589-1>.

-
- [64] Fakruddin M, Hossain MN, Ahmed MM. Antimicrobial, and antioxidant activities of *Saccharomyces cerevisiae* IFST062013, a potential probiotic. *BMC Complement Altern Med* 2017;17(1):64. <https://doi.org/10.1186/s12906-017-1591-9>.
- [65] Chasan-taber L, Schmidt MD, Roberts DE, Hosmer D, Markenson G, Freedson PS. Development and Validation of a Pregnancy Physical Activity Questionnaire [published correction appears in *Med Sci Sports Exerc.* 2011; 43(1):195]. *Med Sci Sports Exerc* 2004;36(10):1750–60.