



Attitudes and barriers to incident reporting: a collaborative hospital study

S M Evans, J G Berry, B J Smith, A Esterman, P Selim, J O'Shaughnessy and M DeWit

Qual. Saf. Health Care 2006;15:39-43
doi:10.1136/qshc.2004.012559

Updated information and services can be found at:

<http://qshc.bmj.com/cgi/content/full/15/1/39>

These include:

References

This article cites 22 articles, 7 of which can be accessed free at:

<http://qshc.bmj.com/cgi/content/full/15/1/39#BIBL>

9 online articles that cite this article can be accessed at:

<http://qshc.bmj.com/cgi/content/full/15/1/39#otherarticles>

Rapid responses

You can respond to this article at:

<http://qshc.bmj.com/cgi/eletter-submit/15/1/39>

Email alerting service

Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article

Notes

To order reprints of this article go to:

<http://journals.bmj.com/cgi/reprintform>

To subscribe to *Quality and Safety in Health Care* go to:

<http://journals.bmj.com/subscriptions/>

ORIGINAL ARTICLE

Attitudes and barriers to incident reporting: a collaborative hospital study

S M Evans, J G Berry, B J Smith, A Esterman, P Selim, J O'Shaughnessy, M DeWit

Qual Saf Health Care 2006;15:39–43. doi: 10.1136/qshc.2004.012559

See end of article for authors' affiliations

Correspondence to:
Ms S Evans, Department of
Epidemiology and
Preventive Medicine,
Monash University, Alfred
Hospital, Prahran,
Victoria, Australia;
sue.evans@med.monash.edu.au

Accepted for publication
22 November 2005

Objectives: To assess awareness and use of the current incident reporting system and to identify factors inhibiting reporting of incidents in hospitals.

Design, setting and participants: Anonymous survey of 186 doctors and 587 nurses from diverse clinical settings in six South Australian hospitals (response rate=70.7% and 73.6%, respectively).

Main outcome measures: Knowledge and use of the current reporting system; barriers to incident reporting.

Results: Most doctors and nurses (98.3%) were aware that their hospital had an incident reporting system. Nurses were more likely than doctors to know how to access a report (88.3% v 43.0%; relative risk (RR) 2.05, 95% CI 1.61 to 2.63), to have ever completed a report (89.2% v 64.4%; RR 1.38, 95% CI 1.19 to 1.61), and to know what to do with the completed report (81.9% v 49.7%; RR 1.65, 95% CI 1.27 to 2.13). Staff were more likely to report incidents which are habitually reported, often witnessed, and usually associated with immediate outcomes such as patient falls and medication errors requiring corrective treatment. Near misses and incidents which occur over time such as pressure ulcers and DVT due to inadequate prophylaxis were least likely to be reported. The most frequently stated barrier to reporting for doctors and nurses was lack of feedback (57.7% and 61.8% agreeing, respectively).

Conclusions: Both doctors and nurses believe they should report most incidents, but nurses do so more frequently than doctors. To improve incident reporting, especially among doctors, clarification is needed of which incidents should be reported, the process needs to be simplified, and feedback given to reporters.

It is a decade since Leape¹ highlighted the need to gather information and redesign hospital systems to minimise errors in health care. Since then, many strategies and tools have been developed to identify and reduce errors.² More than 90% of consumers believe that healthcare workers should report errors,³ and peak quality and safety organisations^{4–6} recommend incident reporting to better understand errors and their contributing factors.

Incidents occurring infrequently, or those not easily coded through limitations in the existing classification system,⁷ can miss detection through medical record review. Incident reporting captures more contextual information about incidents⁸ and, when actively promoted within the clinical setting, it can detect more preventable adverse events than medical record review⁹ at a fraction of the cost.¹⁰ Near misses are rarely documented in medical records,¹¹ yet occur more frequently than adverse events¹² and provide valuable lessons in recovery mechanisms without the detrimental consequences of an adverse event.^{5–6} The subjective data provided by incident reporting enable hypothesis building and preventative strategies to be developed and tested.

Despite its strengths, many incidents are not reported probably for the same reasons they are omitted from medical records; they are simply not recognised, and those that are detected after the event are often not dealt with effectively.¹³ Outside the discipline of anaesthesiology, incident reporting is used predominantly by nurses.^{14–15} The subjective nature of reports, the lack of consistency and validation of incident data classification, and underreporting constrain incident reporting from being used as a reliable epidemiological tool to measure the frequency of events and whether interventions are effective in improving patient safety.^{2–14} Studies which have successfully improved incident reporting have often done so through intense facilitation, either through ward

rounds⁹ or staff reminders,^{10–16} and have questionable sustainability.

For incident reporting to be more reliable, both doctors and nurses must provide a representative account of errors occurring in hospitals. The objectives of this study were therefore to investigate by profession: (1) awareness and use of the current incident reporting system; (2) the types of incidents staff were more likely to report and believe should be reported; and (3) the barriers to reporting.

METHODS

Study design

A cross sectional survey of doctors and nurses was undertaken between November 2001 and June 2003. Hospitals sampled included three principal referral hospitals (each with >300 acute inpatient beds), one major referral hospital (<200 acute inpatient beds), and two major rural base hospitals (each with <120 acute inpatient beds) in South Australia. Rostered doctors and nurses, and casual agency nurses working in one or more of the four intensive care units (two metropolitan and two rural), four emergency departments (two metropolitan and two rural), five surgical units (three metropolitan and two rural), and seven medical units (five metropolitan and two rural) were invited to participate.

Project officers either personally distributed questionnaires to rostered staff and outlined the purpose of the study or, where this was not possible, posted the questionnaire. All doctors were contacted by telephone to encourage participation. To facilitate frank comment without fear of disclosure, the questionnaire was anonymous and self-administered. Ethics committee approval was obtained from each hospital's relevant body.

Questionnaire

The questionnaire was modified from one used in an obstetrics unit by Vincent *et al*¹⁷ to make it generalisable to a wider population. Following review by a panel of clinicians to assess content validity, the questionnaire was piloted on 14 doctors and 10 nurses. Test-retest reliability was determined using a kappa statistic, and only questions for which there was at least moderate reproducibility (kappa ≥ 0.4) or a consistent endorsement of one option were included.

Content

Staff were asked if they knew whether their hospital had an incident reporting system. Those answering in the affirmative were asked whether they knew which form to use, how to access it, and what to do with a completed form.

To measure reporting practice, staff were asked to estimate how often they reported 11 patient incidents representing a diverse range of common iatrogenic injuries,¹⁸ and how often they believed each should be reported using a 4-point Likert scale (never, <50% of occasions, $\geq 50\%$ of occasions, always).

To determine barriers to reporting, staff were provided with 19 potential reasons for not reporting incidents and asked to rate on a 5-point Likert scale (1 = strongly agree, 5 = strongly disagree) the degree to which these acted as a deterrent.

Analysis of data

Comparisons were made for doctors and nurses by profession, level of qualification, years post entry level qualification spent in the acute health sector, and rural/metropolitan location. For knowledge and use of reporting systems and reporting practices, log binomial generalised linear models adjusting for clustering by hospital were used. Likert scales were dichotomised into agree or not agree for reporting barriers and analysed using Fisher's exact test. The conventional level of $p \leq 0.05$ was taken to represent statistical significance. Concordance between views on current reporting behaviour and necessity of reporting was determined using an intraclass correlation coefficient (ICC). Data were analysed using Stata statistical software Version 7.0 (Stata Corporation, College Station, TX, USA).

RESULTS

The overall response rate was 72.8%, and was similar for both doctors and nurses (fig 1). As the questionnaire was anonymous, we were unable to ascertain the demographic features of non-respondents.

Knowledge and use of the incident reporting system

Nurses had a greater awareness of and used the incident reporting system more than doctors (table 1). Senior doctors

(registrars and consultants) were significantly less likely than junior doctors (interns and residents) to have ever completed an incident form (58.4% *v* 85.4%; relative risk (RR) 0.58, 95% CI 0.46 to 0.73). Doctors with >5 years experience post entry level were less likely to have ever completed an incident report than those with less experience (58.1% *v* 79.2%; RR 0.73, 95% CI 0.59 to 0.92). There were no significant differences between rural and metropolitan doctors in knowledge or use of the incident reporting system.

Senior nurses (nurse managers and clinical nurses) were more likely than junior nurses to know how to access a form (100.0% *v* 88.0%; RR 1.14, 95% CI 1.09 to 1.18), to know what to do with it (100.0% *v* 80.9%; RR 1.24, 95% CI 1.13 to 1.35), and to have ever filled one out (100.0% *v* 89.0%; RR 1.12, 95% CI 1.10 to 1.15). Permanently employed nurses were significantly more likely than contract nurses to know how to locate/access an incident form (89.1% *v* 57.1%; RR 1.56, 95% CI 1.25 to 1.95), to know what to do with it once completed (82.7% *v* 50.0%; RR 1.65, 95% CI 1.12 to 2.45), and to have ever filled one out (90.0% *v* 57.1%; RR 1.57, 95% CI 1.21 to 2.06). Nurses with >5 years post entry level experience were more likely to know how to locate a form (91.5% *v* 83.8%; RR 1.09, 95% CI 1.04 to 1.14), to know what to do with it once completed (85.5% *v* 76.7%; RR 1.12, 95% CI 1.06 to 1.17), and to have ever filled out an incident form (94.0% *v* 82.5%; RR 1.14, 95% CI 1.06 to 1.23) than those with less experience. Rural and metropolitan nurses did not differ in their knowledge or use of the incident reporting system.

Staff reporting practices

Figure 2 shows, for 11 patient incidents, the percentages of doctors and nurses who perceive they report the incident always, on 50% or more of occasions, less than 50% of occasions, or never, and their views on the necessity of reporting these incidents.

Doctors reported that they completed incident reports most often for patient falls and least often for pressure sores. Doctors' views ranged from 75.8% who believed that patient falls should always be reported to only 42.1% for drug error "near misses". Agreement between what doctors did compared with what they thought they should report was low, ranging from an ICC of 0.44 for incidents where a patient received the wrong treatment or procedure to an ICC of 0.17 for pressure sores. Senior doctors were less likely than junior doctors to always report patient falls (38.1% *v* 74.4%; RR 0.51, 95% CI 0.30 to 0.87) and patients receiving the wrong treatment or procedure (39.5% *v* 54.1%; RR 0.73, 95% CI 0.63 to 0.84). There were no significant differences in reporting practices among doctors according to rural/metropolitan location.

Nurses reported that they completed incident reports most often for patient falls and least often for pressure sores. Nurses regarded falls as the most important incidents to always report and drug error "near misses" as the least important (97.0% and 41.9%, respectively). The correlation between what nurses did compared with what they thought they should report ranged from an ICC of 0.78 for patient falls to an ICC of 0.27 for deep vein thrombosis (DVT) through inadequate prophylaxis. Nurses with <5 years experience were more likely to always report DVT (23.0% *v* 14.4%; RR 1.60, 95% CI 1.46 to 1.74). There were no significant differences in reporting practices among nurses according to rural/metropolitan location.

Staff views on barriers to reporting

Table 2 shows that major barriers to reporting for doctors were lack of feedback (57.7%), the incident form taking too long to complete (54.2%), and a belief that the incident was too trivial (51.2%). There were no significant differences for




Figure 1 Sampling frame.

Table 1 Awareness and use of the incident reporting system

	Doctors (%)		Nurses (%)		p value*	Relative risk	95% CI
	Yes	N	Yes	N			
Awareness of hospital incident reporting system	93.6%	174	99.8%	586	0.195	1.01	0.99 to 1.03
Ever completed an incident report	64.6%	115	89.2%	520	<0.001	1.38	1.19 to 1.61
Know how to locate/access an incident form	43.0%	77	88.3%	515	<0.001	2.05	1.61 to 2.63
Know what to do with a completed incident form	49.7%	89	81.9%	476	<0.001	1.65	1.27 to 2.13

*Log binomial generalised linear models adjusting for clustering by hospital.

any barriers according to level of qualification, experience, and rural/metropolitan location.

Major barriers to reporting for nurses were lack of feedback (61.8%), a belief that there was no point in reporting near misses (49.0%), and forgetting to make a report when the ward is busy (48.1%, table 2). Nurses with >5 years experience were more likely to believe there was no point reporting near misses (52.5% v 44.0%; RR 1.19, 95% CI 1.06 to 1.34) than nurses with less experience. There were no significant differences for any barriers according to level of qualification and rural/metropolitan location.

DISCUSSION

With adverse event rates estimated to be in the range of 2.9%¹⁹ to 16.6%¹⁸ of acute care hospital admissions, most doctors and nurses working in hospitals will be familiar with a range of adverse events. Despite most staff knowing that an incident reporting system existed, almost a quarter of staff did not know how to access an incident form or what to do with it once completed, and over 40% of consultants and registrars had never completed a report. Nurses were more aware of the reporting system than doctors, although casually

employed nurses were significantly less likely than permanent hospital nurses to know how to access a report, and were a third less likely to have ever completed a report.

At the time of the survey the AIMS reporting system had been used in each hospital for at least 5 years. The reporting system, which offers statutory immunity to reporters, collects data which are entered retrospectively into a stand alone database within each hospital and is managed by only authorised hospital personnel.

In most cases reporting practices were consistent with staff views on the necessity of reporting incidents. Incidents which are immediate, often witnessed, and habitually reported (such as patient falls and medication errors requiring corrective treatment) were better reported than incidents which occurred gradually and were often not attributable to a single event, or were commonly regarded as complications of prolonged hospitalisation (such as pressure ulcers, hospital acquired infections, and postoperative DVT due to inadequate prophylaxis). Only 42.0% of the staff surveyed believed that medication near misses should always be reported, indicating that literature emphasising the importance of reporting near misses¹² is not translating to changes in attitude or clinical




Figure 2 Staff self-perception of reporting of incidents.

Table 2 Self-perceived barriers to reporting (percentage who agree with the statement)

Incident	Doctors (%)		Nurses (%)		p value* (doctors v nurses)
	Agree	N	Agree	N	
I never get any feedback on what action is taken	57.7	170	61.8	570	0.371
The incident form takes too long to fill out and I just don't have the time	54.2	168	44.1	571	0.022
The incident was too trivial	51.2	170	41.2	565	0.027
When the ward is busy I forget to make a report	47.3	167	48.1	574	0.930
I don't know whose responsibility it is to make a report	37.9	169	10.8	573	<0.001
When it is a near miss, I don't see any point in reporting it	36.0	172	49.0	569	0.003
The AIMS+ form is too complicated and requires too much detail	31.9	163	35.0	565	0.512
Junior staff are often blamed unfairly for adverse incidents	31.0	171	25.6	571	0.169
Adverse incident reporting is unlikely to lead to system changes	28.6	171	29.9	568	0.775
I wonder about who else is privy to the information that I disclose	27.1	170	33.8	568	0.112
If I discuss the case with the person involved nothing else needs to be done	24.9	169	11.5	566	<0.001
I don't feel confident the form is kept anonymous	22.6	168	30.0	574	0.065
I am worried about litigation	20.7	169	20.6	574	1.000
It's not my responsibility to report somebody else's mistakes	17.2	169	16.4	567	0.814
My co-workers may be unsupportive	13.8	167	20.8	573	0.045
I don't want to get into trouble	10.6	169	18.6	570	0.014
Even if I don't give my details, I'm sure that they'll track me down	8.4	167	17.0	564	0.006
I am worried about disciplinary action	8.3	168	18.1	570	0.002
I don't want the case discussed in meetings	7.2	167	15.5	574	0.005

*Fisher's exact test.

behaviour. The finding that 80.9% of doctors thought they should always report when a patient gets the wrong treatment—yet only 57.3% believed they should always report when a patient does not receive necessary treatment—is important, given that acts of omission have been implicated in twice as many adverse events as acts of commission.¹⁸

Almost two thirds of respondents believed lack of feedback was the greatest deterrent to reporting. Organisational factors relating to structures and processes for reporting,²⁰ such as inadequate feedback, long forms and insufficient time to report, were identified as the major barriers.

Many of our results support those obtained internationally, including the finding that only a small percentage of doctors formally report incidents,^{20, 21} and unfamiliarity with the reporting process results in a poorer reporting culture.²² Whereas other studies identified cultural issues such as fear of disciplinary action,^{17, 23} legal ramifications, and workplace discrimination²⁴ as barriers to reporting, our study, like that of Uribe *et al*,²⁰ did not identify these issues to be major reporting obstacles. Poor reporting practices by doctors and the fact that they did not identify cultural barriers so much as organisational barriers to reporting probably reflects the prevailing deeply entrenched belief in medicine that only bad doctors make mistakes.

There were a number of limitations to this study. This survey formed baseline data for a matched controlled study in which purposive sampling was undertaken to reduce contamination between intervention and control units and to ensure a variety of areas were represented. Despite non-probability sampling being less ideal than random selection,²⁵ our findings were similar to those determined by Vincent *et al*¹⁷ in a distant healthcare setting, which suggests that the results are representative. Non-responder bias cannot be excluded as we were unable to collect information on non-responders due to the anonymous design of the survey. There may be potentially important variables and barriers not included in the questionnaire because we needed to limit questionnaire burden. Despite being anonymous, respondents may have provided more socially acceptable responses for fear of identification, which might explain why cultural barriers were not reported as significant deterrents to reporting. We did not investigate why staff reported certain incidents more frequently than others. Perhaps staff did not view them as incidents, or believed tools exist to detect/

monitor them or that, in the case of senior medical staff, they delegate reporting to junior staff.

Further research is required to explore why senior medical staff do not support reporting and why iatrogenic injuries with potentially disastrous consequences such as DVT and hospital acquired infections are poorly reported. Our data suggest that the move towards more casual nurses²⁶ could result in a further decline in the number and types of reports submitted, which requires action if incident reporting is to be valued as an important component of each hospital's risk management framework.

Balancing the requirement to receive adequate information on an incident report to enable meaningful analysis and follow up with the clinician's desire to make it less time consuming is an ongoing concern. Faster reporting systems, combined with adequate resources and infrastructure to enable responsive action and feedback, need to be adopted. The use of personal digital assistants,²⁷ call centres to collect information,²⁸ and techniques such as root cause analysis²⁹ to investigate incidents offers possibilities to enable safer health care to be delivered. Perhaps the most challenging task is ensuring that practice improvements resulting from reports are disseminated to clinicians, because only then will incident reporting be seen as worthwhile and relevant.

ACKNOWLEDGEMENTS

The authors thank Rhonda Bills, Clinical Epidemiology Unit for administrative support; Dr Deborah Turnbull, Department of General Practice, University of Adelaide and Lora DalGrande, Centre for Population Studies in Epidemiology, SA Department of Human Services for valuable advice on questionnaire construct and format; and the staff of the participating hospitals for completing the questionnaire.

Authors' affiliations

S M Evans, Department of Medicine, University of Adelaide, South Australia, 5005

J G Berry, Research Centre for Injury Studies, Flinders University, South Australia, 5001

B J Smith, P Selim, J O'Shaughnessy, M DeWit, Clinical Epidemiology and Health Outcomes Unit, The Queen Elizabeth Hospital, South Australia, 5011

A Esterman, School of Nursing and Midwifery, University of South Australia, South Australia, 5000

REFERENCES

- 1 **Leape LL**. Error in medicine. *JAMA* 1994;**272**:1851–7.
- 2 **Shojania KG**, Duncan BW, McDonald KM, *et al*. Making health care safer: a critical analysis of patient safety practices. *Evid Rep Technol Assess (Summ)* 2001;**43**:1–668.
- 3 **Evans SM**, Berry JG, Smith BJ, *et al*. Anonymity or transparency in reporting of medical error: a community-based survey in South Australia. *Med J Aust* 2004;**180**:577–80.
- 4 **Australian Council for Safety and Quality in Health Care**. *Safety in numbers. A technical options paper for a national approach to the use of data for safer health care*. Canberra: Commonwealth of Australia, 2001.
- 5 **Department of Health**. *An organisation with a memory*. London: Stationery Office, 2000.
- 6 **Kohn LT**, Corrigan JM, Donaldson MS. *To err is human: building a safer health system*. Washington, DC: National Academy Press, 2000:1–16.
- 7 **Runciman WB**, Moller J. *Latrogenic injury in Australia*. A report prepared by the Australian Patient Safety Foundation for the National Health Priorities and Quality Branch of the Department of Health and Aged Care of the Commonwealth Government of Australia. Adelaide, South Australia: Australian Patient Safety Foundation, 2001.
- 8 **Runciman WB**, Merry A. A tragic death: a time to blame or a time to learn? *Qual Saf Health Care* 2003;**12**:321–2.
- 9 **Beckmann U**, Bohringer C, Carless R, *et al*. Evaluation of two methods for quality improvement in intensive care: facilitated incident monitoring and retrospective medical chart review. *Crit Care Med* 2003;**31**:1006–11.
- 10 **O'Neil AC**, Petersen LA, Cook EF, *et al*. Physician reporting compared with medical record review to identify adverse medical events. *Ann Intern Med* 1993;**119**:370–6.
- 11 **Neale G**, Woloshynowych M. Retrospective case record review: a blunt instrument that needs sharpening. *Qual Saf Health Care* 2003;**12**:2–3.
- 12 **Barach P**, Small SD. Reporting and preventing medical mishaps: lessons from non-medical near miss reporting systems. *BMJ* 2000;**320**:759–63.
- 13 **Cullen DJ**, Bates DW, Small SD, *et al*. The incident reporting system does not detect adverse drug events: a problem for quality improvement. *Jt Comm J Qual Improv* 1995;**21**:541–8.
- 14 **Johnson CW**. How will we get the data and what will we do with it then? Issues in the reporting of adverse healthcare events. *Qual Saf Health Care* 2003;**12**(Suppl II):ii64–7.
- 15 **Kingston MJ**, Evans SM, Smith BJ, *et al*. Attitudes of doctors and nurses towards incident reporting: a qualitative analysis. *Med J Aust* 2004;**181**:36–9.
- 16 **Welsh CH**, Pedot R, Anderson RJ. Use of morning report to enhance adverse event detection. *J Gen Intern Med* 1996;**11**:454–60.
- 17 **Vincent C**, Stanhope N, Crowley-Murphy M. Reasons for not reporting adverse incidents: an empirical study. *J Eval Clin Pract* 1999;**5**:13–21.
- 18 **Wilson RM**, Runciman WB, Gibberd RW, *et al*. The Quality in Australian Health Care Study. *Med J Aust* 1995;**163**:458–71.
- 19 **Thomas EJ**, Studdert DM, Burstin HR, *et al*. Incidence and types of adverse events and negligent care in Utah and Colorado. *Med Care* 2000;**38**:261–71.
- 20 **Uribe CL**, Schweikhart SB, Pathak DS, *et al*. Perceived barriers to medical-error reporting: an exploratory investigation. *J Healthc Manag* 2002;**47**:263–79.
- 21 **Lawton R**, Parker D. Barriers to incident reporting in a healthcare system. *Qual Saf Health Care* 2002;**11**:15–8.
- 22 **Eland IA**, Belton KJ, van Grootheest AC, *et al*. Attitudinal survey of voluntary reporting of adverse drug reactions. *Br J Clin Pharmacol* 1999;**48**:623–7.
- 23 **Firth-Cozens J**. Barriers to incident reporting. *Qual Health Care* 2002;**11**:7.
- 24 **Elnitsky C**, Nichols B, Palmer K. Are hospital incidents being reported? *J Nurs Admin* 1997;**27**:40–6.
- 25 **Creswell J**. *Research design: qualitative, quantitative and mixed methods approaches*, 2nd edition. California, US: Sage Publications, 2003.
- 26 **Richardson S**, Allen J. Casualization of the nursing workforce: a New Zealand perspective on an international phenomenon. *Int J Nurs Prac* 2001;**7**:104–8.
- 27 **Bent PD**, Bolsin SN, Creati BJ, *et al*. Professional monitoring and critical incident reporting using personal digital assistants. *Med J Aust* 2002;**177**:496–9.
- 28 **Evans S**. *Evaluation of an incident reporting to improve hospital systems (IRIS) project*, April 2003. Available at http://www.safetyandquality.sa.gov.au/site/page.cfm?area_id=10&nav_id=506 (accessed 25 October 2005).
- 29 **New South Wales Department of Health**. *Patient Safety and Clinical Quality Program: First Report on Incident Management in the NSW Public Health System 2003–2004*. NSW, Australia: NSW Department of Health, 2005. Available at http://www.health.nsw.gov.au/pubs/2005/incident_mgmnt.html (accessed 25 October 2005).