



Travelling far but staying close to home

Nicola Low, Sevgi Aral and Jackie A Cassell

Sex. Transm. Inf. 2007;83:501-502
doi:10.1136/sti.2007.028928

Updated information and services can be found at:
<http://sti.bmj.com/cgi/content/full/83/7/501>

These include:

References

This article cites 18 articles, 16 of which can be accessed free at:
<http://sti.bmj.com/cgi/content/full/83/7/501#BIBL>

Rapid responses

You can respond to this article at:
<http://sti.bmj.com/cgi/eletter-submit/83/7/501>

Email alerting service

Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article

Topic collections

Articles on similar topics can be found in the following collections
[Editor's choice](#) (366 articles)

Notes

To order reprints of this article go to:
<http://journals.bmj.com/cgi/reprintform>

To subscribe to *Sexually Transmitted Infections* go to:
<http://journals.bmj.com/subscriptions/>

Travelling far but staying close to home

Nicola Low, Sevgi Aral, Jackie A Cassell

An overwhelming majority of the global burden of morbidity and mortality caused by sexually transmitted infections (STI) and HIV is borne by the world's poorest countries (fig 1). In a world increasingly "shrunk" by modern communications, what are the common issues preoccupying the international community of researchers in our field? In planning this themed issue, published on World AIDS Day 2007, we sought papers on all topics of global relevance: migration, international travel, the extent and impact of antiretroviral rollout programmes, and wide-reaching strategies for the prevention of STI and HIV transmission. We took the opportunity to seek out international diversity in STI and HIV research. Despite our intention to travel far afield, we have found ourselves closer to home than we had hoped, in respect of the authorship, and ownership of the diversity of studies relating to the populations of the world's poorer countries. Our 13 original articles include 9 studies conducted in the countries of sub-Saharan Africa,¹⁻⁶ south and south east Asia,^{7,8} Latin America,⁹ and 4 in Europe.¹⁰⁻¹³ The provenance of the articles, however, does not reflect their scope (fig 2); the e-mail addresses of corresponding authors of seven of these papers are in the United Kingdom,^{1,5,8,10-13} and another five are in the United States, Canada and Australia.^{2,4,6,7,9} On exploring this further, we found that the pattern of the published content was not the result of the selective acceptance of papers authored from the north; we saw the same pattern in papers that were rejected. Neither is ours an isolated experience. In the six leading tropical medicine journals in 2002, 12.5% of the first and 9.4% of the last authors came from countries with the lowest human development indices (usually poor) compared with 50.6% and 58.8% from high human development index (usually rich) countries.¹⁴ Within the increasingly global research community, it is essential that local research capacity is developed in its fullest form. This means that opportunities must be created for local scientists to be full and active partners in the design, analysis, interpretation and presentation of STI and HIV research with a view to the ownership of research by those whose own destiny lies within those countries.¹⁴

The control of HIV infection, and the reduction of associated morbidity in severely affected countries is a major theme. Boulle and Ford¹⁵ from South Africa give an overview of the benefits and challenges of scaling up antiretroviral therapy in resource-poor settings. Encouragingly, they cite some evidence that in South Africa and Brazil death rates and AIDS-related hospital admissions are decreasing at a population level. Vaginal microbicides are a hotly pursued option for HIV prevention, but the failures and closures of clinical trials, despite extensive safety testing, have dampened enthusiasm

recently. Martin Hilber and colleagues¹⁶ suggest that the missing link is a proper understanding of the role of women's use of vaginal products and practices to tighten, dry, warm and clean their vaginas. The goal, they say, is to achieve a safer vaginal environment.

The new global strategy of the World Health Organization (WHO) for the prevention and control of STI¹⁷ is summarised by Lewis and colleagues.¹⁸ It lists the components of comprehensive STI case management, including correct diagnosis, effective treatment, and condom promotion. The need for new technologies to contribute to STI control is further developed by Jones *et al.*,² who show in a randomised controlled trial, that allowing women in Gugulethu, South Africa, to take specimens for STI at home and mail them to the laboratory is a feasible and acceptable alternative to going to a clinic. In the same vein, Romoren *et al.*³ used decision analysis to show that the use of point of care tests for diagnosis and azithromycin




Figure 1 World map of distribution of deaths due to STI. Territories are sized in proportion to the absolute number of people who died from STI other than HIV in one year. Source: http://www.worldmapper.org/display_extra.php?selected=375. Reproduced with permission.




Figure 2 World map of the distribution of authors of scientific papers. Territory size shows the proportion of all scientific papers published in 2001 written by authors living there. Source: <http://www.worldmapper.org/display.php?selected=205>. Reproduced with permission.

for treatment of chlamydia could improve the management of chlamydia in pregnant women, but would cost an additional US\$38 per case cured. Although systematic review of the evidence for the effectiveness of condom promotion interventions in Africa and Asia supports the WHO strategy, Foss and colleagues¹⁰ also draw attention to the need for better evaluation designs using biological outcomes. In practice, levels of risk-taking behaviour remain high and condom use low, as O'Farrell *et al*¹ show in a study of men attending an STI clinic in Durban, South Africa.

An immediate priority of the WHO strategy is the implementation of targeted interventions against STI and HIV in high-risk and vulnerable populations. The Avahan Project is a US\$258 million complex HIV prevention intervention aimed at sex workers and drug users in the six most heavily affected states in India. Boily and colleagues⁸ describe a novel approach to evaluating the effects of this programme in sex workers. They will collect STI, HIV and behavioural data over time from intervention areas and use mathematical models to predict what would have happened to the HIV epidemic in the absence of the intervention. There will be no quick answer though; the evaluation will take over seven years.

The potential of voluntary testing and counselling is also explored. Arthur *et al*⁵ observed some reduction in risk behaviours of 401 people interviewed six months after an HIV test in Kenya, but only 55% of those testing HIV positive disclosed the result to their partner. The benefits of an HIV test in sex workers in Senegal are even less clear, because women who had had an HIV test in the past were less likely to use condoms with their regular partners.⁶ Prost *et al*¹³ conducted focus groups with Africans living in London and concluded that a voluntary counselling and testing approach used in health centres in Kenya could be feasible in the United Kingdom. Promoting HIV testing among Africans in the United Kingdom would certainly be worthwhile. Sadler *et al*¹² report a prevalence of 14% in a community-based survey of 1359 adults

from sub-Saharan Africa, of whom 66% were unaware of their infection.

Finally, Mercer *et al*¹¹ explored the potential of travel for disease transmission, showing that HIV prevention advice in the United Kingdom needs to be targeted to Britons travelling abroad as well as to migrants arriving from high-prevalence HIV countries. Analysis of the second National Survey of Sexual Attitudes and Lifestyles shows that a substantial minority of young single adults reports sex abroad, with its associated risks.

As Peter Piot said in his recent interview with STI, "it is important that young people enter [STI research] because there is still so much work to be done."¹⁹ We hope that this collection of articles contributes to global and local efforts for the prevention and control of STI and HIV to provide a safer sexual environment for all.

Sex Transm Infect 2007;**83**:501–502.
doi: 10.1136/sti.2007.028928

.....

Authors' affiliations

Nicola Low, University of Bern, Bern, Switzerland

Sevgi Aral, Centers for Disease Control and Prevention, Atlanta, Georgia, USA

Jackie A Cassell, Brighton and Sussex Medical School, Brighton, UK

Correspondence to: Nicola Low, Institute of Social and Preventive Medicine, University of Bern, Finkelhubelweg 11, Bern CH-3012, Switzerland; low@ispm.unibe.ch

We thank Professor Danny Dorling for alerting us to the map used in figure 2, and for permission to reproduce these.

REFERENCES

- O'Farrell N, Morison L, Moodley P, *et al*. High risk sexual behaviour in men attending an STI clinic in Durban, South Africa. *Sex Transm Infect* 2007;**83**:530–3.
- Jones HE, Altini L, de Kock, A, *et al*. Home-based versus clinic-based self-sampling and testing for sexually transmitted infections in Gugulethu, South Africa: randomised controlled trial. *Sex Transm Infect* 2007;**83**:552–7.
- Romoren M, Hussein F, Steen TW, *et al*. Costs and health consequences of chlamydia management strategies among pregnant women in sub-Saharan Africa. *Sex Transm Infect* 2007;**83**:558–66.

- Lowndes CM, Alary M, Labbe AC, *et al*. Interventions among male clients of female sex workers in Benin, West Africa: an essential component of targeted HIV preventive interventions. *Sex Transm Infect* 2007;**83**:577–81.
- Arthur G, Nduba V, Forsythe S, *et al*. Behaviour change in clients of health centre based voluntary HIV counselling and testing (VCT) services in Kenya. *Sex Transm Infect* 2007;**83**:541–6.
- Wang C, Hawes SE, Gaye A, *et al*. HIV prevalence, previous HIV testing, and condom use with clients and regular partners among Senegalese commercial sex workers. *Sex Transm Infect* 2007;**83**:534–40.
- Zhang D, Bi P, Lv F, *et al*. Internet use and risk behaviours: an online survey of visitors to three gay websites in China. *Sex Transm Infect* 2007;**83**:571–6.
- Boily MC, Lowndes C, Vickerman P, *et al*. Evaluating large-scale HIV prevention interventions: study design for an integrated mathematical modelling approach. *Sex Transm Infect* 2007;**83**:582–9.
- Blas MM, Alva IE, Cabello R, *et al*. Internet as a tool to access high-risk men who have sex with men from a resource-constrained setting: a study from Peru. *Sex Transm Infect* 2007;**83**:567–70.
- Foss AM, Hossain M, Vickerman PT, Waits CH. A systematic review of published evidence on intervention impact on condom use in sub-Saharan Africa and Asia. *Sex Transm Infect* 2007;**83**:510–16.
- Mercer CH, Fenton KA, Wellings K, *et al*. Sex partner acquisition while overseas: results from a British national probability survey. *Sex Transm Infect* 2007;**83**:517–22.
- Sadler KE, McGarrigle CA, Elam G, *et al*. Sexual behaviour and HIV infection in black-Africans in England: results from the Mayisha II survey of sexual attitudes and lifestyles. *Sex Transm Infect* 2007;**83**:523–9.
- Prost A, Ssanyu Sseruma W, Fakoya I, *et al*. HIV voluntary counselling and testing for African communities in London: learning from experiences in Kenya. *Sex Transm Infect* 2007;**83**:547–51.
- Keiser J, Utzinger J, Tanner M, *et al*. Representation of authors and editors from countries with different human development indexes in the leading literature on tropical medicine: survey of current evidence. *BMJ* 2004;**328**:1229–32.
- Boule A, Ford N. Scaling up antiretroviral therapy in developing countries: what are the benefits and challenges? *Sex Transm Infect* 2007;**83**:502–4.
- Martin Hilber A, Chersich MF, van de Wijgert JHHM, *et al*. Vaginal practices, microbicides and HIV: what do we need to know? *Sex Transm Infect* 2007;**83**:505–6.
- World Health Organization. Global strategy for the prevention and control of sexually transmitted infections: 2006–2015. http://www.who.int/reproductive-health/publications/stisstrategy/stis_strategy.pdf 2007 (accessed 22 Oct 2007).
- Lewis DA, Latif AS, Ndowa F. The global strategy for the prevention and control of sexually transmitted infections: time for action. *Sex Transm Infect* 2007;**83**:507–9.
- Vandenbruene M. Interview with Peter Piot. *Sex Transm Infect* 2007;**83**:436–40.