

## PUBLISHED VERSION

Bulte, Erwin; Boone, Randall B.; Stringer, Randy; Thornton, Philip K.  
Elephants or onions? Paying for nature in Amboseli, Kenya, *Environment and Development Economics*, 2008; 13(3):395-414.

Copyright © 2008 Cambridge University Press

### PERMISSIONS

<http://journals.cambridge.org/action/stream?pageId=4088&level=2#4408>

The right to post the definitive version of the contribution as published at Cambridge Journals Online (in PDF or HTML form) in the Institutional Repository of the institution in which they worked at the time the paper was first submitted, or (for appropriate journals) in PubMed Central or UK PubMed Central, no sooner than one year after first publication of the paper in the journal, subject to file availability and provided the posting includes a prominent statement of the full bibliographical details, a copyright notice in the name of the copyright holder (Cambridge University Press or the sponsoring Society, as appropriate), and a link to the online edition of the journal at Cambridge Journals Online. Inclusion of this definitive version after one year in Institutional Repositories outside of the institution in which the contributor worked at the time the paper was first submitted will be subject to the additional permission of Cambridge University Press (not to be unreasonably withheld).

10<sup>th</sup> December 2010

<http://hdl.handle.net/2440/47136>

# Elephants or onions? Paying for nature in Amboseli, Kenya

ERWIN H. BULTE\*

*Development Economics Group, Wageningen University, and Department of Economics, Tilburg University, P.O. Box 8130, 6700 EW Wageningen, Netherlands. Email: Erwin.bulte@wur.nl*

RANDALL B. BOONE

*Natural Resource Ecology Laboratory, 1499 Campus Delivery – B234 NESB, Colorado State University, Fort Collins, CO 80523-1499, USA. Email: rboone@nrel.colostate.edu*

RANDY STRINGER

*School of Agriculture, Food and Wine, University of Adelaide. Email: randy.stringer@adelaide.edu.au*

PHILIP K. THORNTON

*International Livestock Research Institute, Nairobi, Kenya and the Institute of Atmospheric and Environmental Sciences, School of Geosciences, University of Edinburgh, Edinburgh, UK. Email: p.thornton@cgiar.org*

**ABSTRACT.** Traditional grazing grounds near Amboseli National Park (Kenya) are being rapidly converted to cropland – a process that closes important wildlife corridors. We use a spatially explicit simulation model that integrates ecosystem dynamics and pastoral decision-making to explore the scope for introducing a ‘payments for ecosystem services’ scheme to compensate pastoralists for spillover benefits associated with forms of land use that are compatible with wildlife conservation. Our break-even cost analysis suggests that the benefits of such a scheme likely exceed its costs for a large part of the study area, but that ‘leakage effects’ through excessive stocking rates warrant close scrutiny.

## 1. Introduction

The leading approach to conservation of large mammals is to create protected areas (Brandon *et al.*, 1998; Margules and Pressey, 2000; Adams *et al.*, 2004). Protected area systems involve top-down command-and-control measures, including fencing off specific areas to restrict their use.

\*Corresponding author.

We would like to thank two anonymous referees, Mike Norton Griffiths, Richard Hatfield, Harvey Croze, Paul Ntiati, Alain Lambert, FAO Roles of Agriculture workshop participants, and DIVERSITAS conference participants for helpful suggestions and comments on an earlier version of this paper.

While recent research suggests that protected areas can be effective in some cases (Bruner *et al.*, 2001), many developing country experiences have been disappointing (Brandon *et al.*, 1998; Adams and Hulme, 2001; Western, 2001). The difficulties with protected areas include: (i) their size – often parks are too small to sustain a full range of services; (ii) inadequate administrative and management capacity; (iii) limited resources for monitoring and enforcement; and (iv) denying access to local communities, especially the poor who may depend directly on the restricted ecosystem for survival. When rights and access to ecosystem services shift in ways that adversely impact local communities, poor households have little incentive to maintain or use ecosystems in a sustainable way (Barbier, 1992; Tisdell, 1995; Damania *et al.*, 2003).

The limitations of protected area approaches led to efforts aimed at aligning the interests of the poor with conservation objectives through community-based resource management programs. The importance of integrating local communities into protected-area planning dates back to at least the 1982 World Parks Congress (Adams *et al.*, 2004). This participatory-style conservation approach, mostly driven by encouragement of external donors, seeks to provide local communities with incentives to protect crucial ecosystems through sharing products, responsibilities, and decision-making authority. However, evaluations and studies continue to raise important concerns about the appropriateness of community-based conservation efforts, arguing that positive and lasting success is elusive when development projects combine biodiversity conservation goals with poverty reduction goals (Simpson, 1995; Murombedzi, 1999).

Other studies question the assumptions linking local communities and sustainable resource use across diverse geographic conditions and economic situations (Barrett and Arcese, 1995, 1998). An analysis of community based natural resource management in Kenya concludes that it did *not*: (i) result in more equitable distribution of economic benefits; (ii) reduce conflicts; (iii) consider traditional Maasai grazing and wildlife movement knowledge; (iv) improve biological diversity protection; or (v) improve sustainable resource use (Kellert *et al.*, 2000). The Kenya case focused on the Kimana Community Wildlife Sanctuary, an important area ecologically serving as a wildlife corridor between two protected areas, Amboseli and Tsavo National Parks. The study region of this paper encompasses the Kimana wildlife corridor.

More recently, a variety of compensation and market-related policies have gained prominence to encourage ecosystem and land managers to change behaviour. While direct financial and market incentive schemes, commonly referred to as direct payments for ecosystem services (PES), now exist in many developed countries – e.g., within the European Union there are elaborate payment schemes for the conservation of waders (meadow birds) – experiences in many developing countries (in particular, in countries other than in Latin America) are more limited (e.g., Landell-Mills and Porras, 2002; Wunder, 2005). If conservation can be promoted directly through PES, it would potentially benefit the poor as well (Pagiola *et al.*, 2005).

Ecosystems provide a plethora of services to human communities, and these services benefit people at a wide range of scales – varying from local

to national and even global scale (MEA, 2005). Unless compensation takes place for ecosystem services spilling over to areas beyond their source, such services will likely be ignored by local 'suppliers' and, if those services deteriorate, too little will be provided relative to the desires of the regional, national, or global community. The idea behind PES to remedy this situation is simple: beneficiaries of these services should make 'direct, contractual and conditional payments to local landholders and users in return for adopting practices that secure ecosystem conservation and restoration' (Wunder, 2005). The type of ecosystem service and the geographical location of the beneficiary (*vis-à-vis* the location of the source) are important determinants of the form of the contract that can be written.

To date, PES activities in developing countries most often address watershed issues where feedback loops are 'tight' and where suppliers and demanders are easy to identify. Less work has been done on payments for global values, although there is some experience with carbon sequestration projects in the context of Joint Implementation, or the Clean Development Mechanism. In this study we are concerned with one particular type of ecosystem service, namely benefits associated with biodiversity conservation (more specifically, conservation of elephants, *Loxodonta africana*, in a beautiful landscape). The nature of this service implies two types of service beneficiaries could potentially fund a payment scheme: (i) eco-tourists visiting the area and enjoying non-consumptive use values, and (ii) households in various parts of the world enjoying non-use values of conservation.

From a 'global efficiency perspective' these two types of value should be added, but it is evident that tapping into the latter source of funding is not easy. There is no 'world government' that can tax citizens in the West to finance the provision of public goods in the South. While the Global Environmental Facility (GEF) initiative allows funding for non-use values of international importance, it is not designed for sustained financing. Targeting 'users' of the ecosystem services – tourists visiting the area on a safari – is likely a more viable long-run option for the payment scheme.

This paper explores the opportunity to establish an international payment system for ecosystem services associated with wildlife (elephant) conservation near Kenya's Amboseli National Park (NP). Under current trends the long-term future of the Amboseli ecosystem (and its icon – the elephant) looks rather bleak.<sup>1</sup> Ultimately, impoverished and remote

<sup>1</sup> In addition to the issue of land conversion discussed in this paper, the future of the Amboseli ecosystem may be compromised by the following three factors: (i) the upper forest line on Mount Kilimanjaro shifts down due to an increased frequency and intensity of forest fires, which may have severe repercussions for local climatic and hydrological conditions (for a discussion of the importance of this issue *vis-à-vis* the more conspicuous issue of the disappearing glacier, see Hemp, 2005); (ii) there is a push in the group ranches to sub-divide the communally owned grazing grounds into private plots, which would adversely affect the capacity of the system to support grazing (Boone *et al.*, 2005); and (iii) recently the government of Kenya announced that it would degazette the area from National Park to Game Reserve, and that management would be relegated from the Kenya Wildlife Service

communities are the main drivers of changes in local land use. Because wildlife in many protected areas, including Amboseli, depend on access to (food) resources found on private lands, the success of conservation efforts – elephants or onions? – is determined by the balance between benefits and costs as perceived by these private agents. The objective of this paper, therefore, is twofold. First, we explore whether efforts to promote elephant conservation near Amboseli NP through a PES scheme represent a viable economic proposition, or not. While a cost–benefit analysis is beyond the scope of the study, we aim to provide a rough indication of the costs involved in elephant conservation, and then establish whether these costs are likely to be outweighed by the increase in welfare from ‘more elephants’. The outcome of such a comparison may be used to decide whether strategies should be implemented that provide incentives for local households to sustainably manage their rangelands and share this habitat with wildlife. The second, and closely related objective, is to predict how a PES scheme affects conservation and welfare of the Maasai. To address the second question one would ideally use a household model, but as a fully calibrated Maasai model is not available, we resort to an approximation instead.

The paper is organized as follows. Section 2 provides a brief profile of the Amboseli ecosystem. In section 3, we sketch the bare bones of the SAVANNA and PHEWS models that are used to simulate the impacts, in terms of changes in land use, income, and elephant abundance, from a PES system. Section 4 presents the simulation results as well as the break-even cost analysis. Section 5 concludes.

## **2. The Amboseli Ecosystem**

The Amboseli ecosystem, an area of some 8,000 km<sup>2</sup>, comprises part of the Ilkisongo region of southeastern Kajiado District in Kenya and the Longido region of northern Tanzania. Amboseli is typical of African arid rangelands: rainfall is low and unpredictable in time and space. At the heart of the ecosystem is Amboseli NP, the core of a UNESCO Man and the Biosphere Reserve protecting 392 km<sup>2</sup> (about 5 per cent) of the wildlife dispersal area. Amboseli’s swamps are fed by subsurface water that percolates through volcanic rock from the forested catchment of Kilimanjaro rising spectacularly to the south. Nearly four decades of ecological monitoring and research, as well as two of the world’s longest studies of elephants and primates, have brought Amboseli international scientific and conservation recognition.

Amboseli NP is fundamental to Kenya’s tourist industry, typically ranking second among parks in annual park gate fees – around USD 3.5 million in 2004. In the past, the absence of wide-scale intensive agriculture and the relatively low population density encouraged and provided refuge to a magnificent array of biodiversity, including large and small mammals, birds, reptiles, insects and plants, some of which are rare

(KWS) to the district Maasai Council (but this challenge to the status of the park is still in court). An integrated, international conservation effort would presumably need to tackle these challenges in tandem. The current study provides a first step.

or threatened. Birdlife International has named Amboseli one of the world's Important Bird Areas.

At present, market, policy and institutional incentives interact in ways that weaken the region's ecological integrity, endanger the wildlife tourism industry and threaten the long-term viability of rural households. Wildlife habitats are diminishing, migration corridors are narrowing, water resources are being degraded, livestock-wildlife competition is worsening, income inequality is increasing, and human-wildlife conflicts are mounting (Campbell *et al.*, 2000; Reid *et al.*, 2004). Human-human conflicts are increasing too, as the interests of local communities, park managers, and wildlife tourism providers increasingly clash (Hoare, 1999; Campbell *et al.*, 2003).

The Amboseli ecosystem is home to Maasai pastoralists whose long-practiced livestock activities are well adapted to the variable habitat, and whose land use decisions are a key driver of wildlife abundance in and around the Park. However, the majority of Maasai households receive virtually no direct benefits from the wildlife tourism industry. The cash benefits are not distributed fairly nor equally to the pastoralists (Kellert *et al.*, 2000; Mburu *et al.*, 2003). And the indirect benefits, in the form of reduced school fees, irrigation infrastructure maintenance, livestock sales yards, and other related community goods, often fail to benefit those in most need.<sup>2</sup> The Maasai do bear the costs of managing wildlife habitats, including personal safety, grazing competition, investments to minimize risks, management costs, damage to crops (from eating and trampling), and damage to livestock through the spread of diseases and killing (Norton-Griffiths and Southey, 1995; Campbell *et al.*, 2002).

The Maasai are organized in so-called group ranches, which are communally owned stretches of land. Currently, the property rights system in the traditional Maasai territory is in a state of flux, and many group ranches have been (or are in the process of) subdividing communal land ownership (refer to Boone *et al.*, 2005 and the references therein for more information on the subdivision process). Subdivision implies titling land and allocating it to individuals. The Maasai choose between two decision alternatives. First, they can rent out their land to farmers (or they can farm their land themselves). Depending on the location of the land, irrigated or rain-fed agriculture is feasible, resulting in an area-specific flow of rental payments (or profits). Second, they can use their land for pasture and earn a living as pastoralists, herding goats and cattle. If they do so, we approximate their stocking decisions based on a series of behavioral rules – the PHEWS model discussed below.

<sup>2</sup> The Maasai communities surrounding the Park are themselves divided about the benefits they obtain from the park (in the form of revenue sharing and job opportunities), and are frustrated that certain beneficial policies that were promised have never been implemented, such as water boreholes outside Amboseli NP. Factions within these communities are dissatisfied with the benefits they obtain, and threaten to intensify pressure on key natural resources in the Park (mainly forage and water) unless they will receive a larger share of the Park's proceeds.

From a conservation perspective the pastoral outcome is preferred. Some of the irrigated land was fenced during the late 1990s to protect crops from wildlife, and increasingly those protected croplands impede access to water, food, breeding grounds, and to the seasonal migration of wildlife up and down the slopes of Mount Kilimanjaro, and between Amboseli and other protected areas like Tsavo NP. (In addition, albeit somewhat beside the main point of this paper, there is evidence that agricultural use of the former grazing grounds is not sustainable because of water pollution, agrochemical use, and soil runoff.) Wildlife populations that had access to all of Amboseli's swamps until the 1970s, now have no access to one swamp and only partial access to three others (Reid *et al.*, 2004).

The Maasai have increasingly rented out large areas for irrigated or rain-fed agriculture during the past decade. During the past 20 years, in the adjacent areas to the south and east of Amboseli NP (Loitokitok Division), human populations have more than tripled, rain-fed agricultural areas expanded by 3.5 times, and irrigated area increased by 18 times, from around 250 ha to 4800 ha (Campbell *et al.*, 2003). While cropping may be *privately* rational (the returns of cultivation dominate the private returns of wildlife management), it is an open question whether it is also socially beneficial – i.e., what happens when we include ecosystem services benefiting people outside the Amboseli ecosystem in the picture?

### 3. The model

The majority of elephants and the other migrating species cannot survive without Amboseli's larger ecosystem, migrating seasonally between the Park and its surroundings. The future of much of Amboseli's wildlife lies in the hands of the people surrounding the Park. Six communally owned group ranches surround the Park. The predominant form of land use, livestock raising, is compatible with wildlife conservation. While livestock and wild herbivores may compete for forage, and predators may occasionally kill livestock, historical grazing systems and population pressures seemed to be sustainable and allowed for the co-existence of domesticated and wild animals.

#### 3.1 The SAVANNA model

To address management and policy questions relevant to wildlife and livestock requires an integrated approach (Coughenour *et al.*, 2002). We built upon an integrated assessment of southern Kajiado District (Boone *et al.*, 2005; Thornton *et al.*, 2006) that uses a process-based, spatially explicit ecosystem model called SAVANNA. SAVANNA has been applied at sites throughout the semi-arid areas of the world, but was first developed within Turkana District, Kenya (Coughenour, 1985). Many subsequent improvements and applications have been described (e.g., Coughenour, 1992; Buckley *et al.*, 1993; Ellis and Coughenour, 1998; Boone *et al.*, 2002, 2004, 2005; Thornton *et al.*, 2004, 2006; Boone, 2005). A schematic outline of the model is provided in figure 1a.

SAVANNA is a series of FORTRAN computer programs that join to model primary ecosystem interactions, simulating functional groups for plants and animals (e.g., perennial and annual grasses, cattle, grazing antelopes)




Figure 1a. Schematic outline of the SAVANNA model used to simulate the impact of land use change on wildlife (elephant) abundance




Figure 1b. Schematic outline of the energy flow in the PHEWS model used to simulate behavior of pastoralists

over periods from 10 to 100 or more years (Ellis and Coughenour, 1998). The model represents landscapes by dividing them into a system of square cells, and uses a series of digital maps (GIS maps) to characterize each cell. The model predicts water and nitrogen availability to plants using rainfall and soil properties, for each cell in the map. Based upon water, light, and

nutrient availability, quantities of photosynthate are calculated for each of the plant functional groups. Photosynthate is distributed to the plant parts based on established allometrics, yielding estimates of primary production. Plant populations are calculated from investments in reproductive parts.

A habitat suitability index is calculated for each cell, at weekly intervals and for each animal functional group, based upon forage quality and quantity, slope, elevation, cover, and the density of herbivores. Herbivores are distributed on the landscape based upon these indices. Importantly for this study, where we are interested in the system's response to the presence or absence of fenced-in cropland, maps may be used to modify the distribution of herbivores. Animals feed upon the available vegetation, depending upon dietary preferences and consumption rates. The energy gained is reduced by energy costs associated with basal metabolism, gestation, and lactation. Net energy remaining goes toward weight gain, with weights reflected in animal condition indices. Charts and maps are produced at monthly intervals (e.g., Boone *et al.*, 2002).

Seven plant functional groups are captured in the case study's SAVANNA application: palatable grasses, palatable forbs, unpalatable grasses and forbs, papyrus (*Cyperus papyrus*) swamps, palatable shrubs, unpalatable shrubs, and deciduous woodlands. Nine animal groups are modeled: three livestock species (cattle, goats and sheep), and six wildlife groups (wildebeest, *Connochaetes taurinus*; zebra, *Equus burchellii*; African buffalo, *Syncerus caffer*; grazing antelope; browsing antelope; and elephants). See Boone *et al.* (2005) for species comprising grazing and browsing antelope groups. A variety of data sources were used to parameterize the application for southern Kajiado District, described in Boone *et al.* (2005), including examples of literature used. The ecosystem model was calibrated using sources such as a net primary production database (Kinyamario, 1996), satellite imagery, which relates well to regional stocking levels (Oosterheld *et al.*, 1998), and information from important literature sources (e.g., De Leeuw *et al.*, 1998).

While formal validation studies of the SAVANNA model are unavailable (and perhaps impossible in light of the model's integrative nature), various efforts demonstrate that the model provides simulation results that seem to mirror the 'big pictures' in reality (e.g., Boone *et al.*, 2002, 2004). Adjustment and assessment of the parameterization of SAVANNA applications (such as the Kajiado model used in this study) include comparisons of model output to actual NDVI surfaces (the normalized difference vegetation index made from satellite imagery), NPP estimates based on field data (e.g., Kinyamario, 1996), comparisons of simulated and observed herbivore populations (Rykiel, 1996) from aerial surveys conducted by the Kenyan government, and comparisons to field data (e.g., ecological sampling to quantify standing green and standing dead biomass, as in Boone *et al.*, 2002), sometimes collected for the purpose.

### 3.2 The PHEWS Model

In an ideal world the SAVANNA model would be linked to a process-based, detailed, and fully calibrated household model that captured the myriad of response Maasai may have to changing circumstances. Such a model

Table 1. *Cash flows in the Pastoral Household and Economic Welfare Simulator (PHEWS)*

| <i>Flow</i>                        | <i>How treated in PHEWS</i>   |
|------------------------------------|---|
| <i>Cash in</i>                     |   |
| Livestock sales                    | Triggered by cash needs   |
| Crop sales                         | Calculated as a household characteristic (% sold); remainder is consumed by household |
| Wages                              | From input file   |
| Milk sales                         | Calculated as a household characteristic (% sold); remainder is consumed by household |
| Other (PES, gifts, transfers, etc) | From input file   |
| <i>Cash out</i>                    |   |
| Food purchases                     | Calculated from energy flow (figure 1b)   |
| Household goods                    | From input file   |
| Livestock purchases                | Triggered by cash needs   |
| Other payments out                 | From input file, plus crop inputs   |

Source: Thornton *et al.* (2003, 2006).

would combine the preferences of the Maasai households with respect to goods and services they consume (including their utility from leisure and 'life style' considerations that are undoubtedly relevant in this context) with a set of constraints – a budget constraint, time constraint, production possibilities, etc. However, such a model is not available for the study area. Instead, we use an approximation of such a model, calibrated for pastoral households in East Africa, called PHEWS (Pastoral Household Economic Welfare Simulator – see Thornton *et al.*, 2003, 2004, and 2006 for details). We distinguish between 24 different household 'groups' in this study – the product of eight livelihood strategies and three wealth levels (poor, medium, rich). For details about the classification, please refer to Thornton *et al.* (2006).

PHEWS is based on a set of rules that households follow when trying to secure caloric intake (and also accounts for some probabilistic slaughter for ceremonial occasions). It is well known that rainfall and income from herding are highly volatile in this part of the world. PHEWS keeps track of dietary energy flows and prescribes a certain series of actions when intake falls short of a desired level. An overview of the energy flow in the household, and associated behavioral rules, is provided in figure 1b. The household has access to two types of assets to finance its consumption of energy: (i) a so-called 'cash box' that evolves in accordance with expenditures and income from various sources, and (ii) a livestock herd. An overview of the relevant additions to, and subtractions from, the cash box is provided in table 1. The livestock herd dynamics are governed by a simple set of rules governing livestock sales and purchases. Livestock sales are triggered by specific cash needs above certain levels, and similarly livestock

purchases (which are much rarer) are triggered when the household has accumulated specific levels of available cash. The rules and triggers used were developed from household data collected by BurnSilver (2007) and formed part of the calibration of PHEWS for Kajiado – details are discussed in Thornton *et al.* (2006).

In sum, pastoral households have a target caloric intake and specific cash needs. They are assumed to use livestock as a buffer in periods when household income and consumption are low, and livestock numbers will be built up when income is high and caloric requirements are easily satisfied. Remaining funds are placed in the ‘cash box’ where they are stored for future use when income is low (Thornton *et al.*, 2006). When caloric intake from consumption of animal products, maize and sugar is insufficient to meet the threshold, the household tries to use its ‘cash box’ (if available) to buy maize, to make up for the deficit. If this fails, they sell a goat or cow. If all fails, the model assumes that there will be outside relief from some exogenous source. For this reason the model is not particularly useful for capturing Malthusian population dynamics, say, and we simply assume that the human population is constant.<sup>3</sup>

The SAVANNA and PHEWS models are tightly linked, and are literally part of the same simulation. At each time step modeled, SAVANNA passes information about livestock to PHEWS, and PHEWS passes information about the sale of animals and such back to SAVANNA.

### 3.3 *Three scenarios*

We distinguish between three different scenarios, which are compared to highlight effects of different management practices. In control scenario A we consider the base case where parts of the group ranches that surround the Park are converted to fenced-in cropland (but note that we assume that the fenced-in areas are used for cropping throughout the entire study period – from 1977 to 2000 – and that in reality fencing only started in the 1990s). We use historical rainfall patterns to simulate livestock and wildlife abundance over time and space. In the pastoral scenario B we explore the case where the fenced-in area is returned to grazing ground and accessible for wildlife and livestock alike. One may think of this as a command-and-control approach to conservation, simply banning the use of fences. We simulate the impact on wildlife and livestock, but also on Maasai income. Finally, in PES scenario C we consider what happens if we compensate the Maasai for restoring the grazing grounds. That is, in return for giving up the privately profitable option to rent out land to crop growers, the Maasai are assumed to engage in an easement deal with a funding agency that offers a competitive rate of return on the land. Compared to scenario B the Maasai

<sup>3</sup> In reality of course the population is not constant. It may change because of natural population growth and mortality, but also because of migration patterns. It is possible that both replenishment and migration react endogenously to implementation of a PES scheme. While we ignore this in the analysis, this is something that should be considered when actually transferring money.

budget constraint is therefore relaxed, which means that households are better able to meet their target consumption levels.<sup>4</sup>

This approach involves comparisons between simulations where the only attributes changed are areas available for grazing and payments to Maasai. The model is parameterized to agree with current conditions to the degree possible, but the approach is not predicated on responses being absolutely correct, but rather on comparisons between simulations that are otherwise parameterized identically. Our results are not intended to provide precise predictions about how the elephant population may change in the future; too many unforeseen circumstances may affect that trajectory. Rather, we provide examples of tradeoffs associated with PES systems, and identify the direction and magnitudes of change in wild and domestic ungulates, and in Maasai well-being.

#### 4. Simulation results

In this section we present the simulation results of the three scenarios, and we use these results as input in a break-even cost analysis. We try to address the question whether a PES scheme for elephant conservation is welfare enhancing at the global scale, or not. We also use the output to discuss the form that transfers from conservationists to pastoral households may take.

##### 4.1 Returning cropland to range land

Figure 2 summarizes the impact of returning the fenced-in cropland to grazing grounds on elephant abundance. The dashed upper line reflects elephant abundance in scenario B (no fences) and the solid lower line reflects the number of elephants in control scenario A. The figure also shows the historical pattern of rainfall in the study area (light dashed line).

Not surprisingly, expanding elephant habitat translates into a larger number of elephants. However, during the first 15 years of the simulation exercise, the impact is very modest – typically in the range of only 100 to 300 extra elephants per year, or a modest 15 per cent increase in abundance. It appears as if the pastoral scheme is hardly worthwhile. But the situation abruptly changes after 1992, when stocking rates are rather high and a serious drought hits the area. The elephant population in the control scenario collapses to about 50 per cent of its pre-drought level of abundance, while the elephant population in the pastoral scenario increases. Considering the entire study period from 1977 to 2000, the *average* number of elephants in the pastoral scenario is about 500 heads larger than in the control scenario. The averages tell only part of the story, however; the main benefit from removing fences is that the elephant population is much more resilient to changes in (environmental) conditions when it has access to a wider set of base resources.

<sup>4</sup> In reality a fourth scenario is being discussed: the case where fences are not removed but where agriculture *outside* the fences is controlled to enable a free flow of animals between areas. In theory we could readily analyse this case, but the resolution of the current model is too coarse to yield reliable results. The scenarios considered in this paper are more dramatic cases, sharply illustrating the main tradeoffs.




Figure 2. *Elephant abundance with (solid) and without (heavy dashed) habitat conversion*  
 Precipitation over 12 months (light dashed) is overlaid for comparison

The interpretation of these results is as follows. In times of sufficient rainfall, the swamp areas converted to cropland do not represent a key resource for elephants. Opening up these areas implies they have access to more food, so we observe a modest increase in the population. However, the picture changes in times of drought, when access to the swamps for food and water becomes necessary to support the elephants. If this access is denied, water and food become critical factors and the population crashes.

The elephant population in pastoral scenario B increases during the drought of the mid 1990s because of less competition from livestock. Faced with a drought, the Maasai have no option but to sell part of their large stock to support their families, to buy grain, and to purchase more drought-resistant small stock. The loss of milk from the large stock demands more large animal sales, which in turn means less milk, etc., in the downward spiral seen here and sometimes seen in Maasai communities. In the simulation, goats eventually came to dominate herd composition. This represents a fundamental tradeoff of the command-and-control option to conservation; if it is effective at promoting elephant conservation by restricting the Maasai's use rights of the swamps, the costs of this 'success' are borne entirely by the Maasai who see their herds shrink and income position deteriorate. Since most of the non-use values associated with conservation are transboundary, this is clearly unfair.

#### 4.2 *The effect of paying for ecosystem services*

Figure 3 summarizes the consequences of establishing a PES system, where the Maasai lease their cropland to a conservation agency (as opposed to crop farmers), and where the restored grazing grounds and swamps are




Figure 3. *The effect of paying Maasai for not renting out their lands on the elephant population*  
 A comparison of pastoral systems with (solid) and without compensation (dashed)

available for livestock and wildlife. The upper solid line represents the elephant population when a PES system is in place – the scenario C – and the lower dashed line, again, depicts pastoral scenario B discussed above, where fences have been removed but where no compensation takes place.

The first thing to notice is that a fair transfer to the Maasai did not compromise elephant conservation – the opposite is true. Key resource areas and other rangelands remained available because of the PES agreements limiting cultivation. Elephant numbers exceeded those when the entire area is pastoral because the transfer enabled the Maasai to support a livestock herd that was close to a sustainable size and composition for the ecosystem. While increasing livestock herd size is detrimental for conservation – livestock and wildlife compete for base resources – the same is not true for the changes in composition brought about by the PES system. Specifically, cattle diets overlap less with elephants than do goat diets (e.g., see Skarpe *et al.*, 2007). The PES system enabled the Maasai to gradually expand their cattle stock (towards a herd that exceeds the herd under pure pastoralism by some 4,000 heads, or an increase of some 25 per cent relative to the pastoral scenario B), and move away from goats. In the final periods of the simulation exercise, the goat herd under scenario C is some 10,000 heads smaller than in pastoral scenario B (representing a 33 per cent reduction). Because goats and elephants have overlapping diets – they compete to some degree for food – this change in the composition induced by a relaxed budget constraint favored elephants. By the same token: note that the change in livestock

composition from goats to cattle will adversely impact grazing species of wildlife that compete for food with cows.

The main insight is that poverty alleviation and conservation may go hand-in-hand. Implementation of a PES scheme will both make the Maasai better off (in our specification: they are fully compensated for the foregone returns from leasing out their land, and as a bonus they can use the restored grazing grounds for their own livestock) and will enhance and stabilize elephant populations. The lack of a tradeoff follows from ecological interactions between species, and capturing such interactions implies developing multi-disciplinary models as the one advanced here.

#### *4.3 First attempt at a break-even cost analysis*

The observation that the PES scheme makes elephants and Maasai better off does not necessarily imply that it is welfare enhancing, because there are costs to consider as well. How do the costs and benefits compare? A full cost-benefit analysis may account for the distributional consequences (giving extra weight to income of the Maasai) and should account for transaction costs, etc. The break-even cost analysis ignores these issues and focuses instead on a more narrow question: does the conservation value created by the PES exceed or fall short of the opportunity costs of conservation – the foregone returns to cultivating crops, proxied by the rental payments to Maasai?

Upon comparing control scenario A with PES scenario C, the PES scheme produces benefits of some additional 600–700 elephants per year (average value). How much does the international community value the conservation of some 650 elephants? Answering this question is not easy. First, we are interested in marginal values and this information is not available to our knowledge. Second, the willingness to donate money for elephant conservation projects likely increases with income. Geography matters as well, because elephants are a real threat to the safety of people who live with them (41 per cent of villagers polled in Cameroon wanted elephants removed or fenced in, and a significant minority wanted them shot – see WWF, 2000). When considering the non-use value of charismatic species like elephants, it is not obvious which reference population should be included in the aggregation exercise.

Because of the uncertainties that inevitably surround point estimates of the value of elephants we turn the question around: focus on the costs of conservation first, and then argue whether it is plausible that aggregate conservation values are sufficiently large to overcome these costs or not. Based on observations in the field we first use a payment of US \$10 per acre per year as a proxy for the opportunity costs of conservation. Multiplying the fee by the relevant area of cropland yields a total cost of \$112,500 per year. Assuming constant marginal cost, this translates into a cost of some \$175 per elephant per year (divide by 650).

Assuming that the marginal value of elephant conservation is constant (a strong assumption), a prerequisite for the PES scheme to be globally welfare enhancing is that households in Europe and the United States are

willing to pay \$0.60 per year for African elephant conservation.<sup>5</sup> Of course, it is an open question whether households are indeed willing to pay such amounts, but evidence gleaned from contingent valuation studies into the willingness to pay for other species (for an overview, see Loomis and White, 1996) suggests that this number is not excessive. One specific study aimed at valuing Asian elephants (*E. maximum*) also produced an estimate of willingness to pay amongst the people of Sri Lanka that would have been sufficiently high – some \$12 per household per year (Bandara and Tisdell, 2004). We conclude that a PES effort for the Amboseli region is likely to make good economic sense.

The discussion above, however, is incomplete and misses out on an important source of heterogeneity. While most of the rain-fed land can be obtained at a relatively low cost, this is not true for other areas where irrigated horticulture is possible. Net agricultural profits in such areas are considerably higher. Even if the landowners secure only part of these returns (with the remainder accruing to the farmer who does the actual work and faces most of the production and price risk) as rental payments, it is unlikely that a viable PES scheme could be set up that would be able to induce the Maasai to keep the land under pasture. In other words, while implementing a PES scheme to safeguard critical elephant habitat and wildlife corridors is feasible for large parts of the study area, there are also pockets where the returns to agriculture may dominate the (social) returns to herding and conservation.

#### 4.4 Exploring leakage

This section examines how robust these results are with respect to alternative specifications of Maasai behavior. According to the PHEWS model, pastoralists use PES funds to re-balance the composition of their livestock herd (purchasing extra cattle at the expense of goats and sheep), and store some of the money in their cash box for future use. What happens if, instead, *all* the new funds are used to purchase additional livestock in the *same* proportion as current livestock holdings? This would aggravate competition for food between livestock and wildlife and potentially attenuate the conservation benefits. Following earlier economic literature on such attenuating effects (such as in the literature on carbon emissions, see for example Felder, 1993) we refer to this outcome as ‘leakage’. We have used SAVANNA to explore this issue.

Representative results are provided in figure 4, depicting wildlife populations for three different scenarios: (i) PES payments going to households (solid line), which is just scenario C based on the PHEWS model, (ii) PES based on the assumption that all money is immediately converted into livestock (dashed line), and (iii) the control scenario A above (open line). The curves for scenario’s A and C are different than the ones

<sup>5</sup> The calculation is as follows. Current estimates of the African elephant population amount to some 500,000 head (Blanc *et al.*, 2003). Assuming a minimum benchmark cost of \$175 per elephant per year, the total benefits of elephant conservation should amount to  $\$87.5 \times 10^6$  per year. Dividing by the number of households ( $150 \times 10^6$ ) this amounts to \$0.60 per household per year.




Figure 4. Payments to pastoralists may result in larger livestock herds and accentuated competition for food between livestock and wildlife

Wildlife stocks in the presence (solid) and absence (open) of payments for ecosystem services, as well as if payments are used entirely to purchase livestock (dashed). (The definition for Large Herbivore Units is the same as for a Tropical Livestock Unit, or 250 kg of body mass, with masses used to convert numbers of animals into LHUs as shown in Boone and BurnSilver (2002: 6).)

depicted in figures 2 and 3 because they are based on an aggregate measure of wildlife – they contain, but are not limited to, elephant abundance. Two results follow from figure 4. First, it is clear (and unsurprising) that the conservation effects of PES are attenuated when the Maasai convert all payments into livestock – the dashed curve is below the solid curve. Livestock demand for forage exceeds the carrying capacity by some 20 per cent, and overgrazing and competition for food forces the wildlife population down. In particular smaller-bodied herbivores showed such compensatory changes in abundance in response to a rapid increase in livestock stocking (elephants are less sensitive).

Second, and more interestingly, upon comparing the new scenario where PES payments are used to buy livestock to the control scenario without PES it is evident that it is difficult to unambiguously rank the scenarios in terms of conservation effects. There are periods where the wildlife populations with PES are smaller than those occurring in the control case with farming and fences. Throughout the 1990s this situation reverses, and the conservation effects of PES are positive. The reason for the ambiguity is that PES pushes both the extensive and intensive margin of herding. The extensive margin is pushed out as more rangeland is made available, but the intensive margin shifts simultaneously as Maasai increase their stocking rates. The net effect on the availability of food for wildlife is ambiguous, but will be determined by the *relative price* of livestock. If this price is high

(relative to the PES payment), pastoralists respond by modestly increasing their stocking rates, and the extensive margin effect dominates. However, as the livestock price becomes sufficiently low (or as the payments translates into a sufficiently large number of new livestock heads), the gains from extra rangeland are dissipated through the losses from extra competition for food. In the absence of information on relative prices (context-specific) and a better understanding of the pastoralists' objective function it is hard to predict the outcome of PES systems. This is an area worthy of more research.

## **5. Discussion and conclusion**

This paper explores the opportunities for implementing a payment for ecosystem services scheme on Maasai group ranches near Amboseli NP. Wildlife migrates seasonally in and out of the park, and conserving wildlife in a sustainable fashion implies securing land use types outside the reserve that are compatible with wildlife. Livestock grazing is an example of such a compatible land use type. Fenced-in cropping is not. Due to the many and potentially complex inter-linkages between human and natural systems it is imperative to analyze these issues with a model that integrates insights from ecology and economics.

PES is an increasingly popular instrument for promoting conservation, especially in Europe. In recent years, PES has been introduced in developing countries, in particular in the context of watershed management and carbon storage. However there is no reason to discount the potential use of PES as a mechanism to align potentially opposing interests in the area of wildlife management or biodiversity conservation (areas where non-tangible non-use values are likely important – spilling over national boundaries). We conclude that PES may be a powerful tool in the Amboseli ecosystem because it promotes conservation and contributes to alleviation of poverty (also through stabilization of pastoral income). Nevertheless, the study area displays considerable heterogeneity in terms of the opportunity costs of conservation (i.e. foregone returns from cropping), and perhaps a PES scheme is unlikely to swing Maasai behavior in those irrigated horticulture areas where the returns to agriculture are very high.

Moreover, and interestingly, the basic behavioral model that we employ (PHEWS) suggests that these beneficial effects seem to mutually enforce each other: there is no tradeoff between making the Maasai less poor and protecting elephants. Our analysis also indicates that the cost per household per year in Europe or the USA to support a PES that conserves elephants is modest. One important caveat is the potential issue of 'leakage'. If we use a simple mechanical rule to describe Maasai behavior (i.e. 'use all extra funds to purchase extra livestock'), then much of the gains from habitat expansion are dissipated through extra competition for food between livestock and wildlife – this is clearly an issue that needs to be explored in more detail.

One final issue remains – how should the PES project be funded? In light of the very significant non-use values that represent an ongoing flow of benefits accruing to the world population, the effort should be to implement a matching flow of sustained compensation flowing from North to South. Fortunately matters appear relatively simple for the case of the Amboseli

ecosystem, which is a very popular tourist destination. With 200,000 tourist days a year, the PES program could be easily funded with a relatively minor increase in the Park entrance fee – from US\$30 to US\$31 – or with the introduction of a modest bed tax. Having visitors pay for conservation implies that non-visitors are free riding, and receive their non-use values at zero cost. Clearly such free rides are not always feasible elsewhere.

## References

- Adams, W., R. Aveling, D. Brockington, B. Dickson, J. Elliott, J. Hutton, D. Roe, B. Vira, and W. Wolmer (2004), 'Biodiversity, conservation and the eradication of poverty', *Science* **306**: 1146–1149.
- Adams, W.M. and D. Hulme (2001), 'If community conservation is the answer in Africa, what is the question?', *Oryx* **35**: 193–200.
- Bandara, R. and C. Tisdell (2004), 'The net benefit of saving the Asian elephant: a policy and contingent valuation study', *Ecological Economics* **48**: 93–107.
- Barbier, E.B. (1992), 'Community based development in Africa', in E. Barbier and T. Swanson (eds), *Economics for the Wilds*, Washington, DC: Island Press, pp. 15–33.
- Barrett, C.B. and P. Arcese (1995), 'Are integrated conservation development projects sustainable? On the conservation of large mammals in sub-Saharan Africa', *World Development* **23**: 1073–1084.
- Barrett, C.B. and P. Arcese (1998), 'Wildlife harvesting in integrated conservation and development projects: linking harvest to household demand, agricultural production and environmental shocks in the Serengeti', *Land Economics* **74**: 449–465.
- Blanc, J.J., C.R. Thouless, J.A. Hart, H.T. Dublin, I. Douglas-Hamilton, C.G. Craig, and R.F.W. Barnes (2003), 'African Elephant Status Report 2002: an update from the African Elephant Database', Occasional Paper of the IUCN Species Survival Comm. No. 29, Gland, Switzerland and Cambridge, UK: IUCN, UNEP.
- Boone, R.B. (2005), 'Quantifying changes in vegetation in shrinking grazing areas', *Conservation and Society* **3**: 150–173.
- Boone, R.B. and S.B. BurnSilver (2002), 'Integrated assessment results to aid policy decisions in Ngorongoro Conservation Area, Tanzania', Global Livestock Collaborative Research Support Program, POLEYC Project, University of California, Davis.
- Boone, R.B., S.B. BurnSilver, P.K. Thornton, J.S. Worden, and K.A. Galvin (2005), 'Quantifying declines in livestock due to land subdivision', *Rangeland Ecology and Management* **58**: 523–532.
- Boone, R.B., M.B. Coughenour, K.A. Galvin, and J.E. Ellis (2002), 'Addressing management questions for Ngorongoro Conservation Area using the Savanna Modeling System', *African Journal of Ecology* **40**: 138–150.
- Boone, R.B., K.A. Galvin, M.B. Coughenour, J.W. Hudson, P.J. Weisberg, C.H. Vogel, and J.E. Ellis (2004), 'Ecosystem modeling adds value to a South African climate forecast', *Climatic Change* **64**: 317–340.
- Brandon, K., K.H. Redford, and S.E. Sanderson (1998), *Parks in Peril: People, Politics, and Protected Areas*, Washington, DC: Island Press.
- Bruner, A.G., R.E. Gullison, R.E. Rice, and G.A.B. da Fonseca (2001), 'Effectiveness of parks in protecting tropical biodiversity', *Science* **291**: 125–128.
- Buckley, D.J., M.B. Coughenour, C.B. Blyth, D.J. O'Leary, and J.A. Bentz (1993), 'Ecosystem management model – Elk Island National Park: a case study of integrating environmental models with GIS', Proceedings of the 2nd Int. Conf. Integrating GIS and Environ. Mod., National Center for Geographic Information Analysis, University of California, Santa Barbara, CA, USA.

- BurnSilver, S.B. (2007), *Economic Strategies of Diversification and Intensification among Maasai Pastoralists: Changes in Landscape Use and Movement Patterns in Kajiado District, Kenya*, Graduate Degree Program in Ecology, Colorado State University, Fort Collins (in press).
- Campbell, D.J., H. Gichohi, A. Mwangi, and L. Chege (2000), 'Land use conflicts in S.E. Kajiado District, Kenya', *Land Use Policy* **17**: 339–348.
- Campbell, D.J., H. Gichohi, R. Reid, A. Mwangi, L. Chege, and T. Sawin (2002), 'Competition and conflict between people and wildlife in S.E. Kajiado District, Kenya', Working Paper Number 18, ILRI, Nairobi ([www.lucideastafrica.org](http://www.lucideastafrica.org)).
- Campbell, D.J., D.P. Lusch, T. Smucker, and E.E. Wangui (2003), 'Land use change patterns and root cause in the Loitokitok Area, Kajiado district, Kenya', Working Paper Number 19, ILRI, Nairobi ([www.lucideastafrica.org](http://www.lucideastafrica.org)).
- Coughenour, M.B. (1985), 'Graminoid responses to grazing by large herbivores: adaptations, expectations and interacting processes', *Annals of the Missouri Botanical Garden* **72**: 852–863.
- Coughenour, M.B. (1992), 'Spatial modelling and landscape characterization of an African pastoral ecosystem: a prototype model and its potential use for monitoring drought', in D. McKenzie, D. Hyatt, and V. McDonald (eds), *Ecological Indicators*, Vol. 1, New York: Elsevier, pp. 787–810.
- Coughenour, M., R. Boone, P. Thornton, and R. Howe (2002), 'Integrated modeling and assessment of changing pastoral ecosystems – experience in East Africa, and potential for application in Mongolia', in T. Chuluun and D. Ojima (eds), *Fundamental issues affecting sustainability on the Mongolian steppe*, Ulaanbaatar, Mongolia: International Institute for Study of Nomadic Civilization, pp. 313–326.
- Damania, R., R. Stringer, K.U. Karanth, and B. Stith (2003), 'The economics of protecting tiger populations: linking household behavior to poaching and prey depletion', *Land Economics* **79**: 198–216.
- De Leeuw, J., H.H.T. Prins, E.C. Njuguna, M.Y. Said, and R.A. de By (1998), *Interpretation of DRSRS Animal Counts (1977–1997) in the Rangeland Districts of Kenya*, Enschede, The Netherlands: International Institute for Aerospace Survey and Earth Sciences.
- Ellis, J. and M.B. Coughenour (1998), 'The SAVANNA integrated modeling system', in V. Squires and A. Sidahmed (eds), *Drylands: Sustainable Use of Rangelands in the Twenty-First Century*, IFAD Technical Reports, Rome, pp. 97–106.
- Felder, S. (1993), 'Unilateral CO<sub>2</sub> reductions and leakage: the consequences of international trade in oil and basic materials', *Journal of Environmental Economics and Management* **25**: 162–176.
- Hemp, A. (2005), 'Climate change-driven forest fires marginalize the impact of ice cap wasting on Kilimanjaro', *Global Change Biology* **11**: 1013–1023.
- Hoare, R.E. (1999), 'Determinants of human – elephant conflict in a land–use mosaic', *Journal of Applied Ecology* **36**: 689–700.
- Kellert, S.R., J.N. Mehta, S.A. Ebbin, L.L. Lichtenfeld (2000), 'Community natural resource management: promise, rhetoric, and reality', *Society and Natural Resources* **13**: 705–715.
- Kinyamario, J.I. (1996), 'NPP Grassland: Nairobi, Kenya, 1984–1994', Oak Ridge National Laboratory, Distributed Active Archive Center, Oak Ridge, TN: Available at: <http://www.daac.ornl.gov>
- Landell-Mills, N. and I. Porras (2002), *Silver Bullet or Fools' Gold?* London: International Institute for the Environment and Development (IIED).
- Loomis, J. and D. White (1996), 'Economic benefits of rare and endangered species: summary and meta-analysis', *Ecological Economics* **18**: 197–206.
- Margules, C. and R. Pressey (2000), 'Systematic conservation planning', *Nature* **405**: 243–253.

- Mburu, J., R. Birner, and M. Zeller (2003), 'Relative importance and determinants of landowners' transaction costs in collaborative wildlife management in Kenya: an empirical analysis', *Ecological Economics* **45**: 59–73.
- MEA (Millennium Ecosystem Assessment) (2005), *Ecosystems and Human Well-being: Synthesis*, Washington, DC: Island Press.
- Murombedzi, J.C. (1999), 'Devolution and stewardship in Zimbabwe's CAMPFIRE programme', *Journal of International Development* **11**: 287–293.
- Norton-Griffiths, M. and C. Southey (1995), 'The opportunity costs of biodiversity conservation in Kenya', *Ecological Economics* **12**: 125–139.
- Oosterheld, M., C. DiBella, and H. Kerdules (1998), 'Relation between NOAA-AVHRR satellite data and stocking rate of rangelands', *Ecological Applications* **8**: 207–212.
- Pagiola, S., A. Arcenas, and G. Platais (2005), 'Can payments for environmental services help reduce poverty? An exploration of the issues and the evidence to date from Latin America', *World Development* **33**: 237–253.
- Reid, R., L. Gachimbi, J. Worden, D. Wangui, S. Mathai, S. Mugatha, D. Cambell, J. Maitima, B. Butt, H. Gichohi, and E. Ogot (2004), *Linkages between Changes in Land Use, Biodiversity and Land Degradation in the Loitokitok Area of Kenya*, Lucid Working Paper Series Number 49, Nairobi: ILRI ([www.lucideastafrica.org](http://www.lucideastafrica.org)).
- Rykiel, E.J. (1996), 'Testing ecological models: the meaning of validation', *Ecological Modelling* **90**: 229–244.
- Simpson, D. (1995), *Why Integrated Conservation and Development Projects May Achieve Neither Goal*, Resources for the Future Discussion Paper 95–2, Washington, DC: Resources for the Future.
- Skarpe, C., I. Jansson, L. Seljeli, R. Bergstrom, and E. Roskaft (2007), 'Browsing by goats on three spatial scales in a semi-arid savanna', *Journal of Arid Environments* **68**: 480–491.
- Thornton, P.K., S.B. BurnSilver, R.B. Boone, and K.A. Galvin (2006), 'Modeling the impacts of group ranch subdivision on agro-pastoral households in Kajiado, Kenya', *Agricultural Systems* **87**: 331–356.
- Thornton, P.K., R.H. Fawcett, K.A. Galvin, R.B. Boone, J.W. Hudson, and C.H. Vogel (2004), 'Evaluating management options that use climate forecasts: modeling livestock production systems in the semi-arid zone of South Africa', *Climate Research* **26**: 33–42.
- Thornton, P., K. Galvin, and R. Boone (2003), 'An agro-pastoral household model for the rangelands of East Africa', *Agricultural Systems* **76**: 601–622.
- Tisdell, C.A. (1995), 'Issues in biodiversity conservation including the role of local communities', *Environmental Conservation* **22**: 216–228.
- Western, D. (2001), 'Taking the broad view of conservation: a response to Adams and Hulme', *Oryx* **35**: 201–203.
- Wunder, S. (2005), *Payments for Ecosystem Services: Some Nuts and Bolts*, Bogor: CIFOR Occasional Paper No. 42.
- WWF (World Wide Fund for Nature) (2000), *Elephants in the balance: Conserving Africa's Elephants*, [www.panda.org/resources/publications/species/elephants/index.html](http://www.panda.org/resources/publications/species/elephants/index.html)