

**SMOKE TAIN: Impacts on the Chemical and  
Microbiological Profile of Grapes and Wine**

by

**Kerry Anita Pinchbeck**

B.Sc., Flinders University

B.Sc. (Hons) The University of Adelaide

A thesis submitted in fulfilment of the requirements for  
the degree of Doctor of Philosophy

The University of Adelaide

School of Agriculture, Food and Wine

April, 2011

---

# TABLE OF CONTENTS

ABSTRACT.....	i
DECLARATION.....	iii
STATEMENT OF THE CONTRIBUTIONS OF JOINTLY AUTHORED PAPERS.....	iv
ACKNOWLEDGEMENTS.....	viii
<b>CHAPTER 1: INTRODUCTION.....</b>	<b>1</b>
1.1 Introduction to the Australian wine industry.....	2
1.2 History of vineyard exposure to smoke.....	3
1.3 The effects of smoke on nature.....	8
1.4 Composition of smoke.....	10
1.5 Smoke taint in grapes and wine.....	13
1.6 Research aims.....	19
<b>CHAPTER 2: SYNTHESIS OF GUAIACOL GLUCOSIDES.....</b>	<b>21</b>
2.1 Introduction to glycosides in grapes and wine.....	22
2.2 Introduction to guaiacol $\beta$ -D-glucopyranoside.....	24
2.3 Introduction to glycosylation.....	25
2.4 Glycosylation of guaiacol.....	26
2.5 Results and discussion.....	27
2.5.1 Preparation of guaiacol $\beta$ -D-glucopyranoside.....	27
2.2.5.1 Preparation of guaiacol glucoside using 2,3,4,6-tetra-O-acetyl- $\alpha$ -D-glucopyranosyl bromide (Method 1) .....	27
2.5.1.2 Preparation of guaiacol glucoside using 2,3,4,6-tetra-O-pivaloyl- $\alpha$ -D-glucopyranosyl bromide. (Method 2).....	27
2.5.2 Preparation of deuterated guaiacol $\beta$ -D-glucopyranoside.....	28

---

2.6 Materials and methods.....	30
2.6.1 Solvents and reagents.....	30
2.6.2 Chromatography.....	30
2.6.3 Nuclear magnetic resonance (NMR) spectroscopy.	30
2.6.4 Ultra violet/visible spectroscopy and fluorescence spectroscopy.....	31
2.6.5 Microwave synthesis.....	31
2.6.6 Gas chromatography-mass spectrometry (GC-MS)	31
2.6.7 High performance liquid chromatography tandem mass spectrometry (HPLC-MS/MS).....	32
2.6.8 Synthesis.....	33
2.7 Conclusion.....	42
<b>CHAPTER 3: PROVENENCE OF GUAIACOL GLUCOSIDE IN SMOKE AFFECTED FRUIT.....</b>	<b>43</b>
Paper 1: Identification of a $\beta$ -D-glucopyranoside precursor to guaiacol in grape juice following grapevine exposure to smoke.	45
<b>CHAPTER 4: QUANTIFICATION OF GUAIACOL GLYCOSIDES IN SMOKE AFFECTED FRUIT.....</b>	<b>51</b>
Paper 2: Quantitative analysis of glycoconjugate precursors of guaiacol in smoke-affected grapes using liquid chromatography-tandem mass spectrometry based stable isotope dilution analysis.....	53
<b>CHAPTER 5: QUANTIFICATION OF GUAIACOL GLYCOCONJUGATES IN GRAPES AND WINE.....</b>	<b>59</b>
5.1. Introduction.....	60
5.2. Results and discussion.....	61
5.2.1 Method development.....	61
5.2.1.1 Calibration function for guaiacol $\beta$ -D-glucopyranoside.....	61
5.2.1.2 Mass transitions used for HPLC-SRM analysis.....	62

---

5.2.2 Method validation.....	63
5.2.2.1 Instrument repeatability.....	63
5.2.2.2 Reproducibility.....	63
5.2.2.3 Recovery.....	63
5.2.3 Application of wine based SIDA method to winemaking trials.....	66
5.2.3.1 Hydrolysis of guaiacol glycoconjugates during fermentation.....	66
5.2.3.2 Influence of winemaking techniques on the glycoconjugate content of wine.....	69
5.2.3.3 Glycoconjugate content of wine and potential for smoke taint to intensify with bottle age.....	72
5.2.3.4 Potential for carryover of glycoconjugates between growing seasons.....	74
5.3 Materials and methods.....	76
5.3.1 Method development.....	76
5.3.1.1 Preparation of wine samples for HPLC-MS/MS analysis.....	76
5.3.1.2 Calibration function for guaiacol $\beta$ -D-glucopyranoside.....	77
5.3.1.3 Instrumental analysis.....	77
5.3.1.4 High performance liquid chromatography tandem mass spectrometry (HPLC-MS/MS).....	77
5.3.2 Application of the quantitative guaiacol glycoconjugate method to winemaking trials.....	79
5.3.2.1 Smoke affected grapes.....	79
5.3.2.2 Winemaking.....	80
5.3.3 Statistical analysis.....	81
5.4. Conclusion.....	82
<b>CHAPTER 6: THE EFFECT OF WINEMAKING TECHNIQUES ON THE INTENSITY OF SMOKE TAIN IN WINE.....</b>	<b>83</b>

---

**CHAPTER 6: THE EFFECT OF WINEMAKING TECHNIQUES ON THE INTENSITY OF SMOKE TAIN IN WINE..... 83**

Paper 3: The effect of winemaking techniques on the intensity of smoke taint in wine.....	85
<b>CHAPTER 7: IMPACT OF SMOKE ON GRAPE BERRY MICROFLORA AND YEAST FERMENTATION.....</b>	<b>97</b>
Paper 4: Impact of smoke on grape berry microflora and yeast fermentation.....	99
<b>CHAPTER 8: SUMMARY.....</b>	<b>104</b>
APPENDIX .....	109
REFERENCES.....	117



## ABSTRACT

Guaiacol  $\beta$ -D-glucopyranoside was prepared via a modified Koenigs-Knorr glycosylation method as a reference compound to confirm its presence in smoke affected grapes, using high performance liquid chromatography-tandem mass spectrometry (HPLC-MS/MS) analysis. The  $\beta$ -D-glucopyranoside of guaiacol was identified in extracts of Sangiovese grapes exposed to bushfire smoke and Chardonnay grapes exposed to smoke under experimental conditions. Only trace levels of the glucoside were identified in the corresponding control (i.e. unsmoked) Chardonnay grapes, indicating glycosylation of smoke-derived guaiacol occurred in response to smoke exposure. The reference compound and the glucoside present in smoked juice samples remained largely unaffected following strong acid hydrolysis but were highly susceptible to  $\beta$ -glucosidase enzyme hydrolysis, providing a plausible explanation for the release of guaiacol during fermentation of smoke affected grapes.

Following the identification of additional guaiacol glycoconjugate precursors, the  $d_4$ -labelled analogue of guaiacol  $\beta$ -D-glucopyranoside was synthesised for use as an internal standard in the development of a quantitative stable isotope dilution analysis (SIDA) method, using HPLC-MS/MS. This method was subsequently applied to the analysis of several grape varieties exposed to either experimental or bushfire smoke, to investigate the accumulation of guaiacol glycoconjugates following grapevine smoke exposure. Experimentally smoked grapes contained glycoconjugate concentrations up to 300  $\mu\text{g}/\text{kg}$ ; whereas grapes affected by bushfire smoke contained up to 2,000  $\mu\text{g}/\text{kg}$  glycoconjugates, attributed to the different durations of smoke exposure.

Analysis of separated berry components indicated that the majority of guaiacol glycoconjugates were present in skin and pulp fractions, although approximately 6.7 times higher concentrations were found in the skins by mass. As such,

glycoconjugate extraction from berry homogenate was considered to be more efficient than from juice. To investigate the potential for smoke taint carry over to subsequent growing seasons, grapes were collected from control and smoked Merlot and Viognier grapevines in the season following smoke exposure. Subsequent analysis showed no evidence to suggest grapevine sequestration of glycoconjugates.

The HPLC-MS/MS based SIDA method was adapted for the quantification of guaiacol glycoconjugates in wine and the method applied to several winemaking trials to investigate glycoconjugate metabolism during fermentation. Reduced skin contact achieved using a 'cold soak' winemaking technique, yielded wines with significantly lower concentrations of guaiacol glycoconjugates, compared with traditional red winemaking practices involving extended skin contact at ambient temperature. This suggests winemaking processes which limit precursor extraction might offer opportunities for winemakers to ameliorate the impact of smoke taint in wine. A yeast selection trial demonstrated only partial metabolism of glycoconjugates during fermentation, with glycoconjugate concentrations of finished wines not significantly influenced by choice of yeast strain. The fact that a considerable portion of the glycoconjugate pool remained after fermentation has important implications for industry; i.e. hydrolysis of glycoconjugates after bottling could result in enhanced smoke characters with ageing.

The effect of grapevine smoke exposure on grape berry microflora and the performance of several winemaking yeast in the presence of smoke-derived volatiles were also investigated. The growth of indigenous and winemaking yeast on yeast media agar plates spiked with guaiacol, 4-methylguaiacol or a liquid smoke preparation was investigated to further determine the impact of smoke-derived volatile compounds on yeast performance.

## DECLARATION

This work contains no material which has been accepted for the award of any other degree or diploma in any university or other tertiary institution to Kerry Pinchbeck and, to the best of my knowledge and belief, contains no material previously published or written by another person, except where due reference has been made in the text.

I give consent to this copy of my thesis when deposited in the University Library, being available for loan and photocopying, subject to the provisions of the Copyright Act 1968.

The author acknowledges that copyright of published works contained within the thesis (as listed below) resides with the copyright holder(s) of those works.

I also give permission for the digital version of my thesis to be made available on the web, via the University's digital research repository, the Library catalogue, the Australasian Digital Theses Program (ADTP) and also through web search engines, unless permission has been granted by the University to restrict access for a period of time.

Date 21/4/11

Kerry Pinchbeck

## STATEMENT OF THE CONTRIBUTIONS OF JOINTLY AUTHORED PAPERS

1. Hayasaka, Y. Dungey, K.A. Baldock, G.A. Kennison, K.R. and Wilkinson, K.L. Identification of a  $\beta$ -D-glucopyranoside precursor to guaiacol in grape juice following grapevine exposure to smoke. *Anal. Chim. Acta.*, **2010**, *660*, 143-148.

Author contributions: KLW designed experiments, KAD contributed to the design of experiments and synthesised reference standards, KRK generated control and smoked grape samples through experimental field trials, YH conducted HPLC-MS/MS analysis, GAB conducted GC-MS analysis. All authors contributed to interpretation of data, drafting and/or critical revision of the manuscript.

The following authors agree that the above statement accurately reflects their contribution to this research paper and give consent to its inclusion in this thesis.

..Pinchbeck, K.A. (nee Dungey)

..Hayasaka, Y.

..Wilkinson, K.L.

..Baldock, G.A.

..Kennison, K.R.

2. Dungey, K.A. Hayasaka, Y. Wilkinson, K.L. Quantitative analysis of glycoconjugate precursors of guaiacol in smoke affected grapes using liquid chromatography-tandem mass spectrometry based stable isotope dilution analysis. *Food Chem.* **2011**, 126, 801-806.

Author contributions: KAD and KLW designed experiments and performed field trials, KAD collected and prepared all samples for analysis, YH performed HPLC-MS/MS analysis. All authors contributed to data interpretation, drafting and/or critical revision of the manuscript.

The following authors agree that the above statement accurately reflects their contribution to this research paper and give consent to its inclusion in this thesis.

Pinchbeck, K.A. (nee Dungey)

Hayasaka, Y.

Wilkinson, K.L.

3. Ristic, R. Osidacz, P. Pinchbeck, K.A. Hayasaka, Y. Fudge, A.L. Wilkinson, K.L.  
The effect of winemaking techniques on the intensity of smoke taint in wine. *Aust. J. Grape Wine Res. In Press*, 2011, 17(2).

Author contributions: RR and KLW designed research experiments, RR, KLW and KAP conducted field trials. RR, KAP and ALF conducted winemaking, RR and ALF conducted wine analysis measurements, KAP collected and prepared samples for HPLC-MS/MS analysis, RR and PO performed sensory analysis, YH conducted HPLC-MS/MS analysis. All authors contributed to data interpretation, drafting and/or critical revision of the manuscript.

The following authors agree that the above statement accurately reflects their contribution to this research paper and give consent to its inclusion in this thesis.

..Ristic, R.

..Osidacz, P.

..Pinchbeck, K.A. (nee Dungey)

..Hayasaka, Y.

..Fudge, A.L.

..Wilkinson, K.L.

4. Dungey, K. Grbin, P. Wilkinson, K. Impact of smoke on grape berry microflora and yeast fermentation. *State-of-the-Art in Flavour Chemistry and Biology*. Hofmann, T., Meyerhof, W. and Schieberle, P. Eds; Deutsche Forschungsanstalt für Lebensmittelchemie, Garching (*In Press*)

Author contributions: All authors contributed to the design of research experiments, KD and KW performed field trials, KD collected and analysed samples. All authors contributed to data interpretation, drafting and/or critical revision of the manuscript.

The following authors agree that the above statement accurately reflects their contribution to this research paper and give consent to its inclusion in this thesis.

„Pinchbeck, K. ( nee Dungey)

„Grbin, P.

„Wilkinson, K.

## **ACKNOWLEDGEMENTS**

Throughout the last three years I have been fortunate enough to have a great team of people behind me supporting and helping me the whole way through my study. Firstly many thanks to my principal supervisor Dr. Kerry Wilkinson who convinced me to do a PhD, your guidance and support was crucial to the success of this project and I am grateful to have had such a supporting supervisor. Thankyou also to my supervisors Prof. Dennis Taylor for his help and advice throughout the last three years and Dr. Yoji Hayasaka from the Australian Wine Research Institute, who has enabled all the HPLC-MS analysis for this work, he is truly the master of the HPLC-MS and without his expertise this project would have been a lot more difficult to achieve.

Without valuable funding from the GWRDC this work would not have been conducted so I thank them for their financial support. Also to the Department of Agriculture, Fisheries and Forestry for awarding me the 2009 Science and Innovation award for young people in Agriculture, Fisheries and Forestry and the associated funding that supported the microbiological research to be conducted. The University of Adelaide provided laboratories and facilities for this research for which I am grateful.

I have been fortunate enough to build some important collaborations during my research. Firstly I would like to thank Dr. Alan Pollnitz for his helpful discussions and guidance. Also to Mrs Gayle Baldock from the Australian Wine Research Institute who helped with GC-MS and HPLC-MS analysis.

There are so many people who have helped me with this research, but I have to start by thanking all the members of the Wilkinson group especially Anthea and Renata for their assistance with the challenging field work and great friendship as well. Crista

Burbidge provided me with microbiological technical assistance, without which I would have been lost. Various other people have been essential in helping me with my research and being there as friends, including all the guys and girls in the Taylor group.

Finally and most importantly I need to thank my friends and family for always supporting me with everything I have done. To my parents who helped support me throughout my study, I am so grateful for having those opportunities. My husband Chris you have been there for me when I was stressed and unsure about where I was going, you have pushed me to succeed and I will love you always.