

PUBLISHED VERSION

Burks, Roger A.; Masner, Lubomír; Johnson, Norman Frederick; Austin, Andrew Donald
Taxonomic revision of *Bracalba* Dodd (Hymenoptera, Platygasteridae s.l.), a parasitoid wasp genus
endemic to Australia, *ZooKeys*, 2012; 236:1-53.

Copyright *Roger A. Burks et al.* This is an open access article distributed under the terms of the
[Creative Commons Attribution License 3.0 \(CC-BY\)](https://creativecommons.org/licenses/by/3.0/), which permits unrestricted use, distribution,
and reproduction in any medium, provided the original author and source are credited.

PERMISSIONS

Open Access

<http://www.pensoft.net/page.php?P=2&SESID=0c56c956749477c723152bafac2cf791>



Pensoft adheres to open access as much as possible to accelerate the barrier-free dissemination of scientific knowledge. Several of our journals and e-books can be read, downloaded, printed and distributed at no charge for the reader!

By publishing open access, authors benefit from a higher visibility and increased citation rate of their papers!

All open access papers and news items on this site are licensed under the [Creative Commons Attribution 3.0 \(CC-BY\)](https://creativecommons.org/licenses/by/3.0/).

9th August 2013

<http://hdl.handle.net/2440/75263>

Taxonomic revision of *Bracalba* Dodd (Hymenoptera, Platygasteridae s.l.), a parasitoid wasp genus endemic to Australia

Roger A. Burks^{1,†}, Lubomír Masner^{2,‡}, Norman F. Johnson^{1,§}, Andrew D. Austin^{3,¶}

1 Department of Evolution, Ecology, and Organismal Biology, The Ohio State University, 1315 Kinnear Road, Columbus, OH 43212, U.S.A. **2** Agriculture and Agri-Food Canada, K.W. Neatby Building, Ottawa, ON K1A 0C6, Canada **3** Australian Centre for Evolutionary Biology and Biodiversity, School of Earth and Environmental Sciences, The University of Adelaide, SA 5005, Australia

† [urn:lsid:zoobank.org:author:F792CF45-EB70-458B-85B8-D1AF8A30936F](https://doi.org/urn:lsid:zoobank.org:author:F792CF45-EB70-458B-85B8-D1AF8A30936F)

‡ [urn:lsid:zoobank.org:author:FA505310-F606-4F6C-A1DF-74B9A0055B2E](https://doi.org/urn:lsid:zoobank.org:author:FA505310-F606-4F6C-A1DF-74B9A0055B2E)

§ [urn:lsid:zoobank.org:author:3508C4FF-F027-445F-8417-90AB4AB8FE0D](https://doi.org/urn:lsid:zoobank.org:author:3508C4FF-F027-445F-8417-90AB4AB8FE0D)

¶ [urn:lsid:zoobank.org:author:DE71F924-750D-490D-84A7-F5960066F7CC](https://doi.org/urn:lsid:zoobank.org:author:DE71F924-750D-490D-84A7-F5960066F7CC)

Corresponding author: Roger A. Burks (burks.roger@gmail.com)

Academic editor: M. Buffington | Received 29 May 2012 | Accepted 9 October 2012 | Published 2 November 2012

[urn:lsid:zoobank.org:pub:724528A4-B2C4-4FDB-8702-2734A4DDDC5B](https://doi.org/urn:lsid:zoobank.org:pub:724528A4-B2C4-4FDB-8702-2734A4DDDC5B)

Citation: Burks RA, Masner L, Johnson NF, Austin AD (2012) Taxonomic revision of *Bracalba* Dodd (Hymenoptera, Platygasteridae s.l.), a parasitoid wasp genus endemic to Australia. ZooKeys 236: 1–53. doi: 10.3897/zookeys.236.3434

Abstract

The endemic Australian parasitic wasp genus *Bracalba* (Hymenoptera: Platygasteridae) is revised. Sixteen species are recognized: *B. cuneata* Dodd, *B. laminata* Dodd and *B. nigrescens* (Dodd) are redescribed and thirteen new species are recognized; *B. clavata* Burks, **sp. n.**, *B. globosa* Burks, **sp. n.**, *B. hesperia* Burks, **sp. n.**, *B. intermedia* Burks, **sp. n.**, *B. magnirubra* Burks, **sp. n.**, *B. parvirubra* Burks, **sp. n.**, *B. pinnula* Burks, **sp. n.**, *B. plana* Burks, **sp. n.**, *B. propodealis* Burks, **sp. n.**, *B. sculptifrons* Burks, **sp. n.**, *B. sparsa* Burks, **sp. n.**, *B. tricolorata* Burks, **sp. n.**, and *B. tridentata* Burks, **sp. n.** The genus is found continent-wide but mostly south of the Tropic of Capricorn, and with the highest species diversity occurring in the Pilbara and south-western regions of Western Australia. The hosts of *Bracalba* are unknown but specimens reared from eggs confirm that the genus is associated with orthopteran hosts. A preliminary phylogeny of species did not indicate that species groups were monophyletic, but they are retained despite paraphyly because they are convenient for specimen identification.

Keywords

Platygastroidea, Platygastridae s.l., Scelionidae, *Bracalba*, Scelioninae, key, revision, database, endemism, parasitoid

Introduction

The Platygastroidea are a highly diverse component of the Australian parasitoid wasp fauna with 666 described species in 80 genera (ABRS 2012). However, like most parasitoid families, the real species richness of the group is very much larger than this. The scope of this disparity was recently indicated in a revision of *Idris* (*Ceratobaeus*) (Iqbal and Austin 2000) that resulted in an initial four-fold increase in species to 130+ taxa, and this increase appears to generally hold true for revisions of other platygastroid genera.

The first Australian scelionids were described by Walker (1839) from material collected by Charles Darwin along the east coast of the continent. However, the majority of Australian species (63%) were described by A.P. Dodd between 1913 and 1939 and, testament to his taxonomic abilities, virtually all of his species remain valid, although many have been transferred to other genera (Johnson 1992; ABRS 2012). Other than the taxonomic treatment of a number of small genera by various authors, the most influential post-Dodd studies on the Australian fauna have been the revision of the telenomine genera *Psix* (Johnson and Masner 1985) and *Trissolcus* (Johnson 1991), the spider-associated genera *Baeus* (Stevens and Austin 2007), *Idris* (*Ceratobaeus*) (Iqbal and Austin 2000) and *Mirobaeoides* (Austin 1986), the acridid egg parasitoid genus *Scelio* (Dangerfield et al. 2001), and the generic overview provided by Galloway and Austin (1984). Platygastrids are mostly endoparasitoids of insect and spider eggs, including all taxa previously accommodated in the family Scelionidae, while many platygastroids (*s.str.*) oviposit either in the egg or early larva of gall midges (Cecidomyiidae) and complete their development in the larvae (Austin and Field 1997; Austin et al. 2005).

The current study focuses on one of the few genera, *Bracalba* Dodd, that is endemic to the Australian continent. Among other characters, the genus is recognised by its stout, sculptured body, obviously setose eyes, face with pronounced frontal depression, and very long fore wing postmarginal vein (R1). Previously known from only three species, two from south-eastern Queensland and one from south-western Western Australia, this revision treats 16 species in total. In addition, we document the distribution of all species, present a key to their identification, and undertake a preliminary analysis of species-level relationships.

The contributions of the individual authors are as follows; R.A. Burks: character definition, species concept development; key development, imaging, capture of specimen data, manuscript preparation, phylogenetic analysis and illustration; L. Masner: specimen acquisition, and generic overview; N.F. Johnson: generic concept development and manuscript preparation ; A.D. Austin: initial species concept development, manuscript preparation, and taxonomic overview.

Materials and methods

Specimens examined were provided by the following collections: The American Entomological Institute, Gainesville, Florida, USA (AEIC)¹; Australian Museum, Sydney, Australia (AMSA)²; Australian National Insect Collection, Canberra, Australia (ANIC)³; The Natural History Museum, London, United Kingdom (BMNH)⁴; Canadian National Collection of Insects, Arachnids and Nematodes, Ottawa, Canada (CNCI)⁵; Museum of Victoria, Entomology, Melbourne, Australia (MVMA)⁶; Museum of Comparative Zoology, Harvard University, Cambridge, Massachusetts, USA (MCZC)⁷; C.A. Triplehorn Insect Collection, Ohio State University, Columbus, Ohio (OSUC)⁸; Queensland Primary Industries and Fisheries Insect Collection, Indooroopilly, Australia (QDPC)⁹; Queensland Museum, Brisbane, Australia (QMBA)¹⁰; R.M. Bohart Museum, University of California, Davis, USA (UCDC)¹²; Western Australian Museum, Perth, Australia (WAMP)¹³; Waite Insect and Nematode Collection, Adelaide, Australia (WINC)¹⁴. South Australian and Northern Territory holotypes for species newly described in this manuscript are deposited in the South Australian Museum, Adelaide, Australia (SAMA)¹¹.

This revision is a product of the Platygastroidea Planetary Biodiversity Inventory, funded by the U.S. National Science Foundation (N.F. Johnson, Ohio State University; A.D. Austin, University of Adelaide; Principal Investigators). One objective of this project is to use biodiversity informatics resources to accelerate taxonomic processes, making real-time collaboration possible within a community of researchers. Data associated with specimens examined in this study can be accessed at hol.osu.edu and by entering the unique specimen identifier (e.g. OSUC 238448) in the search form. Morphological terminology follows Mikó et al. (2007), except for: 1) antennal features discussed by Bin (1981), 2) anteclypeus and postclypeus used sensu Dangerfield et al. (2001), 3) dorsal epomial carina and vertical epomial carina, which are mentioned by Masner and Johnson (2007) and defined by Talamas et al. (2011), 4) a newly coined “metasomal bend,” which is a ventrad bend occurring near the midlength of the 4th metasomal segment, and 5) ovipositor features described by Austin and Field (1997). Surface sculptural terms are used in part based on Eady (1968), with general terms added for descriptive purposes: “mesh” to define a polygonal, ovate or circular area enclosed by raised or sunken borders; “septa” to define distinctive borders that enclose meshes; and “interspaces” to define areas separating individual sculptural features. Note that sculptural septa can be differentiated, as in the longitudinal septa being much stronger than the transverse septa. They are also repeated corresponding to the number of sculptural meshes present. The aforementioned terminology is used instead of calling all raised sculpture “carinae,” which would remove all recognizable meaning from the word. Instead, “carina” is used for a raised sculptural element that does not correspond to repeated enclosed meshes. Life science identifiers (LSIDs) can be resolved at <http://lsid.tdwg.org> (i.e. urn:lsid:zoobank.org:act:E5095F58-4603-4796-9D22-03A759B7B29).

Illustration and data citations. Photographs were taken using a Visionary Digital BK+ Imaging System, November 2010 model, with either a K2 Long Distance Microscope or a 65 mm varifocal lens. Source photos were stacked using Zerene Stacker version 1.04 and enhanced using Adobe Photoshop CS5.

Phylogenetic analysis. An exact search using implicit enumeration (branch-and-bound) was performed using TNT (Tree analysis using New Technology) version 1.1 (Goloboff et al. 2003, 2008). Bootstrapping was performed with 1,000 replicates using New Technology searches at an initial level of 95. The outgroup specimens coded in this analysis were from an undescribed species of *Chromoteleia* Ashmead from Belize, (OSUC 064028, OSUC 064021), chosen because of the overall morphological similarity between *Chromoteleia* and *Bracalba*. A total of 42 characters were used, of which 38 were parsimony informative (see Appendix II for characters and matrix).

Taxonomy

Bracalba Dodd

urn:lsid:zoobank.org:act:E5095F58-4603-4796-9D22-03A759B7B29

urn:lsid:biosci.ohio-state.edu:osuc_concepts:457

<http://species-id.net/wiki/Bracalba>

Bracalba Dodd 1931: 78 (original description. Type: *Bracalba laminata* Dodd, by original designation); Muesebeck and Walkley 1956: 336 (citation of type species); Masner 1976: 22, 23 (description, key to separate *Baryconus* Förster, *Bracalba* Dodd, *Chromoteleia* Ashmead, *Oxyscelio* Kieffer); Galloway and Austin 1984: 8, 13 (diagnosis, list of species described from Australia, keyed); Johnson 1992: 354 (catalog of world species); Austin and Field 1997: 18, 68 (structure of ovipositor system, discussion of phylogenetic relationships, genus misplaced in Baryconini).

Diagnosis. Eye setose; frontal depression present; antenna 12-segmented; netrion present; postmarginal vein of fore wing present, longer than marginal and stigmal veins; mesotibia and metatibia each with 1 spur; metascutellum setose dorsally and ventrally; ovipositor *Scelio*-type.

Description. Body length: 2.75–6.88 mm (n=78).

Head. Head shape in dorsal view: transverse. Hyperoccipital carina: absent. Occipital carina: present laterally, broadly interrupted medially; present, complete medially. Occipital carina sculpture: crenulate. OOL: lateral ocellus nearly contiguous with inner orbits, OOL < 0.5 OD. Upper frons: convex, without frontal shelf. Scrobe shape: frons with shallow unmarginated depression above antennal foramina. Frons sculpture: areolate rugose, transversely striate within scrobe; areolate rugose,

scrobe sparsely punctate. Submedian carina: absent. Orbital carina: absent. Inner orbits: diverging ventrally. IOS/EH: IOS slightly less than EH. Interantennal process: triangular in lateral view, well-developed. Central keel: absent. Antennal foramen opening: nearly anteriorly. Lower frons striae: absent. Malar sulcus: present. Compound eye size: of normal proportions, not significantly reduced. Compound eye setation: densely setose; sparsely setose. Gena: narrow, weakly convex, receding behind posterior orbit. Clypeus shape: transversely rectangular. Apical margin of clypeus: rounded; with a small median point. Anteclypeus: absent. Postclypeus: absent. Labrum: not visible. Mandible shape: moderate. Mandibular teeth: apex with 2, acute, subequal teeth; apex tridentate, teeth acute, middle tooth distinctly shortest. Arrangement of mandibular teeth: transverse. Number of maxillary palpomeres: 4. Shape of maxillary palpomeres: cylindrical. Number of labial palpomeres: 2.

Antenna. Number of antennomeres in female: 12. Number of antennomeres in male: 12. Insertion of radicle into A1: parallel to longitudinal axis of A1. Shape of A1: more or less cylindrical, not flattened. Length of A3 of female: distinctly longer than A2. Number of clavomeres in female antenna: 8. Claval formula of female antenna: A12-A7/1-2-2-2-2-2; A12-A6/1-2-2-2-2-2-2; A12-A6/1-2-2-2-2-2-1. Arrangement of doubled multiporous plate sensilla on female clava: in longitudinal pairs. Tyloid distribution on male antenna: A5 only. Shape of male flagellum: filiform.

Mesosoma. Mesosoma shape in dorsal view: longer than wide. Mesosoma shape in lateral view: longer than high. Medial portion of transverse pronotal carina: absent. Posterior apex of pronotum in dorsal view: straight, bifid apically to articulate with tegula. Vertical epomial carina: absent. Dorsal epomial carina (lateral portion of transverse pronotal carina of Vilhelmsen et al. (2010)): absent. Central pronotal area: vertical, not visible in dorsal view. Lateral face of pronotum: weakly concave below position of dorsal epomial carina. Netrion: present. Netrion shape: moderately wide, closed ventrally. Anterior portion of mesoscutum: vertical, flexed ventrally to meet pronotum. Mesoscutum shape: pentagonal in outline, posterolateral corner rounded. Skaphion: absent. Notauli: present, percurrent. Parapsidal lines: absent. Admedial lines: absent. Transscutal articulation: well-developed, crenulate. Shape of mesoscutellum: quadrate to trapezoidal. Armature of mesoscutellum: absent. Surface of mesoscutellum: convex throughout. Median longitudinal furrow on mesoscutellum: absent; present. Shape of axillula: small, dorsal margin sinuate. Metascutellum in dorsal view: clearly differentiated. Metascutellar armature: produced into flattened plate. Metascutellar setation: setose dorsally and ventrally. Metapostnotum: not defined externally. Extent of metasomal depression of propodeum: percurrent, extending anteriorly to anterior margin of propodeum. Lateral propodeal projection: well-developed, extending clearly beyond anterior margin of T1. Mesopleural carina: absent or strongly abbreviated, present only near mid coxa. Mesal course of acetabular carina: projecting anteriorly, but too short to intercede between fore coxae. Mesopleural pit: absent. Sternaulus: absent. Posterodorsal corner of mesopleuron: rounded anteriorly.

Legs. Number of mid tibial spurs: 1. Number of hind tibial spurs: 1. Dorsal surface of hind coxa: smooth. Hind tibia shape: cylindrical, ecarinate. Trochantellus: indicated by transverse sulcus on femur.

Wings. Wing development of female: macropterous. Wing development of male: macropterous. Tubular veins in fore wing: present. Bulla of fore wing R: absent. Extent of marginal venation of fore wing: distinct marginal or postmarginal veins developed. Origin of r-rs in fore wing: arising at junction of R/R1 with costal margin. Development of basal vein (Rs+M) in fore wing: nebulous, weakly pigmented. Development of R in hind wing: elongate, extending to costal margin.

Metasoma. Number of external terga in female: 6. Number of external sterna in female: 6. Number of external terga in male: 8. Number of external sterna in male: 7. Shape of metasoma: acuminate, widest submedially. Laterotergites: present, narrow. Laterosternites: present. T1 of female: raised medially into low, rectangular or subelliptical platform, laterally depressed. Relative size of metasomal terga: T2–T4 largest, subequal in size. Terga with basal crenulae: T1–T4. Sublateral carinae on terga: absent. Median longitudinal carina on metasomal terga: absent; present. Distribution of felt fields: absent. Ovipositor type: Scelio-type (Austin and Field 1997).

Etymology. Dodd did not specify the source of the name for this genus, but presumably it is derived the name of the Queensland town of Bracalba. (Dodd also used the names of Merriwa, NSW and Nyleta, QLD for other genera.) He originally combined this name with two species epithets of variable gender, both coined in feminine form: *Bracalba cuneata* and *Bracalba laminata*, clearly indicating that he intended *Bracalba* to be a feminine noun.

Distribution. *Bracalba* has been collected only from Australia (Fig. 1), and has seldom been collected north of the Tropic of Capricorn. The furthest north that any *Bracalba* has been collected was at 24°39'S, and only two species have been found north of Alice Springs, NT. Location records show that the highest species diversity occurs in the Pilbara and south-western regions of Western Australia.

Biology. The hosts of *Bracalba* are unknown but the structure of the ovipositor (*Scelio*-type; Austin and Field 1997) and specimens reared from eggs confirm that the genus is associated with orthopteran hosts. These eggs could not be identified beyond the ordinal level.

Relationships among species. The implicit enumeration search found two optimum trees of 166 steps (strict consensus: Fig. 2). *Bracalba* was monophyletic with respect to the *Chromoteleia* outgroup (bootstrap = 100). Intuitively-based species groups were not monophyletic in the analysis, but this was likely due to homoplasy resulting from inclusion of many sculptural and coloration characters that have mainly descriptive value. Species with a bend at metasomal segment 4 in females did not form a monophyletic group, but this was complicated by the smooth transition in this character from being distinctively present to clearly absent. One feature that was helpful in determining species group, but which was not reported in the key because it was too difficult to accurately and consistently assess, was the lateral margin of the dorsal axillar area. In the *cuneata*-group, the dorsal axillar area was essentially triangular, broadening posteriorly,




Figure 1. Collection events for each *Bracalba* species. * = species known from only one locality.

with the lateral margin sometimes forming a posterior tooth. In the *laminata*-group the dorsal axillar area was generally more nearly semicircular, broadest at midlength, and with the lateral margin not forming a posterior tooth. Apparent variation, and difficulty in observing this character with confidence, caused us to omit it from the key.

Key to species of *Bracalba*

Note: It is imperative to consult figures when using the below key, as shape features of the metascutellum and metasoma are very important in identifying *Bracalba* species with any hope of accuracy.

- 1 Mandibles bidentate with no hint of a third tooth (Fig. 11), teeth narrowly separated and of nearly the same length. Females: T4 without bend or hump (Figs 4, 7); fore wings extending to middle of T5 or beyond when folded over body **2 (*cuneata*-group)**
- Mandibles either tridentate with a small middle tooth (Fig. 40) or very rarely bidentate, but with teeth broadly separated and/or dorsal tooth much longer than ventral tooth (Figs 52, 58). Females: T4 with or without bend or hump (bend: Fig. 33, hump: Fig. 56); fore wings extending only to middle or apex of T4 when folded over body..... **4 (*laminata*-group)**

- 2 Metascutellum flat (Figs 12–13), not projecting dorsally. Male: A3 short: 1.1–1.2× as long as broad (Fig. 13)..... ***Bracalba plana* Burks, sp. n.**
- Metascutellum projecting dorsally (Figs 4, 7). Male: A3 over 1.5× as long as broad..... **3**
- 3 Mesoscutellum sparsely foveolate, with smooth interspaces (Fig. 8). Mesoscutum smoothly and regularly convex anteriorly and posteriorly (Figs 7–8)..
.....***Bracalba globosa* Burks, sp. n.**
- Mesoscutellum densely foveolate, interspaces with irregular longitudinal rugae (Fig. 3). Mesoscutum abruptly convex only anteriorly, flattened or irregularly convex posteriorly (Figs 3–4).....***Bracalba cuneata* Dodd**
- 4 Propodeum with sharp protrusions (from lateral propodeal area) protruding dorsally immediately posterior to metascutellum, reaching at least the same height as the dorsal surface of metascutellum (Figs 47–48). Males unknown
.....***Bracalba propodealis* Burks, sp. n.**
- Propodeum without such protrusions..... **5**
- 5 Metascutellum strongly narrowed apically (Figs 28, 32, 36, 40, 55), tapering to an incised or narrowly truncate apex that is narrower than the maximum metascutellar length; metasomal bend or hump present in females (Figs 33, 56)..... **6**
- Metascutellum with apex broadly truncate, as broad or broader than total metascutellar length (Figs 14, 19, 24, 43, 47, 60); metasomal bend present or absent in females..... **10**
- 6 Mesosoma truncate anteriorly (Figs 28, 32), with antero-lateral pronotal corners protruding anteriorly. Metascutellum incised apically..... **7**
- Mesosoma smoothly rounded anteriorly (Fig. 36), with antero-lateral pronotal corners much weaker, more rounded, and not protruding anteriorly. Metascutellar apex variable **8**
- 7 Mesoscutellum without median carina (Fig. 32). Females with a mostly reddish metasoma and very large body (at least 6.5 mm long).....
.....***Bracalba magnirubra* Burks, sp. n.**
- Mesoscutellum with median carina (Fig. 28). Color of metasoma in females unknown. Female body length 4.25 mm [using Dodd’s (1930) description, which is probably an underestimate].....***Bracalba laminata* Dodd**
- 8 Antennal scrobes crossed by transverse carinae that meet at torular triangle (Fig. 38). Females unknown***Bracalba nigrescens* (Dodd)**
- Antennal scrobes foveolate or smooth. Females with hump near midlength T4 (Figs 41, 56)..... **9**
- 9 Antennal scrobes smooth and distinct. Clypeus with blunt median carina (Fig. 42). Female: S6 without apical incision (Fig. 42); T5, S5 with many strong longitudinal carinae (Figs 40, 42).....***Bracalba parvirubra* Burks, sp. n.**
- Antennal scrobes foveolate and indistinct (Fig. 57). Clypeus without median carina (Fig. 58). Female: S6 with strong apical incision (Fig. 59); T5, S5 with at most a median carina (Figs 55, 59) ***Bracalba sparsa* Burks, sp. n.**

- 10 Frontal depression with strong longitudinal carina that is crossed by a strong transverse carina dorsally. Clypeus with median carina and transverse dorsal carina (Fig. 53). Metascutellum broadly trapezoidal (Fig. 51). Females: metasoma reddish, T5 and S5 with many longitudinal carinae, S6 without apical notch (Fig. 54).....***Bracalba sculptifrons* Burks, sp. n.**
- Frontal depression without such a set of carinae. Clypeus with weaker or no carinae (Figs 20, 45, 64). Metascutellum variable (Figs 19, 62). Females: metasoma brown to black, T5, S5, and S6 variable.....**11**
- 11 Metascutellum elongate-trapezoidal with concave lateral margins (Fig. 62). Female: S6 with strong apical notch (Fig. 65); T5, S5 with distinct longitudinal carinae (Figs 62, 65). Males unknown.....***Bracalba tridentata* Burks, sp. n.**
- Metascutellum broadly trapezoidal with straight lateral margins (Figs 19, 43, 60), or subrectangular to weakly convex with irregular lateral margins (Figs 14, 24). Female: S6 without apical notch (Figs 17, 23, 46); T5, S5 variable.....**12**
- 12 Mesoscutellum with a set of incomplete median longitudinal grooves (Fig. 19). Middle mandibular tooth almost as long as the other two (Fig. 20). Clypeus with weak median carina. Female: S5 with longitudinal carinae (Fig. 23). Males unknown.....***Bracalba hesperia* Burks, sp. n.**
- Mesoscutellum without median grooves (Fig. 43). Middle mandibular tooth tiny, much smaller than the other two (Figs 16, 26, 45). Clypeus without median carina. Female: S5 with or without carinae.....**13**
- 13 Mesoscutellum sparsely foveolate, with broad interspaces (Fig. 60). Female: S5 with longitudinal carinae (Fig. 61).....***Bracalba tricolorata* Burks, sp. n.**
- Mesoscutellum densely foveolate (interspaces very narrow) except sometimes a small smooth antero-median area (Figs 14, 24, 43). Female: S5 finely foveolate, without longitudinal carinae (Figs 17, 46).....**14**
- 14 Metascutellum trapezoidal in dorsal view, with straight lateral margins (Fig. 43). Female: Antenna with seven apical segments bearing large ventral sensilla, forming a more or less distinct 7-segmented club that includes A6***Bracalba pinnula* Burks, sp. n.**
- Metascutellum in dorsal view irregularly subrectangular or cushion-like, with irregular margins (Figs 14, 24). Female: Antenna with large ventral sensilla on only the six apical segments, these expanded into a 6-segmented club that is distinct from A6 (Figs 14–15)**15**
- 15 Mesoscutellum with fine wrinkles or carinae in interspaces between foveolae (Fig. 14). Female: T6 broader than long (Fig. 14)***Bracalba clavata* Burks, sp. n.**
- Mesoscutellum with smooth interspaces between foveolae (Fig. 24); occasionally smaller foveolae may be present in the interspaces, or the interspaces themselves could be raised to form large wrinkles. Female: T6 longer than broad (Fig. 24).....***Bracalba intermedia* Burks, sp. n.**




Figure 2. Strict consensus phylogram of two most parsimonious trees for species of *Bracalba* using implicit enumeration (branch-and-bound exact search), score = 166. Bootstrap support values found using TNT new technology search (set initial level = 95). Bremer support values above 1, and bootstrap values about 50% indicated on branches.

Bracalba cuneata Species Group

Bracalba cuneata Dodd

urn:lsid:zoobank.org:act:10EC007E-A303-4847-BDD9-29AF68935211

urn:lsid:biosci.ohio-state.edu:osuc_concepts:4126

http://species-id.net/wiki/Bracalba_cuneata

Figures 3–6; Morphbank¹⁵

Bracalba cuneata Dodd 1931: 80 (original description); Galloway 1976: 87 (type information); Galloway and Austin 1984: 99 (figure of antenna); Johnson 1992: 354 (catalogued).

Description. *Female.* Body length 3.25–4.00 mm (n=29). Color of antenna beyond radicle: yellowish-brown. Radicle color: same as scape. Number of claval segments with ventral gustatory sensilla: 7. Number of ventral gustatory sensilla on A6: 2.



Figures 3–6. *Bracalba cuneata* Dodd, holotype female (QMBA HY4733). **3** Dorsal habitus **4** Lateral habitus, female (OSUC 238135) **5** Head, anterior view **6** Metasomal sterna 5–6, ventral view. Morphbank¹⁵

Ocular setae: short and dense. Frontal depression: smooth dorsally, ventrally with longitudinal carina and with additional carinae arising from medial margins of antennal foramina. Smooth depression extending dorsolaterally from antennal foramen: present. Dorsal clypeal margin: but without other sculpture. Clypeal median carina: absent. Ventral clypeal margin: with a small median point. Mandibular teeth: two, separated by narrow incision. Smooth area obliquely posterior to lateral ocellus: present. Genal sculpture: reticulate-rugose with strong dorsoventral carinae. Mandibular color: dark basally and at teeth, becoming lighter reddish brown between these areas.

Dorsal pronotal area: not set off by carina ventrally. Anterolateral corner of dorsal pronotal area: truncate anteriorly. Sculpture of posteromedian area of mesoscutum: foveolate with broad longitudinal septa. Lateral margin of dorsal axillar area: triangularly expanded or with posterior tooth, broadest posteriorly. Mesoscutellar sculpture: densely foveolate. Metascutellum in dorsal view: trapezoidal with broad apex. Dorsal surface of metascutellum: apex protruding dorsally. Femoral depression: crossed by rounded carinae. Anterior corner of lateral propodeal carina: flat, without tooth. Posteromedial corner of lateral propodeal area: protruding posteriorly. Leg color: yellowish-brown except for dorsal external part of coxae and sometimes tarsomeres 2-5.

Metasoma color: black to dark reddish brown. Median lobe of T1: with 7 or more longitudinal carinae, or with median smooth area interrupting carinae. Metasoma at middle of T4: without bend. T5 median carina: absent. Longitudinal sculptural septa

on T5: weak, blunt and hardly raised. Transverse sculptural septa on T5: about as strong as the longitudinal septa. T5 setae: directed posteriorly, arising from anterior edge of sculptural mesh. T6: as broad or broader than long. T6 laterotergite: overlapping S6. S4 median carina: absent. Transverse sculptural septa on S5: weak or absent, much weaker than the longitudinal septa. S5 setae: directed posteriorly, arising from anterior edge of sculptural mesh. Lateral carinae of S6: absent. Apex of S6: without notch.

Male. Body length 3.12–3.80 mm (n=15). Flagellomere length: A3 over 1.5× as long as broad, most others as long or longer than broad. T7: flat and posteriorly truncate.

Diagnosis. *Female.* A6 with 2 large ventral sensilla; 2 mandibular teeth; antero-lateral corner of dorsal pronotal area truncate anteriorly; metascutellum trapezoidal in dorsal view, with a broad apex, protruding dorsally; metasomal bend absent; T4–T6 without median carina, S4–S6 without median carina; T6 broader than long; S6 without apical notch.

Link to distribution map. <http://hol.osu.edu/map-full.html?id=4126>

Associations. Collected near flowering *Scaevola spinescens* R. Br.: [Asterales: Goodeniaceae]

Material examined. Holotype, female: AUSTRALIA: QLD, Chinchilla, 8.II.1928, A. P. Dodd, QMBA HY4733 (deposited in QMBA). Paratypes: AUSTRALIA: 2 females, OSUC 238448-238449 (ANIC). Other material: AUSTRALIA: 32 females, 20 males, OSUC 238516 (BMNH); OSUC 238120, 238123-238128, 238130-238153, 238159-238166, 238171-238173, 238176, 238180-238181, 238188, 238201-238202 (CNCI); OSUC 376915 (MCZC); OSUC 238451, QM Reg. No. T35161 (QDPC); OSUC 231779 (WINC).

Comments. *Bracalba cuneata* is the most commonly collected species of its genus. Our concept of this species includes some slight variation in metascutellar length and pronotal sculpture. This variation included many intermediates and did not correlate with other characters or with collection locality.

***Bracalba globosa* Burks, sp. n.**

urn:lsid:zoobank.org:act:68361C59-4610-4DD5-94CB-0BEB86CA2BB1


urn:lsid:biosci.ohio-state.edu:osuc_concepts:302153

http://species-id.net/wiki/Bracalba_globosa

Figures 7–11; Morphbank¹⁶

Description. *Female.* Body length 3.12–3.25 mm (n=2). Color of antenna beyond radicle: entirely dark. Radicle color: lighter than scape. Number of claval segments with ventral gustatory sensilla: 7. Number of ventral gustatory sensilla on A6: 2.

Ocular setae: short and dense. Frontal depression: smooth dorsally, torular triangle foveolate with 1 transverse carina extending laterally from inner margin of antennal foramen. Smooth depression extending dorsolaterally from antennal foramen: present. Dorsal clypeal margin: but without other sculpture. Clypeal median carina: absent. Ventral clypeal margin: with a small median point. Mandibular color: dark basally



Figures 7–11. *Bracalba globosa* sp. n., paratype female (OSUC 148612). **7** Lateral habitus **8** Head and mesosoma, dorsal view **9** Metasomal sterna 5–6, ventral view **11** Head, ventral view, holotype female (OSUC 148701) **10** Head, anterior view. Morphbank¹⁶

and at teeth, becoming lighter reddish brown between these areas. Mandibular teeth: two, separated by narrow incision. Smooth area obliquely posterior to lateral ocellus: present. Genal sculpture: reticulate-rugose without any strong carinae.

Dorsal pronotal area: not set off by carina ventrally. Anterolateral corner of dorsal pronotal area: weakly rounded anteriorly. Sculpture of posteromedian area of mesoscutum: foveolate with slightly stronger longitudinal septa. Lateral margin of dorsal axillar area: triangularly expanded or with posterior tooth, broadest posteriorly. Mesoscutellar sculpture: densely foveolate with slightly stronger longitudinal septa. Metas-

cutellum in dorsal view: trapezoidal with broad apex. Dorsal surface of metascutellum: apex protruding dorsally. Femoral depression: crossed by rounded carinae. Leg color: dark except for tibiae, 2nd trochanter, and tarsomeres 2–5. Anterior corner of lateral propodeal carina: flat, without tooth. Posteromedial corner of lateral propodeal area: protruding posteriorly.

Metasoma color: black to dark reddish brown. Median lobe of T1: with a set of rugae that merge with one another. Metasoma at middle of T4: without bend. Posterolateral margins of metasomal terga: without protrusions. T5 median carina: absent. Longitudinal sculptural septa on T5: weak, blunt and hardly raised. Transverse sculptural septa on T5: about as strong as the longitudinal septa. T5 setae: directed posteriorly, arising from anterior edge of sculptural mesh. T6: as broad or broader than long. T6 laterotergite: overlapping S6. S4 median carina: absent. Transverse sculptural septa on S5: weak or absent, much weaker than the longitudinal septa. S5 setae: directed posteriorly, arising from anterior edge of sculptural mesh. Lateral carinae of S6: absent. Apex of S6: without notch.

Male. Body length 2.88 mm (n=1). Flagellomere length: A3 over 1.5× as long as broad, most others as long or longer than broad. T7: flat and posteriorly truncate.

Diagnosis. *Female.* A6 with 2 large ventral sensilla; 2 mandibular teeth; anterolateral corner of dorsal pronotal area weakly rounded anteriorly; metascutellum trapezoidal with a broad apex, extending dorsally; metasomal bend absent; T4–T6 without median carina, S4–S6 without median carina; T6 broader than long; S6 without apical notch. *Bracalba globosa* is very similar to *B. cuneata*, but differs mainly in mesosomal shape and sculpture. The metascutellum also differs subtly between most specimens of the two species, being smaller in most *B. globosa*. *Bracalba plana* is also very similar to *B. globosa*, but has a very different mesosomal shape.

Etymology. Latin adjective, named after the convex mesoscutum of most specimens in this species.

Link to distribution map. <http://hol.osu.edu/map-full.html?id=302153>

Material examined. Holotype, female: AUSTRALIA: WA, Kalbarri National Park, 12.XII-19.XII.1986, malaise trap/pan trap, J. S. Noyes, OSUC 148701 (deposited in WAMP). Paratype: AUSTRALIA: 1 female, OSUC 148612 (CNCI).

***Bracalba plana* Burks, sp. n.**

urn:lsid:zoobank.org:act:DF77087E-2227-4528-854C-9EE7F7BC7AF1

urn:lsid:biosci.ohio-state.edu:osuc_concepts:302159

http://species-id.net/wiki/Bracalba_plana

Figures 12–13; Morphbank¹⁷

Description. *Female.* Body length 2.75–3.13 mm (n=3). Color of antenna beyond radicle: mostly dark, ventral parts of pedicel apex, A4–A12 variably lighter. Radicle color: lighter than scape. Number of claval segments with ventral gustatory sensilla: 7. Number of ventral gustatory sensilla on A6: 2.



Figures 12–13. *Bracalba plana* sp. n., holotype female (OSUC 230804). **12** Dorsal habitus, paratype male (OSUC 148706) **13** Dorsal habitus. Morphbank¹⁷

Ocular setae: short and dense. Frontal depression: with many irregularly transverse rugae. Smooth depression extending dorsolaterally from antennal foramen: present. Dorsal clypeal margin: absent between antennal foramina. Clypeal median carina: absent. Ventral clypeal margin: with a small median point. Mandibular color: dark basally and at teeth, becoming lighter reddish brown between these areas. Mandibular teeth: two, separated by narrow incision. Smooth area obliquely posterior to lateral ocellus: present. Genal sculpture: deeply reticulate-rugose with some septa much stronger than others, forming distinct rows differing in height.

Dorsal pronotal area: not set off by carina ventrally. Anterolateral corner of dorsal pronotal area: weakly rounded anteriorly. Sculpture of posteromedian area of mesoscutum: densely foveolate. Lateral margin of dorsal axillar area: triangularly expanded or with posterior tooth, broadest posteriorly. Mesoscutellar sculpture: sparsely foveolate, with large smooth interspaces. Metascutellum in dorsal view: semicircular. Dorsal surface of metascutellum: flat. Femoral depression: centrally smooth, peripherally foveolate. Leg color: coxae dark, leg becoming gradually lighter apically. Anterior corner of lateral propodeal carina: flat, without tooth. Posteromedial corner of lateral propodeal area: protruding posteriorly.

Metasoma color: black to dark reddish brown. Median lobe of T1: with 7 or more longitudinal carinae. Metasoma at middle of T4: without bend. Posterolateral margins of metasomal terga: without protrusions. T5 median carina: absent. Longitudinal sculp-

tural septa on T5: weak, blunt and hardly raised. Transverse sculptural septa on T5: about as strong as the longitudinal septa. T5 setae: directed posteriorly, arising from anterior edge of sculptural mesh. T6: as broad or broader than long. T6 laterotergite: overlapped by rim from S6. S4 median carina: absent. Transverse sculptural septa on S5: about as strong as the longitudinal septa. S5 setae: directed posteriorly, arising from anterior edge of sculptural mesh. Lateral carinae of S6: absent. Apex of S6: without notch.

Males. Body length 2.75–3.00 mm (n=2). Flagellomere length: A3 1–1.2× as long as broad, most others transverse. T7: flat and posteriorly truncate.

Diagnosis. *Female.* A6 with 2 large ventral sensilla; 2 mandibular teeth; anterolateral corner of dorsal pronotal area weakly rounded anteriorly; metascutellum trapezoidal with a broad apex, flat dorsally; metasomal bend absent; T4–T6 without median carina, S4–S6 without median carina; T6 broader than long; S6 without apical notch. This species is very similar to *B. globosa*, but differs mainly in having a shorter, more flat metascutellum and shorter flagellomeres in males.

Etymology. Latin adjective, referring to the flat mesoscutum, mesoscutellum, and metascutellum of most specimens of this species.

Link to distribution map. <http://hol.osu.edu/map-full.html?id=302159>

Material examined. Holotype, female: AUSTRALIA: NT, Centre for Zone Research (CSIRO), Alice Springs, XI-1992, malaise trap, Austin & Dangerfield, OSUC 230804 (deposited in SAMA). Paratypes: AUSTRALIA: 5 females, 3 males, OSUC 148706, 238115, 238168, 238170 (CNCI); OSUC 230802-230803, 230805-230806 (WINC).

Comments. Most specimens have a flat mesoscutum and mesoscutellum, but these features were not constant within this species nor in others of the genus.

Bracalba laminata species group

Bracalba clavata Burks, sp. n.

urn:lsid:zoobank.org:act:C75BF55D-ADB0-417C-8AF3-EE4E46370C15

urn:lsid:biosci.ohio-state.edu:osuc_concepts:302166

http://species-id.net/wiki/Bracalba_clavata

Figures 14–18; Morphbank¹⁸

Description. *Female.* Body length 3.25–3.38 mm (n=2). Color of antenna beyond radicle: mostly dark, ventral parts of pedicel apex, A4–A12 variably lighter. Radicle color: base lighter than scape. Number of claval segments with ventral gustatory sensilla: 6. Number of ventral gustatory sensilla on A6: 0.

Ocular setae: short and dense. Frontal depression: with many irregularly transverse rugae, or smooth dorsally, ventrally with oblique carinae converging on a longitudinal ruga. Smooth depression extending dorsolaterally from antennal foramen: present. Dorsal clypeal margin: forming a complete connection between antennal foramina medially. Clypeal median carina: absent. Ventral clypeal margin: with a small median point. Mandibular color: dark basally and at teeth, becoming lighter reddish brown



Figures 14–18. *Bracalba clavata* sp. n., holotype female (OSUC 384555). **14** Dorsal habitus **15** Lateral habitus, paratype female (OSUC 384558) **16** Head, ventral view, paratype female (OSUC 384559) **17** Metasomal sterna 5–6, ventral view, paratype male (OSUC 384556) **18** Dorsal habitus. Morphbank¹⁸

between these areas. Mandibular teeth: three, but middle tooth tiny. Smooth area obliquely posterior to lateral ocellus: present. Genal sculpture: deeply reticulate-rugose with some septa much stronger than others, forming distinct rows differing in height.

Dorsal pronotal area: not set off by carina ventrally. Anterolateral corner of dorsal pronotal area: weakly rounded anteriorly. Sculpture of posteromedian area of mesoscutum: foveolate with broad longitudinal septa. Lateral margin of dorsal axillar area: triangularly expanded or with posterior tooth, broadest posteriorly. Mesoscutellar sculpture: densely foveolate but with smooth central area. Metascutellum in dorsal

view: strongly transverse, subrectangular. Dorsal surface of metascutellum: flat, or convex. Femoral depression: centrally smooth, peripherally foveolate. Leg color: coxae, femora (aside from their apices), and at least the last two tarsomeres dark, otherwise yellowish brown. Anterior corner of lateral propodeal carina: flat, without tooth. Posteromedial corner of lateral propodeal area: not protruding posteriorly.

Metasoma color: black to dark reddish brown. Median lobe of T1: with 7 or more longitudinal carinae. Metasoma at middle of T4: with very weak bend. Posterolateral margins of metasomal terga: with tooth-like protrusions. T5 median carina: absent. Longitudinal sculptural septa on T5: strong, sharply raised. Transverse sculptural septa on T5: about as strong as the longitudinal septa. T5 setae: directed posteriorly, arising from anterior edge of sculptural mesh. T6: as broad or broader than long. T6 laterotergite: overlapping S6. S4 median carina: present. Transverse sculptural septa on S5: weak or absent, much weaker than the longitudinal septa. S5 setae: directed posteriorly, arising from anterior edge of sculptural mesh. Lateral carinae of S6: absent. Apex of S6: without notch.

Male. Body length 2.75–3.25 mm (n=5). Flagellomere length: A3 over 1.5× as long as broad, most others as long or longer than broad. T7: flat and posteriorly truncate.

Diagnosis. *Female.* A6 without large ventral sensilla, separate and much smaller than A7 (therefore only 6 distinct claval segments present); 3 mandibular teeth with the middle tooth much smaller than the others; metascutellum subrectangular and very short; metasomal bend present but very weak; sculpture posterior to metasomal bend with longitudinal and transverse septa of about equal strength and hardly different from those anterior to the bend; T4–T6 without longitudinal carina; S4–S6 with longitudinal carina; T6 about as long as broad; S6 without apical notch. This species is very similar to *B. pinnula* and to some species without a T4 bend. It differs in the unusual antenna, short metascutellum, and in its shorter wings that reach only to the middle of T4.

Etymology. Latin adjective meaning “clavate,” named for the clavate antenna with its distinct separation between the 6-segmented club and the basal flagellomeres.

Link to distribution map. <http://hol.osu.edu/map-full.html?id=302166>

Material examined. Holotype, female: AUSTRALIA: SA, Brookfield Conservation Park, 34°21'S, 139°29'E, 24.XI–26.XI.1992, yellow pan trap, I. Naumann & J. C. Cardale, OSUC 384555 (deposited in ANIC). Paratypes: AUSTRALIA: 2 females, 6 males, OSUC 230820, 367520, 384556, 384558–384559, 384564–384566 (ANIC).

***Bracalba hesperia* Burks, sp. n.**

[urn:lsid:zoobank.org:act:1D575B7A-FD40-4392-B7A5-B0EF403B0168](http://zoobank.org/act:1D575B7A-FD40-4392-B7A5-B0EF403B0168)

[urn:lsid:biosci.ohio-state.edu:osuc_concepts:302154](http://biosci.ohio-state.edu/osuc_concepts:302154)

http://species-id.net/wiki/Bracalba_hesperia

Figures 19–23; Morphbank¹⁹

Description. *Female.* Body length 5.76–6.06 mm (n=2). Color of antenna beyond radicle: entirely dark. Radicle color: same as scape. Number of claval segments with ventral gustatory sensilla: 7. Number of ventral gustatory sensilla on A6: 1.



Figures 19–23. *Bracalba hesperia* sp. n., holotype female (OSUC 148713). **19** Head and mesosoma, dorsal view **20** Head, ventral view **21** Lateral habitus; **22** Metasomal terga 5–6, dorsal view **23** Metasomal sterna 5–6, ventral view. Morphbank¹⁹

Ocular setae: short and sparse. Frontal depression: smooth dorsally, torular triangle foveolate with transverse carinae lateral to this area. Smooth depression extending dorsolaterally from antennal foramen: present. Dorsal clypeal margin: wrinkle-like with a median peak. Clypeal median carina: present. Ventral clypeal margin: with a small median point. Mandibular color: dark basally and at teeth, becoming lighter reddish brown between these areas. Mandibular teeth: three of roughly equal size. Smooth area obliquely posterior to lateral ocellus: present. Genal sculpture: reticulate-rugose without any strong carinae.

Dorsal pronotal area: not set off by carina ventrally. Anterolateral corner of dorsal pronotal area: weakly rounded anteriorly. Sculpture of posteromedian area of mesoscutum: densely foveolate. Lateral margin of dorsal axillar area: with a semi-circular expansion, broadest near midlength. Mesoscutellar sculpture: densely foveolate with one or two median longitudinal channels. Metascutellum in dorsal view: trapezoidal with broad apex. Dorsal surface of metascutellum: convex. Femoral depression: centrally smooth, peripherally foveolate. Leg color: dark except for trochanters, tips of femora and tibiae, and tarsomeres 1–4. Anterior corner of lateral propodeal carina: flat, without tooth. Posteromedial corner of lateral propodeal area: protruding posteriorly.

Metasoma color: black to dark reddish brown. Median lobe of T1: with 7 or more longitudinal carinae. Metasoma at middle of T4: with metasomal bend and abrupt transition in sculpture. Posterolateral margins of metasomal terga: without protrusions. T5 median carina: absent. or present. Longitudinal sculptural septa on T5: weak, blunt and hardly raised. Transverse sculptural septa on T5: about as strong as the longitudinal septa. T5 setae: directed posteriorly, arising from anterior edge of sculptural mesh. T6: longer than broad. T6 laterotergite: overlapped by rim from S6. S4 median carina: absent. Transverse sculptural septa on S5: weak or absent, much weaker than the longitudinal septa. S5 setae: not directed posteriorly, arising from center of sculptural mesh. Lateral carinae of S6: absent. Apex of S6: without notch.

Male. unknown.

Diagnosis. *Female.* A6 with 1 large ventral sensillum; 3 mandibular teeth of approximately equal length; metascutellum broadly trapezoidal with a broad and slightly concave apex; metasomal bend present but weak; sculpture posterior to metasomal bend with transverse septa about as strong as the longitudinal septa; T4–T6 and S4–S6 without median carina; T6 longer than broad; S6 without apical notch. This species is similar to *B. tridentata*, but in that species S6 has a strong apical notch. It is also near *B. nigrescens*, but no complete female specimens of that species are known.

Etymology. Latin adjective, referring to the geographic distribution of this species.

Link to distribution map. <http://hol.osu.edu/map-full.html?id=302154>

Material examined. Holotype, female: AUSTRALIA: WA, 3km W Walpole, Keystone Road, 34°59.01'S, 116°40.76'E, no date, yellow pan trap, George, Hawks & Munro, OSUC 148713 (deposited in WAMP). Paratypes: AUSTRALIA: 2 females, OSUC 148702, 238154 (CNCI).

Comments. Some singleton specimens similar to *B. hesperia* have been examined but left undescribed, most collected from Western Australia. They all exhibit a stronger metasomal bend and stronger longitudinal sculpture posterior to the bend than in *B. hesperia*, but are variable in these and other characters. Together these specimens and *B. hesperia* may represent members of a species complex. Some specimens within this complex possess longitudinal median grooves on the mesoscutellum, but this character may be variable within species.

***Bracalba intermedia* Burks, sp. n.**

urn:lsid:zoobank.org:act:D1FA6338-2827-4D30-AF06-849208F4A6E6

urn:lsid:biosci.ohio-state.edu:osuc_concepts:302155

http://species-id.net/wiki/Bracalba_intermediaFigures 24–27; Morphbank²⁰

Description. *Female.* Body length 3.25–3.88 mm (n=7). Color of antenna beyond radicle: entirely dark. Radicle color: lighter than scape. Number of claval segments with ventral gustatory sensilla: 6. Number of ventral gustatory sensilla on A6: 0.

Ocular setae: short and dense. Frontal depression: medially smooth, torular triangle sparsely foveolate, or foveolate dorsally, torular triangle foveolate, areas lateral to it with transverse carinae. Smooth depression extending dorsolaterally from antennal foramen: absent. Dorsal clypeal margin: bordering antennal foramina, absent between them. Clypeal median carina: absent. Ventral clypeal margin: convex. Mandibular color: dark basally and at teeth, becoming lighter reddish brown between these areas. Mandibular teeth: three, but middle tooth tiny. Smooth area obliquely posterior to lateral ocellus: present. Genal sculpture: reticulate-rugose without any strong carinae.

Dorsal pronotal area: not set off by carina ventrally. Anterolateral corner of dorsal pronotal area: weakly rounded anteriorly. Sculpture of posteromedian area of mesoscutum: foveolate with slightly stronger longitudinal septa. Lateral margin of dorsal axillar area: triangularly expanded or with posterior tooth, broadest posteriorly. Mesoscutellar sculpture: densely foveolate with slightly stronger longitudinal septa. Metascutellum in dorsal view: strongly transverse, subrectangular. Dorsal surface of metascutellum: flat. Femoral depression: centrally smooth, peripherally foveolate. Leg color: coxae, femora (aside from their apices), and at least the last two tarsomeres dark, otherwise yellowish brown. Anterior corner of lateral propodeal carina: flat, without tooth. Posteromedial corner of lateral propodeal area: protruding posteriorly.

Metasoma color: black to dark reddish brown. Median lobe of T1: with a set of rugae that merge with one another. Metasoma at middle of T4: without bend. Posterolateral margins of metasomal terga: without protrusions. T5 median carina: absent. Longitudinal sculptural septa on T5: weak, blunt and hardly raised. Transverse sculptural septa on T5: about as strong as the longitudinal septa. T5 setae: directed posteriorly, arising from anterior edge of sculptural mesh. T6: longer than broad. T6 laterotergite: overlapping S6. S4 median carina: present. Transverse sculptural septa on S5: weak or absent, much weaker than the longitudinal septa. S5 setae: directed posteriorly, arising from anterior edge of sculptural mesh. Lateral carinae of S6: absent. Apex of S6: without notch.

Male. Body length 2.62–3.25 mm (n=18). Flagellomere length: A3 over 1.5× as long as broad, most others as long or longer than broad. T7: flat and posteriorly truncate.

Diagnosis. *Female.* A6 without large ventral sensilla; 3 mandibular teeth with middle tooth much smaller than the others; metascutellum broad and subrectangular,



Figures 24–27. *Bracalba intermedia* sp. n., paratype female (OSUC 180713). **24** Dorsal habitus **25** Ventral habitus, paratype female (OSUC 148707) **26** Head, ventral view, paratype male (OSUC 227576) **27** Dorsal habitus. Morphbank²⁰

very short; metasomal bend absent; T4–T6 without median carina, S4–S6 with at least a vague median carina; T6 longer than broad; S6 without apical notch. The antenna and short metascutellum help distinguish it from similar species. It is very similar to *B. pinnula*, but differs in having a much shorter metascutellum. It is also very similar to *B. clavata*, but has no distinct metasomal bend and has a longer T6 in females.

Etymology. Latin adjective, referring to its unusual mixture of characters.

Link to distribution map. <http://hol.osu.edu/map-full.html?id=302155>

Material examined. Holotype, female: AUSTRALIA: WA, via Dwellingup, Yarragil 4P Catchment, 18.II–25.II.1981, malaise trap, A. Postle, OSUC 230811 (deposited in QMBA). Paratypes: AUSTRALIA: 8 females, 23 males, OSUC 148705, 148707–148708, 180713, 238105–238109, 238111, 238179, 238182, 238184–238186 (CNCI); OSUC 230809–230810, 230812, 230814, 238450 (QDPC); OSUC 227572–227578, 230801, 238454, 238457, 384567 (WINC).

Comments. *Bracalba intermedia* lacks a metasomal bend, but the three mandibular teeth, elongate metasoma, and sculptural features suggest that it belongs in the *laminata* group. Specimens from the eastern localities tend to be slightly smaller and have some reduction in sculpture, but are retained in our concept of this species.

***Bracalba laminata* Dodd**

urn:lsid:zoobank.org:act:4C6F2E7E-3E8C-4973-AC72-D02632852D69

urn:lsid:biosci.ohio-state.edu:osuc_concepts:4127

http://species-id.net/wiki/Bracalba_laminata

Figures 28–31; Morphbank²¹

Bracalba laminata Dodd 1931: 78 (original description); Galloway 1976: 88 (type information); Johnson 1992: 354 (catalogued).

Description. *Female.* Number of claval segments with ventral gustatory sensilla: 7. Number of ventral gustatory sensilla on A6: 2.

Male. Body length = 3.38 mm (n=1). Flagellomere length: A3 over 1.5× as long as broad, most others as long or longer than broad. Ocular setae: long and dense. Frontal depression: densely foveolate. Smooth depression extending dorsolaterally from antennal foramen: present. Dorsal clypeal margin: absent between antennal foramina. Clypeal median carina: absent. Ventral clypeal margin: with a small median point. Mandibular teeth: three, but middle tooth tiny. Smooth area obliquely posterior to lateral ocellus: present. Genal sculpture: deeply reticulate-rugose with some septa much stronger than others, forming distinct rows differing in height. Mandibular color: dark basally and at teeth, becoming lighter reddish brown between these areas.


Dorsal pronotal area: not set off by carina ventrally. Anterolateral corner of dorsal pronotal area: truncate anteriorly. Sculpture of posteromedian area of mesoscutum: foveolate with slightly stronger longitudinal septa. Lateral margin of dorsal axillar area: with a semicircular expansion, broadest near midlength. Mesoscutellar sculpture: densely foveolate with a longitudinal carina. Metascutellum in dorsal view: elongate-trapezoidal but with incised apex. Dorsal surface of metascutellum: apex protruding dorsally. Femoral depression: crossed by rounded carinae. Anterior corner of lateral propodeal carina: flat, without tooth. Posteromedial corner of lateral propodeal area: protruding posteriorly. T7: arched and posteriorly concave.

Diagnosis. Males: Lateral margin of dorsal axillar area semicircularly expanded; mesosoma broadly truncate anteriorly; mesoscutellum with median carina; metascutellum elongate-trapezoidal with incised apex.

Link to distribution map. <http://hol.osu.edu/map-full.html?id=4127>

Material Examined. Holotype, female: AUSTRALIA: QLD, Gogango, 40mi W Rockhampton, III-1928, A. P. Dodd, QMBA HY4732H (deposited in QMBA). Allotype: AUSTRALIA: 1 male, QMBA HY4732A (QMBA). Other material: AUSTRALIA: 3 males, OSUC 365210-365212 (UQIC).

Comments. Only the antenna and a few legs of the female holotype of *B. laminata* remain. The syntopic allotype male strongly resembles the holotype of *B. nigrescens*, but differs in the presence of a partial median carina on the mesoscutellum, the anteriorly truncate mesosoma, and denser sculpturing on the thoracic dorsum. No other female specimens are known aside from the holotype. The allotype shares characteris-



Figures 28–31. *Bracalba laminata* Dodd, allotype male (QMBA HY4732A). **28** Head and mesosoma, dorsal view **29** Head, anterior view **30** Metasomal terga 6–8, posterior view, holotype female (QMBA HY4732H) **31** Antenna, ventral view. Morphbank²¹

tics with *B. hesperia*, *B. magnirubra*, and *B. tridentata*, but differs enough that it cannot definitively be associated with any of these species.

Because the antenna and some legs of the holotype remains, they now represent the holotype. A neotype cannot be properly designated without first requesting that the existing holotype be set aside. This was considered unnecessary, as the antenna clearly belongs to *Bracalba*, based on the number and arrangement of ventral sensilla. Therefore, there is no doubt that the holotype agrees with our concept of *Bracalba*. There is also no reason to conclude that the allotype male would be from a different species than the female, based on its morphology compared with Dodd's (1931) description of the now lost parts of the female's body.

***Bracalba magnirubra* Burks, sp. n.**


urn:lsid:zoobank.org:act:48277EF5-2342-478C-8BDE-3B3F2AF0790E

urn:lsid:biosci.ohio-state.edu:osuc_concepts:302156

http://species-id.net/wiki/Bracalba_magnirubra

Figures 32–35; Morphbank²²

Description. *Female.* Body length 6.50–6.88 mm (n=3). Color of antenna beyond radicle: mostly dark, extreme base of scape becoming reddish. Radicle color: lighter



Figures 32–35. *Bracalba magnirubra* sp. n., paratype female (OSUC 238178). **32** Dorsal habitus **34** Head, ventral view, holotype female (OSUC 148715) **33** Lateral habitus, paratype female (OSUC 238112) **35** Metasomal sterna 5–6, ventral view. Morphbank²²

than scape. Number of claval segments with ventral gustatory sensilla: 7. Number of ventral gustatory sensilla on A6: 1.

Ocular setae: long and sparse. Frontal depression: medially smooth, torular triangle sparsely foveolate. Smooth depression extending dorsolaterally from antennal foramen: present. Dorsal clypeal margin: interrupted by a median areole below interantennal process. Clypeal median carina: present. Ventral clypeal margin: with a small median point. Mandibular color: dark basally and at teeth, becoming lighter reddish brown between these areas. Mandibular teeth: three, but middle tooth tiny. Smooth area obliquely posterior to lateral ocellus: present. Genal sculpture: reticulate-rugose without any strong carinae.

Dorsal pronotal area: not set off by carina ventrally. Anterolateral corner of dorsal pronotal area: protruding anteriorly. Sculpture of posteromedian area of mesoscutum: densely

foveolate. Lateral margin of dorsal axillar area: with a semicircular expansion, broadest near midlength. Mesoscutellar sculpture: densely foveolate. Metascutellum in dorsal view: elongate-trapezoidal but with incised apex. Dorsal surface of metascutellum: apex protruding dorsally, or convex. Femoral depression: crossed by rounded carinae. Leg color: reddish but with at least last two tarsomeres dark. Anterior corner of lateral propodeal carina: flat, without tooth. Posteromedial corner of lateral propodeal area: protruding posteriorly.

Metasoma color: reddish, with last segment variably dark. Median lobe of T1: with 7 or more longitudinal carinae. Metasoma at middle of T4: with metasomal bend and abrupt transition in sculpture. Posterolateral margins of metasomal terga: with tooth-like protrusions. T5 median carina: present. Longitudinal sculptural septa on T5: strong, sharply raised. Transverse sculptural septa on T5: weak or absent, much weaker than the longitudinal septa. T5 setae: not directed posteriorly, arising from center of sculptural mesh. T6: longer than broad. T6 laterotergite: overlapping S6. S4 median carina: present. Transverse sculptural septa on S5: weak or absent, much weaker than the longitudinal septa. S5 setae: not directed posteriorly, arising from center of sculptural mesh. Lateral carinae of S6: not meeting apically. Apex of S6: with notch.

Male. Body length 4.75–5.00 mm (n=5). Flagellomere length: A3 over 1.5× as long as broad, most others as long or longer than broad. T7: arched and posteriorly concave.

Diagnosis. *Female.* A6 with 1 large ventral sensillum; 3 mandibular teeth with middle tooth smaller than the others; metascutellum trapezoidal and relatively long, with a truncate or slightly incised apex; metasomal bend present; sculpture posterior to metasomal bend with strong longitudinal septa but without transverse septa; T4–T6 and S4–S6 with median carina; T6 longer than broad; S6 with apical notch.

Etymology. Compound adjective using the Latin adjectives *magnus* and *ruber*.

Link to distribution map. <http://hol.osu.edu/map-full.html?id=302156>

Material examined. Holotype, female: AUSTRALIA: SA, 32km N Renmark, Amalia Dam, xeric mallee scrub, MT 4, ROM 2000040, Bookmark Biosphere Reserve, 33°53'S, 140°43'E, 263m, 15.II–15.IV.2000, malaise trap, D. C. Darling, OSUC 148715 (deposited in SAMA). Paratypes: AUSTRALIA: 4 females, 6 males, OSUC 365208, 367506–367507 (ANIC); OSUC 238112, 238117–238119, 238121–238122, 238178 (CNCI).

Comments. This is the largest-bodied species of *Bracalba* known. It resembles *B. sculptifrons* and *B. sparsa*, but differs from both in a large number of characters.

Bracalba nigrescens (Dodd)

urn:lsid:zoobank.org:act:3692A700-53E1-4B03-8247-BEE9AC332619

urn:lsid:biosci.ohio-state.edu:osuc_concepts:4128

http://species-id.net/wiki/Bracalba_nigrescens

Figures 36–39; Morphbank²³

Chromoteleia nigrescens Dodd 1920: 329 (original description); Masner 1965: 71 (type information).

Bracalba nigrescens (Dodd): Dodd 1931: 80 (generic transfer); Galloway 1976: 88 (type information); Johnson 1992: 354 (catalogued).

Description. *Male.* Body length 3.38 mm (n=1). Flagellomere length: A3 over 1.5× as long as broad, most others as long or longer than broad. Ocular setae: long and dense. Frontal depression: with many irregularly transverse rugae. Smooth depression extending dorsolaterally from antennal foramen: present. Dorsal clypeal margin: absent between antennal foramina. Clypeal median carina: absent. Ventral clypeal margin: with a small median point. Mandibular teeth: three, but middle tooth tiny. Smooth area obliquely posterior to lateral ocellus: present. Genal sculpture: deeply reticulate-rugose with some septa much stronger than others, forming distinct rows differing in height. Mandibular color: mostly reddish brown, dark at teeth.

Dorsal pronotal area: not set off by carina ventrally. Anterolateral corner of dorsal pronotal area: weakly rounded anteriorly. Sculpture of posteromedian area of mesoscutum: sparsely foveolate, with large smooth interspaces. Lateral margin of dorsal axillar area: with a semicircular expansion, broadest near midlength. Mesoscutellar sculpture: sparsely foveolate, with large smooth interspaces. Metascutellum in dorsal view: elongate-trapezoidal but with incised apex. Dorsal surface of metascutellum: apex protruding dorsally. Femoral depression: crossed by rounded carinae. Anterior corner of lateral propodeal carina: flat, without tooth. Posteromedial corner of lateral propodeal area: protruding posteriorly. Leg color: coxae and at least the last three tarsomeres dark, otherwise yellowish. T7: arched and posteriorly concave.

Diagnosis. Males: Lateral margin of dorsal axillar area semicircularly expanded; mesosoma weakly rounded anteriorly; mesoscutellum without median carina; metascutellum elongate-trapezoidal with incised apex.

Link to distribution map. <http://hol.osu.edu/map-full.html?id=4128>

Material examined. Holotype, male, *C. nigrescens*: **AUSTRALIA:** WA, Yallingup, 1.XII-12.XII.1913, R. E. Turner, B.M. TYPE HYM. 9.499 (deposited in BMNH).

Comments. The male holotype of *B. nigrescens* cannot be associated with any known female *Bracalba*. It is very similar to *B. laminata* even though the two species occur on opposite sides of the Australian continent. These species may be closely related to *B. hesperia* and *B. tridentata*.

***Bracalba parvirubra* Burks, sp. n.**

urn:lsid:zoobank.org:act:9B615296-D426-4E5E-937A-9AC299199003

urn:lsid:biosci.ohio-state.edu:osuc_concepts:302157

http://species-id.net/wiki/Bracalba_parvirubra

Figures 40–42; Morphbank²⁴

Description. *Female.* Body length 3.38–3.63 mm (n=3). Color of antenna beyond radicle: mostly dark, extreme base of scape becoming reddish. Radicle color: same as scape. Number of claval segments with ventral gustatory sensilla: 7. Number of ventral gustatory sensilla on A6: 1.

Ocular setae: long and sparse. Frontal depression: smooth dorsally, torular triangle foveolate, areas lateral to it smooth. Smooth depression extending dorsolaterally from



Figures 36–39. *Bracalba nigrescens* (Dodd) holotype male (B.M. TYPE HYM. 9.499). **36** Head and mesosoma, dorsal view **37** Metasomal terga 6–8, postero-lateral view **38** Head, anterior view **39** Head, ventral view. Morphbank²³

antennal foramen: present. Dorsal clypeal margin: arched, interrupted by broad median carina. Clypeal median carina: present. Ventral clypeal margin: with a small median point. Mandibular color: dark basally and at teeth, becoming lighter reddish brown between these areas. Mandibular teeth: three, but middle tooth tiny. Smooth area obliquely posterior to lateral ocellus: present. Genal sculpture: deeply reticulate-rugose with some septa much stronger than others, forming distinct rows differing in height.

Dorsal pronotal area: not set off by carina ventrally. Anterolateral corner of dorsal pronotal area: protruding anteriorly. Sculpture of posteromedian area of mesoscutum: foveolate with slightly stronger longitudinal septa. Lateral margin of dorsal axillar area: with a semicircular expansion, broadest near midlength. Mesoscutellar sculpture: densely foveolate with one or two median longitudinal channels. Metascutellum in dorsal view: elongate-trapezoidal but with incised apex. Dorsal surface of metascutellum: flat, or convex. Femoral depression: crossed by many sharply defined carinae. Leg color: yellowish-brown except for dorsal external part of coxae and sometimes tarsomeres 2–5. Anterior corner of lateral propodeal carina: flat, without tooth. Posteromedial corner of lateral propodeal area: protruding posteriorly.

Metasoma color: centrally reddish with some vague dark intrusions, peripherally becoming black. Median lobe of T1: with 7 or more longitudinal carinae. Metasoma at



Figures 40–42. *Bracalba parvirubra* sp. n., holotype female (OSUC 238194). **40** Dorsal habitus **41** Lateral habitus **42** Ventral habitus. Morphbank²⁴

middle of T4: with metasomal bend and abrupt transition in sculpture. Posterolateral margins of metasomal terga: with tooth-like protrusions. T5 median carina: absent. Longitudinal sculptural septa on T5: strong, sharply raised. Transverse sculptural septa on T5: about as strong as the longitudinal septa. T5 setae: directed posteriorly, arising from anterior edge of sculptural mesh. T6: as broad or broader than long. T6 laterotergite: overlapped by rim from S6. S4 median carina: present. Transverse sculptural septa on S5: weak or absent, much weaker than the longitudinal septa. S5 setae: directed posteriorly, arising from anterior edge of sculptural mesh. Lateral carinae of S6: forming complete peripheral carina. Apex of S6: without notch.

Male. unknown.

Diagnosis. *Female.* A6 with 1 large ventral sensillum; 3 mandibular teeth with middle tooth much smaller than the others; metascutellum trapezoidal and strongly narrowing to an incised apex; metasomal bend present; sculpture posterior to metasomal bend with strong longitudinal and transverse septa; T4–T6 without distinct median carina; S4–S6 with median carina; T6 about as broad as long; S6 without apical notch. This species is similar to *B. sculptifrons* and *B. sparsa*, but lacks the S6 notch and frontal carinae.

Etymology. Compound adjective using the Latin adjectives parvus and ruber.

Link to distribution map. <http://hol.osu.edu/map-full.html?id=302157>

Material examined. Holotype, female: AUSTRALIA: WA, 20km N Denmark, 16.I.1987, J. S. Noyes, OSUC 238194 (deposited in WAMP). Paratypes: AUSTRALIA: 3 females, OSUC 238191, 238193, 238196 (CNCI).

***Bracalba pinnula* Burks, sp. n.**

urn:lsid:zoobank.org:act:29EE41F2-7AE8-46F5-AE0C-C9250E6A5771

urn:lsid:biosci.ohio-state.edu:osuc_concepts:302158

http://species-id.net/wiki/Bracalba_pinnula

Figures 43–46; Morphbank²⁵

Description. *Female.* Body length 3.37–3.75 mm (n=7). Color of antenna beyond radicle: entirely dark. Radicle color: lighter than scape. Number of claval segments with ventral gustatory sensilla: 7. Number of ventral gustatory sensilla on A6: 1, or 2.

Ocular setae: short and dense. Frontal depression: with many strong transverse carinae, sparsely foveolate at torular triangle. Smooth depression extending dorsolaterally from antennal foramen: present. Dorsal clypeal margin: absent between antennal foramina. Clypeal median carina: absent. Ventral clypeal margin: with a small median point. Mandibular color: dark basally and at teeth, becoming lighter reddish brown between these areas. Mandibular teeth: three of roughly equal size. Smooth area obliquely posterior to lateral ocellus: present. Genal sculpture: deeply reticulate-rugose with some septa much stronger than others, forming distinct rows differing in height.

Dorsal pronotal area: not set off by carina ventrally. Anterolateral corner of dorsal pronotal area: weakly rounded anteriorly. Sculpture of posteromedian area of mesoscutum: densely foveolate. Lateral margin of dorsal axillar area: with a semicircular expansion, broadest near midlength. Mesoscutellar sculpture: densely foveolate. Metascutellum in dorsal view: trapezoidal with broad apex. Dorsal surface of metascutellum: flat, or convex. Femoral depression: centrally smooth, peripherally foveolate. Leg color: coxae and femora dark, but tibiae, tarsi, and sometimes 2nd trochanters yellowish brown. Anterior corner of lateral propodeal carina: flat, without tooth. Posteromedial corner of lateral propodeal area: protruding posteriorly.

Metasoma color: black to dark reddish brown. Median lobe of T1: with a set of rugae that merge with one another. Metasoma at middle of T4: with metasomal bend



Figures 43–46. *Bracalba pinnula* sp. n., paratype female (OSUC 231794). **43** Dorsal habitus **45** Head, ventral view, holotype female (OSUC 231797) **44** Lateral habitus **46** Metasomal sterna 5–6, ventral view. Morphbank²⁵

and abrupt transition in sculpture. Posterolateral margins of metasomal terga: without protrusions. T5 median carina: absent. Longitudinal sculptural septa on T5: weak, blunt and hardly raised. Transverse sculptural septa on T5: about as strong as the longitudinal septa. T5 setae: directed posteriorly, arising from anterior edge of sculptural mesh. T6: as broad or broader than long. T6 laterotergite: overlapping S6. S4 median carina: present. Transverse sculptural septa on S5: about as strong as the longitudinal septa. S5 setae: directed posteriorly, arising from anterior edge of sculptural mesh. Lateral carinae of S6: absent. Apex of S6: without notch.

Male. Body length 2.75–3.13 mm (n=4). Flagellomere length: A3 over 1.5× as long as broad, most others as long or longer than broad. T7: flat and posteriorly truncate.

Diagnosis. *Female.* A6 with 1 or 2 large sensilla; 3 mandibular teeth of approximately equal length; metascutellum trapezoidal with a broad and truncate or slightly concave apex; metasomal bend present but very weak; sculpture posterior to metasomal bend with longitudinal and transverse septa of about equal height and hardly differing from those anterior to the bend; T4–T6 and S4–S5 without median carina; S4 sometimes with a slight median carina; T6 longer than broad; S5 with longitudinal septa not stronger than transverse septa; S6 without apical notch. *Bracalba pinnula* has only a very slight metasomal bend, and therefore it can easily be confused

with those species that lack the bend. It differs from most of these species in that the fore wings reach to the middle of T₄, and from *B. intermedia* in having a long and trapezoidal metascutellum.

Etymology. Latin noun, meaning “a small fin.” This is considered to be a noun in apposition to the generic name.

Link to distribution map. <http://hol.osu.edu/map-full.html?id=302158>

Material examined. Holotype, female: AUSTRALIA: WA, WA122401, 7km N Stirling Range National Park, 34°19'S, 118°11'E, 900ft, 24.XII.1994, malaise trap, L. S. Kimsey & R. B. Kimsey, OSUC 231797 (deposited in WAMP). Paratypes: AUSTRALIA: 12 females, 11 males, OSUC 149756 (AEIC); OSUC 230822–230824, 367508, 367512–367519 (ANIC); OSUC 148700, 231780–231781 (CNCI); OSUC 238452 (QDPC); OSUC 179086, 179088–179089, 231794–231796 (UCDC).

***Bracalba propodealis* Burks, sp. n.**

urn:lsid:zoobank.org:act:F867EC04-0D6C-43C9-8BF8-E134A5C8D488

urn:lsid:biosci.ohio-state.edu:osuc_concepts:302160

http://species-id.net/wiki/Bracalba_propodealis

Figures 47–50; Morphbank²⁶

Description. *Female.* Body length 3.62–3.88 mm (n=3). Color of antenna beyond radicle: entirely dark. Radicle color: base lighter than scape. Number of claval segments with ventral gustatory sensilla: 7. Number of ventral gustatory sensilla on A6: 1.

Ocular setae: short and dense. Frontal depression: present as a vague smooth triangle. Smooth depression extending dorsolaterally from antennal foramen: present. Dorsal clypeal margin: absent between antennal foramina. Clypeal median carina: absent. Ventral clypeal margin: with a small median point. Mandibular color: dark basally and at teeth, becoming lighter reddish brown between these areas. Mandibular teeth: three, but middle tooth tiny. Smooth area obliquely posterior to lateral ocellus: present. Genal sculpture: deeply reticulate-rugose with some septa much stronger than others, forming distinct rows differing in height.

Dorsal pronotal area: not set off by carina ventrally. Anterolateral corner of dorsal pronotal area: weakly rounded anteriorly. Sculpture of posteromedian area of mesoscutum: foveolate with broad longitudinal septa. Lateral margin of dorsal axillar area: with a semicircular expansion, broadest near midlength. Mesoscutellar sculpture: sparsely foveolate, with large smooth interspaces. Metascutellum in dorsal view: very short, subrectangular. Dorsal surface of metascutellum: convex. Femoral depression: irregularly foveolate but not crossed by carinae. Leg color: coxae, femora (aside from their apices), and at least the last two tarsomeres dark, otherwise yellowish brown. Anterior corner of lateral propodeal carina: with longitudinal tooth-like crest extending dorsally above level of metascutellum. Posteromedial corner of lateral propodeal area: protruding posteriorly.

Metasoma color: black to dark reddish brown. Median lobe of T₁: with 7 or more longitudinal carinae. Metasoma at middle of T₄: with metasomal bend and abrupt



Figures 47–50. *Bracalba propodealis* sp. n., holotype female (OSUC 227590). **47** Dorsal habitus, paratype female (OSUC 227591) **48** Lateral habitus, paratype female (OSUC 238190) **49** Head, ventral view **50** Metasomal sterna 5–6, ventral view. Morphbank²⁶

transition in sculpture. Posterolateral margins of metasomal terga: without protrusions. T5 median carina: absent. Longitudinal sculptural septa on T5: weak, blunt and hardly raised. Transverse sculptural septa on T5: about as strong as the longitudinal septa. T5 setae: directed posteriorly, arising from anterior edge of sculptural mesh. T6: longer than broad. T6 laterotergite: overlapping S6. S4 median carina: present. Transverse sculptural septa on S5: about as strong as the longitudinal septa. S5 setae: not directed posteriorly, arising from center of sculptural mesh. Lateral carinae of S6: absent. Apex of S6: without notch.

Male. unknown.

Diagnosis. *Female.* A6 with 1 large ventral sensillum; 3 mandibular teeth with the middle tooth much smaller than the others; metascutellum subrectangular, very short and broad; anteromedial margins of lateral propodeal area extending dorsally above the metascutellar surface as distinct protrusions; metasomal bend extremely weak; sculpture posterior to metasomal bend with longitudinal septa stronger than those anterior to the bend, transverse septa as strong or slightly weaker than longitudinal septa; T4–T6 and S6 without median carina, S4–S5 with median carina; S6 without apical notch.

Etymology. An adjective, combining the anatomical term propodeum with the Latin adjectival suffix *-alis*, referring to its unique propodeum.

Link to distribution map. <http://hol.osu.edu/map-full.html?id=302160>

Material examined. Holotype, female: AUSTRALIA: WA, Mount Cooke, 28.I-17.II.1991, malaise trap, M. S. Harvey & J. M. Waldock, OSUC 227590 (deposited in WAMP). Paratypes: AUSTRALIA: 2 females, OSUC 238190 (CNCI); OSUC 227591 (WINC).

Comments. The propodeum in this species is unique among the specimens studied. It is very difficult to determine in all examined specimens if the metasomal bend is present or not.

***Bracalba sculptifrons* Burks, sp. n.**

urn:lsid:zoobank.org:act:F99233BD-F270-46FF-8E14-9507C07E6D9C

urn:lsid:biosci.ohio-state.edu:osuc_concepts:302161


http://species-id.net/wiki/Bracalba_sculptifrons

Figures 51–54; Morphbank²⁷

Description. *Female.* Body length 5.75–6.13 mm (n=2). Color of antenna beyond radicle: entirely dark. Radicle color: same as scape. Number of claval segments with ventral gustatory sensilla: 7. Number of ventral gustatory sensilla on A6: 2.

Ocular setae: short and dense. Frontal depression: with irregular rugae indicating large foveae. Smooth depression extending dorsolaterally from antennal foramen: present. Dorsal clypeal margin: angular, emarginate medially. Clypeal median carina: present. Ventral clypeal margin: with a small median point. Mandibular color: dark basally and at teeth, becoming lighter reddish brown between these areas. Mandibular teeth: three, but middle tooth tiny. Smooth area obliquely posterior to lateral ocellus: present. Genal sculpture: reticulate-rugose with strong dorsoventral carinae.

Dorsal pronotal area: not set off by carina ventrally. Anterolateral corner of dorsal pronotal area: truncate anteriorly. Sculpture of posteromedian area of mesoscutum: densely foveolate. Lateral margin of dorsal axillar area: with a semicircular expansion, broadest near midlength. Mesoscutellar sculpture: densely foveolate. Metascutellum in dorsal view: trapezoidal with broad apex. Dorsal surface of metascutellum: apex protruding dorsally. Femoral depression: centrally smooth, peripherally foveolate. Leg color: coxae and at least the last three tarsomeres dark, otherwise reddish. Anterior



Figures 51–54. *Bracalba sculptifrons* sp. n., paratype female (OSUC 148704). **51** Dorsal habitus **53** Head, ventral view **54** Metasomal sterna 5–6, ventral view, holotype female (OSUC 231799) **52** Lateral habitus. Morphbank²⁷

corner of lateral propodeal carina: flat, without tooth. Posteromedial corner of lateral propodeal area: protruding posteriorly.

Metasoma color: T2 to flat part of T4, troughs of S1 to variable part of S5 reddish, otherwise dark. Median lobe of T1: with 7 or more longitudinal carinae. Metasoma at middle of T4: with metasomal bend and abrupt transition in sculpture. Posterolateral margins of metasomal terga: without protrusions. T5 median carina: absent. Longitudinal sculptural septa on T5: strong, sharply raised. Transverse sculptural septa on T5: weak or absent, much weaker than the longitudinal septa. T5 setae: not directed

posteriorly, arising from center of sculptural mesh. T6: longer than broad. T6 laterotergite: overlapping S6. S4 median carina: present. Transverse sculptural septa on S5: weak or absent, much weaker than the longitudinal septa. S5 setae: not directed posteriorly, arising from center of sculptural mesh. Lateral carinae of S6: not meeting apically. Apex of S6: without notch.

Male. unknown.

Diagnosis. *Female.* A6 with 2 large ventral sensilla; 3 mandibular teeth, with middle tooth much smaller than the others; metascutellum broadly trapezoidal with a broad truncate apex; metasomal bend present; sculpture posterior to metasomal bend strong longitudinal septa and weaker transverse septa; T4–T6 and S4–S6 with or without median carina; T6 longer than broad; S6 without apical notch. This species is very similar to *B. magnirubra* in coloration and in having a relatively large body, but it differs in many characters, especially in lacking the apical S6 notch in females.

Etymology. Used as a noun in apposition to the generic name, derived from a combination of the Latin participle sculptus and noun frons, referring to its strong network of frontal carinae.

Link to distribution map. <http://hol.osu.edu/map-full.html?id=302161>

Material examined. Holotype, female: AUSTRALIA: WA, ~25km E Perth, John Forrest National Park, 23.XII-27.XII.1986, J. S. Noyes, OSUC 231799 (deposited in WAMP). Paratype: AUSTRALIA: 1 female, OSUC 148704 (CNCI).

***Bracalba sparsa* Burks, sp. n.**

url:lsid:zoobank.org:act:C947CBC7-D26E-4001-AE11-3A8EDD4873A3

url:lsid:biosci.ohio-state.edu:osuc_concepts:302162


http://species-id.net/wiki/Bracalba_sparsa

Figures 55–59; Morphbank²⁸

Description. *Female.* Body length 3.37–4.00 mm (n=6). Color of antenna beyond radicle: reddish-brown, darker at scape apex, pedicel, and A3. Radicle color: lighter than scape. Number of claval segments with ventral gustatory sensilla: 7. Number of ventral gustatory sensilla on A6: 1.

Ocular setae: long and sparse. Frontal depression: with irregular rugae indicating large foveae. Smooth depression extending dorsolaterally from antennal foramen: present. Dorsal clypeal margin: forming a complete connection between antennal foramina medially. Clypeal median carina: absent. Ventral clypeal margin: with a small median point. Mandibular color: dark basally and at teeth, becoming lighter reddish brown between these areas. Mandibular teeth: two, separated by narrow incision. Smooth area obliquely posterior to lateral ocellus: present. Genal sculpture: reticulate-rugose without any strong carinae.

Dorsal pronotal area: not set off by carina ventrally. Anterolateral corner of dorsal pronotal area: truncate anteriorly. Sculpture of posteromedian area of mesoscutum: densely foveolate. Lateral margin of dorsal axillar area: with a semicircular expansion,



Figures 55–59. *Bracalba sparsa* sp. n., holotype female (OSUC 231798). **55** Dorsal habitus **56** Lateral habitus **58** Head, ventral view, paratype female (OSUC 238205) **57** Head, anterior view **59** Metasomal sterna 5–6, ventral view. Morphbank²⁸

broadest near midlength. Mesoscutellar sculpture: densely foveolate. Metascutellum in dorsal view: trapezoidal but tapering to narrow apex. Dorsal surface of metascutellum: apex protruding dorsally. Femoral depression: crossed by rounded carinae. Leg color: entirely reddish. Anterior corner of lateral propodeal carina: flat, without tooth. Posteromedial corner of lateral propodeal area: protruding posteriorly.

Metasoma color: mostly reddish, variably dark at T4 bend, T5-T6, with variable smaller dark areas elsewhere. Median lobe of T1: with 7 or more longitudinal carinae. Metasoma at middle of T4: with metasomal bend and abrupt transition in sculpture.

Posterolateral margins of metasomal terga: with tooth-like protrusions. T5 median carina: absent. Longitudinal sculptural septa on T5: strong, sharply raised. Transverse sculptural septa on T5: weak or absent, much weaker than the longitudinal septa. T5 setae: not directed posteriorly, arising from center of sculptural mesh. T6: as broad or broader than long. T6 laterotergite: overlapping S6. S4 median carina: present. Transverse sculptural septa on S5: weak or absent, much weaker than the longitudinal septa. S5 setae: not directed posteriorly, arising from center of sculptural mesh. Lateral carinae of S6: not meeting apically. Apex of S6: with notch.

Male. Body length 3.5 mm (n=1). Flagellomere length: A3 over 1.5× as long as broad, most others as long or longer than broad. T7: arched and posteriorly concave.

Diagnosis. *Female.* A6 with 1 large ventral sensillum; 2 mandibular teeth; metascutellum subtrapezoidal but strongly tapering to a narrow truncate apex; metasomal bend expressed as a raised hump with posterior metasomal segments not tilted downwards; sculpture posterior to metasomal hump with strong longitudinal and transverse septa; T4–T6 without longitudinal carina; T6 broader than long; S4–S6 with longitudinal carina; S6 with apical notch.

Etymology. Latin participle meaning “scattered,” recalling the sparsely distributed setae of this species.

Link to distribution map. <http://hol.osu.edu/map-full.html?id=302162>

Material examined. Holotype, female: AUSTRALIA: WA, Toodyay, 31.X.1979, R. M. Bohart, OSUC 231798 (deposited in WAMP). Paratypes: AUSTRALIA: 4 females, 3 males, OSUC 230821 (ANIC); OSUC 148703, 238204–238206, 238208 (CNCI); OSUC 55870 (OSUC).

Comments. *Bracalba sparsa* represents a distinctive element within the genus, with its unique metasoma and unusual metascutellum.

***Bracalba tricolorata* Burks, sp. n.**

urn:lsid:zoobank.org:act:D54F676D-1EEF-4B40-88DA-CFB9583628C5

urn:lsid:biosci.ohio-state.edu:osuc_concepts:302163

http://species-id.net/wiki/Bracalba_tricolorata

Figures 60–61; Morphbank²⁹

Description. *Female.* Body length 3.88–4.25 mm (n=4). Color of antenna beyond radicle: entirely dark. Radicle color: same as scape. Number of claval segments with ventral gustatory sensilla: 7. Number of ventral gustatory sensilla on A6: 1.

Ocular setae: short and sparse. Frontal depression: with many strong transverse carinae, sparsely foveolate at torular triangle. Smooth depression extending dorsolaterally from antennal foramen: present. Dorsal clypeal margin: bordering antennal foramina, absent between them. Clypeal median carina: absent. Ventral clypeal margin: with a small median point. Mandibular color: mostly yellowish brown, dark at teeth. Mandibular teeth: three, but middle tooth tiny. Smooth area obliquely posterior to lat-



Figures 60–61. *Bracalba tricolorata* sp. n., holotype female (OSUC 238541). **60** Dorsal habitus **61** Ventral habitus. Morphbank²⁹

eral ocellus: present. Genal sculpture: deeply reticulate-rugose with some septa much stronger than others, forming distinct rows differing in height.

Dorsal pronotal area: not set off by carina ventrally. Anterolateral corner of dorsal pronotal area: weakly rounded anteriorly. Sculpture of posteromedian area of mesoscutum: foveolate with slightly stronger longitudinal septa. Lateral margin of dorsal axillar area: with a semicircular expansion, broadest near midlength. Mesoscutellar sculpture: sparsely foveolate, with large smooth interspaces. Metascutellum in dorsal view: trapezoidal with broad apex. Dorsal surface of metascutellum: convex. Femoral depression: crossed by rounded carinae. Leg color: coxae, femora (aside from their apices), and at least the last two tarsomeres dark, otherwise yellowish brown. Anterior corner of lateral propodeal carina: flat, without tooth. Posteromedial corner of lateral propodeal area: protruding posteriorly.

Metasoma color: black to dark reddish brown. Median lobe of T1: with a set of rugae that merge with one another. Metasoma at middle of T4: with very weak bend. Posterolateral margins of metasomal terga: without protrusions. T5 median carina: absent. Longitudinal sculptural septa on T5: weak, blunt and hardly raised. Transverse sculptural septa on T5: about as strong as the longitudinal septa. T5 setae: directed posteriorly, arising from anterior edge of sculptural mesh. T6: as broad

or broader than long. T6 laterotergite: overlapped by rim from S6. S4 median carina: absent. Transverse sculptural septa on S5: weak or absent, much weaker than the longitudinal septa. S5 setae: directed posteriorly, arising from anterior edge of sculptural mesh. Lateral carinae of S6: forming complete peripheral carina. Apex of S6: without notch.

Male. Body length 3.75–4.00 mm (n=3). Flagellomere length: A3 over 1.5× as long as broad, most others as long or longer than broad. T7: arched and posteriorly concave.

Diagnosis. *Female.* A6 with 1 large ventral sensillum; 3 mandibular teeth with the middle tooth much smaller than the others; metascutellum broadly trapezoidal with a broad apex that is truncate or sometimes slightly incised; metasomal bend present but very weak; sculpture posterior to metasomal bend with transverse septa about as strong as the longitudinal septa; T4–T6 without median carina, S4–S6 with or without median carina; T6 about as broad as long; S5 with longitudinal septa much stronger than transverse septa; S6 without apical notch. This species is very similar to *B. pinnula*, but has distinctly different ventral metasomal sculpture.

Etymology. Latin participle meaning “tricky,” named for its strong similarity to some other species and its very subtle metasomal bend.

Link to distribution map. <http://hol.osu.edu/map-full.html?id=302163>

Material examined. Holotype, female: AUSTRALIA: WA, 20km N Denmark, 16.I.1987, J. S. Noyes, OSUC 238541 (deposited in WAMP). Paratypes: AUSTRALIA: 3 females, 3 males, OSUC 148699, 238192, 238195, 238197, 238199-238200 (CNCI).

***Bracalba tridentata* Burks, sp. n.**

urn:lsid:zoobank.org:act:193C973C-CFBA-48E9-8F82-34589E44F32E

urn:lsid:biosci.ohio-state.edu:osuc_concepts:302164

http://species-id.net/wiki/Bracalba_tridentata

Figures 62–65; Morphbank³⁰

Description. *Female.* Body length 5.12–5.38 mm (n=3). Color of antenna beyond radicle: entirely dark. Radicle color: same as scape. Number of claval segments with ventral gustatory sensilla: 7. Number of ventral gustatory sensilla on A6: 1.

Ocular setae: long and sparse. Frontal depression: with many irregularly transverse rugae. Smooth depression extending dorsolaterally from antennal foramen: present. Dorsal clypeal margin: arched, interrupted by broad median carina. Clypeal median carina: present. Ventral clypeal margin: with a small median point. Mandibular color: dark basally and at teeth, becoming lighter reddish brown between these areas. Mandibular teeth: three, but middle tooth tiny. Smooth area obliquely posterior to lateral ocellus: present. Genal sculpture: deeply reticulate-rugose with some septa much stronger than others, forming distinct rows differing in height.

Dorsal pronotal area: not set off by carina ventrally. Anterolateral corner of dorsal pronotal area: truncate anteriorly. Sculpture of posteromedian area of mesoscutum:



Figures 62–65. *Bracalba tridentata* sp. n., paratype female (OSUC 238116). **62** Dorsal habitus **64** Head, ventral view, metasomal sterna 5–6, ventral view **65** Metasomal sterna 5–6, ventral view, holotype female (OSUC 238113) **63** Lateral habitus. Morphbank³⁰

densely foveolate. Lateral margin of dorsal axillar area: with a semicircular expansion, broadest near midlength. Mesoscutellar sculpture: densely foveolate. Metascutellum in dorsal view: trapezoidal with broad apex, or elongate-trapezoidal but with incised apex. Dorsal surface of metascutellum: apex protruding dorsally. Femoral depression: crossed by 4–6 dorsal carinae, 7–8 ventral carinae interrupted by central smooth area. Leg color: yellowish-brown except for dorsal external part of coxae and sometimes tarsomeres 2–5. Anterior corner of lateral propodeal carina: flat, without tooth. Posteromedial corner of lateral propodeal area: protruding posteriorly.

Metasoma color: black to dark reddish brown. Median lobe of T1: with 7 or more longitudinal carinae. Metasoma at middle of T4: with metasomal bend and abrupt transition in sculpture. Posterolateral margins of metasomal terga: without protrusions. T5 median carina: absent. Longitudinal sculptural septa on T5: strong, sharply raised. Transverse sculptural septa on T5: about as strong as the longitudinal septa. T5 setae: not directed posteriorly, arising from center of sculptural mesh. T6: as broad or broader than long. T6 laterotergite: overlapping S6. S4 median carina: absent. Transverse sculptural septa on S5: weak or absent, much weaker than the longitudinal septa. S5 setae: not directed posteriorly, arising from center of sculptural mesh. Lateral carinae of S6: not meeting apically. Apex of S6: with notch.

Male. unknown.

Diagnosis. *Female.* A6 with 1 large ventral sensillum; middle mandibular tooth present but much smaller than the others; metascutellum broadly trapezoidal and with a broadly truncate apex that may have a conspicuous median notch; metasomal bend strong; sculpture posterior to metasomal bend distinctly different from that anterior to it, with very weak transverse septa and strong longitudinal septa; T6 about as long as broad; S6 with apical notch. This species is very similar to *B. hesperia*, but S6 in that species does not have an apical notch. It is also similar to *B. sparsa*, but that species has a unique metasomal bend and metascutellar shape.

Etymology. Latin adjective, named for the usually 3-pronged pattern formed apically by dark cuticle on T6.

Link to distribution map. <http://hol.osu.edu/map-full.html?id=302164>

Material examined. Holotype, female: AUSTRALIA: SA, 32km N Renmark, Amalia Dam, xeric mallee scrub, MT 4, ROM 2000041, Bookmark Biosphere Reserve, 33°53'S, 140°43'E, 263m, 15.II-15.IV.2000, malaise trap, D. C. Darling, OSUC 238113 (deposited in SAMA). Paratypes: AUSTRALIA: 2 females, OSUC 238114, 238116 (CNCI).

Acknowledgments

We thank L. Musetti, J. Jennings, J. Cora, A. Valerio, S. Mantel, D. Stringer, and other members of the PBI team, for assistance with specimens, photographs, and databasing. This work was funded by the National Science Foundation under grant No. DEB-0614764 to N.F. Johnson and A.D. Austin, and by the Australian Biological Resources Study grant 208-058 to A.D. Austin.

References

ABRS (2012) Australian Faunal Directory. Platygastridae. Australian Biological Resources Study, Canberra. <http://www.environment.gov.au/biodiversity/abrs/online-resources/fauna/afd/taxa/PLATYGASTROIDEA> [accessed on 21 May 2012]

- Austin AD (1986) A taxonomic revision of *Mirobaeoides* Dodd (Hymenoptera Scelionidae). *Australian Journal of Zoology* 34: 315–337.³¹ doi: 10.1071/ZO9860315
- Austin AD, Field SA (1997) The ovipositor system of scelionid and platygastrid wasps (Hymenoptera: Platygastroidea): comparative morphology and phylogenetic implications. *Invertebrate Taxonomy* 11: 1–87.³² doi: 10.1071/IT95048
- Austin AD, Johnson NF, Downton M (2005) Systematics, evolution and biology of scelionid and platygastrid wasps. *Annual Review of Entomology* 50: 553–582.³³ doi: 10.1146/annurev.ento.50.071803.130500
- Bin F (1981) Definition of female antennal clava based on its plate sensilla in Hymenoptera Scelionidae Telenominae. *Redia* 64: 245–261.³⁴
- Dangerfield P, Austin A, Baker G (2001) *Biology, Ecology and Systematics of Australian Scelio*, wasp parasitoids of locust and grasshopper eggs. CSIRO Publications: Collingwood, Victoria, 254 pp.³⁵
- Dodd AP (1920) Notes on the exotic Proctotrupeoidea in the British and Oxford University Museums, with descriptions of new genera and species. *Transactions of the Entomological Society of London* 1919: 321–382.³⁶ doi: 10.1111/j.1365-2311.1920.tb00008.x
- Dodd AP (1931) The genus *Oxyscelio* Kieffer, its synonymy and species, with a description of one new genus (Hymenoptera: Proctotrypoidea). *Proceedings of the Royal Society of Queensland* 42: 71–81.³⁷
- Eady RD (1968) Some illustrations of microsculpture in the Hymenoptera. *Proceedings of the Royal Entomological Society of London, (A)* 43: 66–72.³⁸ doi: 10.1111/j.1365-3032.1968.tb01029.x
- Galloway ID (1976) The types of Australian species of the subfamily Scelioninae (Hymenoptera: Scelionidae). *Queensland Journal of Agricultural and Animal Sciences* 33: 83–114.³⁹
- Galloway ID, Austin AD (1984) Revision of the Scelioninae (Hymenoptera: Scelionidae) in Australia. *Australian Journal of Zoology Supplementary Series* 99: 1–138.⁴⁰ doi: 10.1071/AJZS099
- Goloboff PA, Farris JS, Nixon KC (2003) T.N.T. – Tree Analysis Using New Technology, version 1.1. Computer software and manual, available at: (<http://www.zmuc.dk/public/phylogeny>).
- Goloboff PA, Farris JS, Nixon KC (2008) TNT: a free program for phylogenetic analysis. *Cladistics* 24: 774–786.⁴¹ doi: 10.1111/j.1096-0031.2008.00217.x
- Iqbal M, Austin AD (2000) Systematics of the wasp genus *Ceratobaenus* Ashmead (Hymenoptera: Scelionidae) from Australasia: parasitoids of spider eggs. *Records of the South Australian Museum Monograph Series* 6: 1–164.⁴²
- Johnson NF (1991) Revision of Australasian *Trissolcus* species (Hymenoptera: Scelionidae). *Invertebrate Taxonomy* 5: 211–39.⁴³ doi: 10.1071/IT9910211
- Johnson NF (1992) Catalog of world Proctotrupeoidea excluding Platygastridae. *Memoirs of the American Entomological Institute* 51: 1–825.⁴⁴
- Johnson NF, Masner L (1985) Revision of the genus *Psix* Kozlov & Lê (Hymenoptera: Scelionidae). *Systematic Entomology* 10: 33–58.⁴⁵ doi: 10.1111/j.1365-3113.1985.tb00562.x
- Masner L (1976) Revisionary notes and keys to world genera of Scelionidae (Hymenoptera: Proctotrupeoidea). *Memoirs of the Entomological Society of Canada* 97: 1–87.⁴⁶ doi: 10.4039/entm10897fv

- Masner L, Johnson NF (2007) *Xentor*, a new endemic genus from Fiji (Hymenoptera: Platygastroidea: Scelionidae) and description of three new species. *Fiji Arthropods*, 9: 11–20.⁴⁷ doi: 10.1206/0003-0082(2007)3551[1:TANGON]2.0.CO;2
- Mikó I, Vilhelmsen L, Johnson NF, Masner L, Péntzes Z (2007) Skeletomusculature of Scelionidae (Hymenoptera: Platygastroidea): head and mesosoma. *Zootaxa* 1571: 1–78.⁴⁸ doi: 10.1206/0003-0082(2007)3551[1:TANGON]2.0.CO;2
- Muesebeck CFW, Walkley LM (1956) Type species of the genera and subgenera of parasitic wasps comprising the superfamily Proctotrupoidea (order Hymenoptera). *Proceedings of the United States National Museum* 105: 319–419.⁴⁹ doi: 10.5479/si.00963801.3359.319
- Stevens NB, Austin AD (2007) Systematics, distribution and biology of the Australian ‘micro-flea’ wasps, *Baeus* spp. (Hymenoptera: Scelionidae): parasitoids of spider eggs. *Zootaxa* 1499: 1–45.⁵⁰
- Talamas EJ, Masner L, Johnson NF (2011) Revision of the *Paridris nephta* species group (Hymenoptera, Platygastroidea, Platygastriidae). *ZooKeys* 133: 49–94.⁵¹ doi: 10.3897/zookeys.133.1613
- Vilhelmsen L, Mikó I, Krogmann L (2010) Beyond the wasp-waist: structural diversity and phylogenetic significance of the mesosoma in apocritan wasps (Insecta: Hymenoptera). *Zoological Journal of the Linnean Society* 159: 22–194.⁵² doi: 10.1111/j.1096-3642.2009.00576.x
- Walker F (1839) *Monographia Chalciditum*. Vol. II. Hyppolite Baillière, London, 100 pp.⁵³

Endnotes

- 1 urn:lsid:biocol.org:col:1008
- 2 urn:lsid:biocol.org:col:32975
- 3 urn:lsid:biocol.org:col:32981
- 4 urn:lsid:biocol.org:col:1009
- 5 urn:lsid:biocol.org:col:1012
- 6 urn:lsid:biocol.org:col:33928
- 7 urn:lsid:biocol.org:col:33791
- 8 urn:lsid:biocol.org:col:1014
- 9 urn:lsid:biocol.org:col:34157
- 10 urn:lsid:biocol.org:col:34161
- 11 urn:lsid:biocol.org:col:34982
- 12 urn:lsid:biocol.org:col:34411
- 13 urn:lsid:biocol.org:col:34578
- 14 urn:lsid:biocol.org:col:34593
- 15 <http://www.morphbank.net/?id=799328>
- 16 <http://www.morphbank.net/?id=799329>
- 17 <http://www.morphbank.net/?id=799330>
- 18 <http://www.morphbank.net/?id=799331>
- 19 <http://www.morphbank.net/?id=799332>

- 20 <http://www.morphbank.net/?id=799333>
- 21 <http://www.morphbank.net/?id=799334>
- 22 <http://www.morphbank.net/?id=799335>
- 23 <http://www.morphbank.net/?id=799336>
- 24 <http://www.morphbank.net/?id=799337>
- 25 <http://www.morphbank.net/?id=799338>
- 26 <http://www.morphbank.net/?id=799339>
- 27 <http://www.morphbank.net/?id=799340>
- 28 <http://www.morphbank.net/?id=799341>
- 29 <http://www.morphbank.net/?id=799342>
- 30 <http://www.morphbank.net/?id=799343>
- 31 urn:lsid:biosci.ohio-state.edu:osuc_pubs:111
- 32 urn:lsid:biosci.ohio-state.edu:osuc_pubs:20940
- 33 urn:lsid:biosci.ohio-state.edu:osuc_pubs:21201
- 34 urn:lsid:biosci.ohio-state.edu:osuc_pubs:131
- 35 urn:lsid:biosci.ohio-state.edu:osuc_pubs:9560
- 36 urn:lsid:biosci.ohio-state.edu:osuc_pubs:409
- 37 urn:lsid:biosci.ohio-state.edu:osuc_pubs:404
- 38 urn:lsid:biosci.ohio-state.edu:osuc_pubs:21506
- 39 urn:lsid:biosci.ohio-state.edu:osuc_pubs:353
- 40 urn:lsid:biosci.ohio-state.edu:osuc_pubs:463
- 41 <http://dx.doi.org/10.1111/j.1096-0031.2008.00217.x>
- 42 urn:lsid:biosci.ohio-state.edu:osuc_pubs:8079
- 43 urn:lsid:biosci.ohio-state.edu:osuc_pubs:633
- 44 urn:lsid:biosci.ohio-state.edu:osuc_pubs:229
- 45 urn:lsid:biosci.ohio-state.edu:osuc_pubs:657
- 46 urn:lsid:biosci.ohio-state.edu:osuc_pubs:311
- 47 urn:lsid:biosci.ohio-state.edu:osuc_pubs:21304
- 48 urn:lsid:biosci.ohio-state.edu:osuc_pubs:21300
- 49 urn:lsid:biosci.ohio-state.edu:osuc_pubs:240
- 50 urn:lsid:biosci.ohio-state.edu:osuc_pubs:21213
- 51 urn:lsid:biosci.ohio-state.edu:osuc_pubs:23583
- 52 urn:lsid:biosci.ohio-state.edu:osuc_pubs:23221
- 53 urn:lsid:biosci.ohio-state.edu:osuc_pubs:528

Appendix I

Characters

Color of antenna beyond radicle

1. entirely dark
2. entirely reddish
3. mostly dark, extreme base of scape becoming reddish
4. reddish-brown, darker at scape apex, pedicel, and A3
5. mostly dark, ventral parts of pedicel apex, A4-A12 variably lighter
6. yellowish-brown
7. dark except for pedicel apex
8. claval segments brown, basal segments yellowish

Radicle color

1. same as scape
2. base lighter than scape
3. lighter than scape

Number of claval segments with ventral gustatory sensilla

1. 6
2. 7

Number of ventral gustatory sensilla on A6

1. 1
2. 2
3. 0

Ocular setae

1. short and dense
2. long and sparse
3. short and sparse
4. long and dense
5. absent

Frontal depression ventral sculpture

1. with transverse rugae
2. with large areoles
3. smooth or foveolate
4. with median longitudinal carina
5. inapplicable (frontal depression absent)

Frontal depression dorsal sculpture

1. with transverse rugae
2. with large areoles
3. smooth or foveolate
4. with median longitudinal carina
5. inapplicable (frontal depression absent)

Smooth depression extending dorsolaterally from antennal foramen

1. absent
2. present

Dorsal clypeal margin

1. absent between antennal foramina
2. interrupted by a median areole below interantennal process
3. interrupted by median carina
4. complete only between antennal foramina medially
5. wrinkle-like with median peak
6. angular, emarginate medially
7. uniformly arched without interruption

Clypeal median carina

1. present
2. absent

Ventral clypeal margin

1. straight
2. with a small median point
3. convex
4. concave

Mandibular color

1. dark basally and at teeth, becoming lighter reddish brown between these areas
2. entirely dark
3. mostly yellowish brown, dark at teeth
4. mostly reddish brown, dark at teeth

Mandibular teeth

1. three of roughly equal size
2. two, separated by narrow incision
3. three, but middle tooth tiny
4. two, broadly separated by an angled flat area

Smooth area obliquely posterior to lateral ocellus

1. present
2. absent

Genal sculpture

1. reticulate-rugose with strong dorsoventral carinae
2. reticulate-rugose without any strong carinae
3. deeply reticulate-rugose with some septa much stronger than others, forming distinct rows differing in height

Dorsal pronotal area

1. not set off by carina
2. set off by carina

Anterolateral corner of dorsal pronotal area

1. protruding anteriorly
2. truncate anteriorly
3. weakly rounded anteriorly

Sculpture of posteromedian area of mesoscutum

1. sparsely foveolate, with large smooth interspaces
2. with many longitudinal carinae
3. densely foveolate
4. foveolate with broad longitudinal septa
5. foveolate with slightly stronger longitudinal septa
6. foveolate, with some interspersed additional tiny foveae

Lateral margin of dorsal axillar area

1. not expanded
2. broadly expanded
3. triangularly expanded or with posterior tooth, broadest posteriorly
4. with a semicircular expansion, broadest near midlength

Mesoscutellar sculpture

1. densely foveolate
2. sparsely foveolate, with large smooth interspaces
3. densely foveolate with one or two median longitudinal channels
4. densely foveolate but with smooth central area
5. densely foveolate with slightly stronger longitudinal septa
6. foveolate, with some interspersed additional tiny foveae
7. densely foveolate with a longitudinal carina
8. medially smooth except for a few irregular carinae

Metascutellum in dorsal view

1. strongly transverse, subrectangular
2. trapezoidal with broad apex
3. trapezoidal but tapering to narrow apex
4. elongate-trapezoidal but with incised apex
5. very short, subrectangular
6. semicircular
7. elongate-rectangular

Dorsal surface of metascutellum

1. flat
2. apex protruding dorsally
3. convex

Femoral depression

1. crossed by rounded carinae
2. irregularly foveolate but not crossed by carinae
3. smooth, not crossed by carinae
4. crossed by 4-6 dorsal carinae, 7-8 ventral carinae interrupted by central smooth area
5. centrally smooth, peripherally foveolate
6. shallowly sculptured but not crossed by carinae
7. crossed by many sharply defined carinae

Leg color

1. entirely dark
2. entirely reddish
3. coxae dark, all else reddish
4. coxae, femora (aside from their apices), and at least the last two tarsomeres dark, otherwise yellowish brown
5. coxae and at least the last three tarsomeres dark, otherwise reddish
6. reddish but with at least last two tarsomeres dark
7. dark except for 2nd trochanter of fore leg and mid leg, tip of mid and hind metatibia
8. reddish except for variable antero-dorsal portion of coxae
9. dark except for trochanters, tips of femora and tibiae, and tarsomeres 1-4
10. coxae and femora dark, but tibiae, tarsi, and sometimes 2nd trochanters yellowish brown
11. dark except for tips of tibiae and at least 1st tarsomere
12. yellowish-brown except for dorsal part of coxae and sometimes tarsomeres 2-5
13. dark except for tibiae, 2nd trochanter, and tarsomeres 2-5
14. dark, except trochanters, femoral apices, variable parts of tibiae, and tarsi pale
15. coxae, femora, and apical two tarsomeres dark, trochanters, tibiae, and other tarsomeres yellowish brown
16. coxae dark, leg becoming gradually lighter apically

17. coxae and at least the last three tarsomeres dark, otherwise yellowish
18. entirely yellowish

Anterior corner of lateral propodeal carina

1. flat, without tooth
2. with longitudinal tooth-like crest extending dorsally above level of metascutellum

Posteromedial corner of lateral propodeal area

1. not protruding posteriorly
2. protruding posteriorly

Metasoma color

1. black to dark reddish brown
2. entirely reddish
3. reddish, with last segment variably dark
4. reddish, with last two segments dark
5. reddish, with last 2.5 segments dark
6. T2 to flat part of T4, troughs of S1 to variable part of S5 reddish; otherwise dark
7. mostly reddish, variably dark at T4 bend, T5-T6, with variable smaller dark areas elsewhere
8. centrally reddish with some vague dark intrusions, peripherally becoming black

Median lobe of T1

1. with 7 or more longitudinal carinae
2. with a set of rugae that merge with one another
3. with median smooth area interrupting carinae
4. with 6 carinae
5. with 5 carinae

Metasoma at middle of T4

1. with metasomal bend and abrupt transition in sculpture
2. without bend
3. with very weak bend

Posterolateral margins of metasomal terga

1. with tooth-like protrusions
2. without protrusions

T5 median carina

1. absent
2. present

Longitudinal sculptural septa on T5

1. strong, sharply raised
2. weak, blunt and hardly raised

Transverse sculptural septa on T5

1. about as strong as the longitudinal septa
2. weak or absent, much weaker than the longitudinal septa

T5 setae

1. directed posteriorly, arising from anterior edge of sculptural mesh
2. not directed posteriorly, arising from center of sculptural mesh

T6

1. longer than broad
2. as broad or broader than long

T6 laterotergite

1. overlapped by rim from S6
2. overlapping S6

S4 median carina

1. absent
2. present

Transverse sculptural septa on S5

1. weak or absent, much weaker than the longitudinal septa
2. about as strong as the longitudinal septa

S5 setae

1. directed posteriorly, arising from anterior edge of sculptural mesh
2. not directed posteriorly, arising from center of sculptural mesh

Lateral carinae of S6

1. forming complete peripheral carina
2. not meeting apically
3. absent
4. weak and ruga-like, becoming indistinct posteriorly

Apex of S6

1. without notch
2. with notch
3. roundly concave

Flagellomere length

1. A3 over 1.5× as long as broad, most others as long or longer than broad
2. A3 1–1.2× as long as broad, most others transverse

T7

1. arched and posteriorly concave
2. flat and posteriorly truncate

Matrix

Chromoteleia sp.	7002444010221101223052200H02111000100020011
Bracalba_cuneata	50110[23][23]16111000[01]132[04]11001B0[02]101001100020011
Bracalba_globosa	02110[23][23]161110100242111001C01101001100020011
Bracalba_plana	4211000101110200222150401F00101001001020111
Bracalba_clavata	41020[03][02]13112020023230[02]400300200001110020001
Bracalba_hesperia	00102021401001002232124018000[01]1000000120?1?
Bracalba_intermedia	0202022061220100242400401301101000110020011
Bracalba_laminata	?11302101120200143631001?????????????0?0
Bracalba_magnirubra	221012211012010002303[12]001520010110110111000
Bracalba_nigrescens	???300101120230203131001G?????????????0?0
Bracalba_parvirubra	201012012012020004323[02]601B70000001010000?0?
Bracalba_pinnula	021[01]00010110020022301[02]401901001001111020011
Bracalba_sculptifrons	00110[13][13]150120000123011401450000110110110?1?
Bracalba_propodealis	0110022101120200233142111300001000111120?1?
Bracalba_tricolorata	0010200101120220243112001301201001000000010
Bracalba_sparsa	3210122131110100123021001160000111110111000
Bracalba_tridentata	00101001201202001230[13]1301B00000011100111?1?

Appendix 2

Taxonomic records for all records used in the present paper. (doi: 10.3897/zookeys.236.3434.app2) File format: DarwinCore Archive.

Explanation note: Darwin Core Archive (DwC-A) is the preferred format for publishing data to the Global Biodiversity Information Facility (GBIF) network. Together with Dublin Core (on which its ideas are based), it is used by GBIF and others to encode data about organism names, taxonomies and species information.

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Citation: Burks RA, Masner L, Johnson NF, Austin AD (2012) Taxonomic revision of *Bracalba Dodd* (Hymenoptera, Platygastridae s.l.), a parasitoid wasp genus endemic to Australia. *ZooKeys* 236: 1–53. doi: 10.3897/zookeys.236.3434.app2

Appendix 3

Locality records for all records used in the present paper. (doi: 10.3897/zookeys.236.3434.app3) File format: DarwinCore Archive.

Explanation note: Darwin Core Archive (DwC-A) is the preferred format for publishing data to the Global Biodiversity Information Facility (GBIF) network. Together with Dublin Core (on which its ideas are based), it is used by GBIF and others to encode data about organism names, taxonomies and species information.

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Citation: Burks RA, Masner L, Johnson NF, Austin AD (2012) Taxonomic revision of *Bracalba Dodd* (Hymenoptera, Platygastridae s.l.), a parasitoid wasp genus endemic to Australia. *ZooKeys* 236: 1–53. doi: 10.3897/zookeys.236.3434.app3
