

PUBLISHED VERSION


M. E. Carrillo-Serrano, I. C. Cloët, K. Tsushima, A. W. Thomas, I. R. Afnan
Quark mass variation and its effect on nuclear binding
XII International Symposium on Nuclei in the Cosmos, 2012 / J. Lattanzio, A. Karakas, M. Lugaro, G. Dracoulis (eds.), article 147

© Copyright owned by the author(s) under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike Licence

PERMISSIONS

<http://pos.sissa.it/POSlicense.html>

Given the Open Access nature of this proceedings collection, PoS has chosen to apply the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License.



This is a human-readable summary of (and not a substitute for) the [license](#).

[Disclaimer](#)

You are free to:

Share — copy and redistribute the material in any medium or format

The licensor cannot revoke these freedoms as long as you follow the license terms.

Under the following terms:



Attribution — You must give **appropriate credit**, provide a link to the license, and **indicate if changes were made**. You may do so in any reasonable manner, but not in any way that suggests the licensor endorses you or your use.



NonCommercial — You may not use the material for **commercial purposes**.



NoDerivatives — If you **remix, transform, or build upon** the material, you may not distribute the modified material.

No additional restrictions — You may not apply legal terms or **technological measures** that legally restrict others from doing anything the license permits.

<http://hdl.handle.net/2440/83761>

Quark Mass Variation and its effect on nuclear binding

M. E. Carrillo-Serrano*

*CSSM and ARC Centre of Excellence for Particle Physics at the Tera-scale,
School of Chemistry and Physics, University of Adelaide, Adelaide SA 5005, Australia
E-mail: manuel.carrilloserrano@adelaide.edu.au*

I. C. Cloët

*CSSM, School of Chemistry and Physics, University of Adelaide, Adelaide SA 5005, Australia
E-mail: ian.cloet@adelaide.edu.au*

K. Tsushima

*CSSM, School of Chemistry and Physics, University of Adelaide, Adelaide SA 5005, Australia
E-mail: kazuo.tsushima@adelaide.edu.au*

A. W. Thomas

*CSSM and ARC Centre of Excellence for Particle Physics at the Tera-scale,
School of Chemistry and Physics, University of Adelaide, Adelaide SA 5005, Australia
E-mail: anthony.thomas@adelaide.edu.au*

I. R. Afnan

*School of Chemical and Physical Sciences, Flinders University, GPO Box 2100, Adelaide SA
5001 Australia
E-mail: Iraj.Afnan@Flinders.edu.au*

We consider the variations of the nucleon mass and the masses of the mesons typically used in a one-boson exchange model arising from possible changes in light quark masses. These results are used to calculate the corresponding changes of the energy of the 1S_0 anti-bound state and the binding energies of ^2H , ^3H , and other nuclei. The possible consequences for the standard scenario for big bang nucleosynthesis are discussed [9]. In particular the possible solution of the ^7Li problem through a variation of quark masses is not supported.

*XII International Symposium on Nuclei in the Cosmos,
August 5-12, 2012
Cairns, Australia*

*Speaker.

1. INTRODUCTION

Recently there has been growing evidence from quasar absorption spectra showing that the fine structure constant may have changed by an amount of around 10^{-5} in a billion years [1, 2, 3, 4]. Although this evidence is still debatable [5], that variation can be related to higher changes in other fundamental quantities such as Λ_{QCD} and quark masses [6, 7, 8]. In particular a variation of the light quark masses $\delta m_q/m_q$ would be of order 38 times that of the fine structure constant $\delta\alpha/\alpha$ [6].

Different approaches to calculate the effect of these apparent quark mass changes in the nucleon-nucleon force have been studied. These include: effective field theory [10, 11], constraints from lattice QCD [12, 13], and calculations using the Argonne potential and the Schwinger-Dyson method [14].

In this work we calculate the variations with quark mass of the 1S_0 anti-bound state, and the binding energies of ^2H , ^3H , and other selected light nuclei [9]. This is studied considering the variations of the mass of each of the mesons that are employed in a one-boson exchange (OBE) picture of the nucleon-nucleon interaction.

These results are included into the quark-meson coupling model (QMC) for the calculations of the binding energies of ^7Li , ^{12}C , and ^{16}O as well as using a typical OBE model for ^2H , the 1S_0 antibound state and ^3H [9].

Section II deals with the calculations for each of the mesons included and the nucleon. Sections III and IV give the results for the deuteron, triton and some other nuclei. Finally the possible effect to these results for big bang nucleosynthesis (BBN) is discussed [9], showing that in our model the ^7Li problem cannot be solved in this way.

2. MESON AND NUCLEON MASSES

We use three steps to calculate the changes in the meson masses with the variation of the current quark mass [9]. Firstly we used the Nambu-Jona-Laisino (NJL) model [16, 17] to calculate the relationship between the bare mass of the sigma meson ($m_\sigma^{(0)}$) with the pion mass (m_π), then we introduce the self energy contributions for the σ , ρ , and ω mesons. In the case of the σ the inclusion was subjected to the position of the pole on the second sheet of the energy plane found by Leutwyler *et. al.* [18]; for the ρ and ω we use chiral fits from lattice QCD [22]. Finally all the results are translated to relationships with the quark mass through the Gell-Mann-Oakes-Renner (GMOR) relation. In the case of the nucleon mass, we use the recent calculations of the sigma commutator $\sigma_{\pi N}$ [23].

2.1 VARIATION IN m_σ WITH m_q

With an OBE approach we parametrize the intermediate range attraction of the nucleon-nucleon interaction through the exchange of a σ meson [9]. Our results can be compared to [14] where a different approach is used. This should serve to pin down the uncertainties on this calculation. In our model the σ can be composed of a “bare” part which coupled to two pions. The mass of the bare part is computed using the NJL model. For the rest of the contribution we calculate the propagator of the dressed σ , which can be described as an infinite sum of terms of the form of (a) in Fig. 1 (replacing ρ by σ). Denoting the self-energy as $\Sigma_{\pi\pi}^\sigma$, the total propagator

Δ_σ can be resummed to: $\Delta_\sigma = \frac{i}{p^2 - (m_\sigma^{(0)})^2 + \Sigma_{\pi\pi}^\sigma}$. The pole of this propagator was calculated before by Leutwyler using the model independent method of the Roy equations [19], their result was: $p = m_\sigma - \frac{i}{2}\Gamma = 441_{-8}^{+16} - i272_{-12.5}^{+9}$ MeV, in agreement with the results given by the Particle Data Group (PDG) [20]. Our calculations should match this pole, and this allow us to fix $\Sigma_{\pi\pi}^\sigma$ in the form

$$\sqrt{(m_\sigma^{(0)})^2 - \Sigma_{\pi\pi}^\sigma(m_\sigma^2)} \simeq 441 - i272 \text{ MeV}. \quad (2.1)$$

Respecting chiral symmetry, the expression for this self-energy is found with a derivative coupling between the bare σ and the two pions

$$i\Sigma_{\pi\pi}^\sigma = \frac{3}{2}\gamma_0^2 \int \frac{d^4k}{(2\pi)^4} \frac{\left[1 - \frac{(\frac{p}{2}-k)^2}{\Lambda^2}\right]^{-4} \left[k^\mu (p-k)_\mu\right]^2}{(k^2 - m_\pi^2) \left((p-k)^2 - m_\pi^2\right)}, \quad (2.2)$$

with k the momentum of the pion loop, p the momentum of σ , γ the coupling constant, and the first term in the numerator of the integrand is a dipole regulator with mass Λ . This integration is performed doing the temporal part first using contour integration, whereas the spatial component includes a rotation in the complex plane of \vec{k} to ensure that the pole is in the second sheet. We then fit the three parameters: γ , λ , and $m_\sigma^{(0)}$ with the pole position [9].

Defining $m_\sigma^2(OBE) = (m_\sigma^{(0)})^2 - \Sigma_{\pi\pi}^\sigma(0)$ (i.e. at zero momentum transfer), any variation on $m_\sigma(OBE)$ with respect to m_π is given by changes in $m_\sigma^{(0)}$ and $\Sigma_{\pi\pi}^\sigma(0)$. In addition, using the Gell-Mann-Oakes-Renner (GMOR) relation [21], we finally can get the variation of $m_\sigma(OBE)$ with m_q by: $\frac{\delta m_{\sigma_{0(1)}}(OBE)}{m_{\sigma_{0(1)}}(OBE)} = 0.089(0.072) \frac{\delta m_q}{m_q}$, with the indices 0 and 1 respectively correspond the scalar σ_0 and the second one to the isoscalar σ_1 .

2.2 VARIATIONS IN m_ρ AND m_ω WITH RESPECT TO m_q

With the ρ meson we follow the analysis done by Armour *et. al.* [22] using partially quenched lattice QCD data, and extrapolating to the physical limit for their fit. The self-energy diagrams relevant are shown in Fig. 1, where diagram (a) will be denoted as $\Sigma_{\pi\pi}^\rho$ and (b) as $\Sigma_{\pi\omega}^\rho$.




Figure 1: Self-energy contributions for the ρ -meson mass.

The expressions for these diagrams are regularized using dipole cutoffs in a similar way that was done for the σ meson. The fit is then given by the following relation

$$m_\rho = \sqrt{(a_0 + a_2 m_\pi^2 + a_4 m_\pi^4)^2 + \Sigma_{TOT}}, \quad (2.3)$$

where $\Sigma_{TOT} = \Sigma_{\pi\pi}^{\rho} + \Sigma_{\pi\omega}^{\rho}$, and the coefficients, a_i , are: $a_0 = 832.00$ MeV, $a_2 = 4.94 \times 10^{-4}$ MeV $^{-1}$, $a_4 = -6.10 \times 10^{-11}$ MeV $^{-3}$ and $\Lambda = 655.00$ MeV (up to errors). Similarly to the σ case we consider almost zero momentum transferred in the OBE potential. We then can relate the changes in m_{ρ} with m_{π} according to the fit given in (2.3). The final variation gives: $\frac{\delta m_{\rho}}{m_{\rho}} = 0.034 \frac{\delta m_q}{m_q}$.

In the case of the ω the procedure is basically the same, but with the difference that there is not an analogue of diagram (a) in Fig. 1, because it violates G-parity. In addition, since we have three ways to combine ρ and π : $\Sigma_{\rho\pi}^{\omega} = 3 \times \Sigma_{\omega\pi}^{\rho}$. We use the same fit with the same coefficients as ρ , simply because their mass difference is around 10 MeV. For the coefficient for the variation in m_{ω} , we obtain is: $\frac{\delta m_{\omega}}{m_{\omega}} = 0.024 \frac{\delta m_q}{m_q}$ [9].

2.3 NUCLEON MASS

For the nucleon mass we invoke the πN sigma commutator

$$\sigma_{\pi N} = m_q \langle N | \bar{q}q | N \rangle = m_q \frac{\delta m_N}{\delta m_q}, \quad (2.4)$$

where $\bar{q}q \equiv \bar{u}u + \bar{d}d$. This result comes from the Feynman-Hellmann theorem. To extract this value we use the analysis done by Shanahan *et. al.* [23] of the PACS-CS data, $\sigma_{\pi N} = 45 \pm 6$ MeV. This finally gives a relation: $\frac{\delta m_N}{m_N} = 0.048 \frac{\delta m_q}{m_q}$.

3. ${}^7\text{Li}$, ${}^{12}\text{C}$ AND ${}^{16}\text{O}$ NUCLEI

To calculate the single-particle energies of ${}^7\text{Li}$, ${}^{12}\text{C}$ and ${}^{16}\text{O}$ nuclei we use the quark-meson coupling model, a nuclear model based on quark degrees of freedom [24]. We then calculate the changes in these binding energies with variations in the current quark mass m_q [9].

We took standard values for mesons and nucleon masses as well as the σ - N coupling constant and the rms charge radius of ${}^{40}\text{Ca}$ [25].

The final results for binding energies per nucleon as a function of the quark mass are:

$$\frac{\delta |E_{7\text{Li}}|/\text{nucleon}}{|E_{7\text{Li}}|/\text{nucleon}} = -2.571 \frac{\delta m_q}{m_q}, \quad (3.1)$$

$$\frac{\delta |E_{12\text{C}}|/\text{nucleon}}{|E_{12\text{C}}|/\text{nucleon}} = -1.438 \frac{\delta m_q}{m_q}, \quad (3.2)$$

$$\frac{\delta |E_{16\text{O}}|/\text{nucleon}}{|E_{16\text{O}}|/\text{nucleon}} = -1.082 \frac{\delta m_q}{m_q}. \quad (3.3)$$

4. VARIATION IN THE ENERGIES OF THE TWO- AND THREE-NUCLEON SYSTEMS WITH VARIATION IN THE MESON AND NUCLEON MASSES

For the deuteron, the antibound 1S_0 state and the triton; we consider an OBE potential. We choose the Bryan-Scott (BS) approach because it employs the mesons we have considered ($(\pi, \eta, \sigma_0, \sigma_1, \rho, \omega)$) [26]. For the anti-bound state 1S_0 , the location of the pole is in the second energy sheet, which makes the evaluation more complicated in order to apply the method of moments. This problem is overcome by approximating to the zero energy point, because its energy is

close to zero ($E_p = -0.066$ MeV). The next step is to include a separable potential that allows us to write the Fadeev equations as a set of coupled one dimensional integral equations.

Varying the values of the masses of the mesons included permits us to find the changes in the energies of the deuteron, the 1S_0 state and the triton. Finally, using the GMOR relation, and the coefficients relating the meson masses and the quark mass, we get the following expressions [9]:

$$\frac{\delta E_D}{E_D} = -0.912 \frac{\delta m_q}{m_q}, \quad (4.1)$$

$$\frac{\delta E_t}{E_t} = -0.980 \frac{\delta m_q}{m_q}, \quad (4.2)$$

$$\frac{\delta E_P}{E_P} = -2.839 \frac{\delta m_q}{m_q}. \quad (4.3)$$

5. CONCLUSION

Defining the parameter K_A as

$$\frac{\delta BE(A)}{BE(A)} = K_A \frac{\delta m_q}{m_q}, \quad (5.1)$$

our results for K_D and K_t are in agreement with [14]. However, for the 1S_0 state the disagreement is considerable, since the signs are different (in our case K_P is negative whereas using the relationship with E_D in [15] it is positive).

For the light nuclei the values of K_A deduced for ^{12}C and ^{16}O agree with those found in [14]. However it is noticeable that for ^7Li our value is larger than the value they reported. These authors claim that the uncertainty can be as large as a factor of two, and our value is within this range.

Our motivation for the study of these variations with quark mass, comes from the possible effects on BBN. In particular, the discrepancy of the abundance of ^7Li with the present baryon to photon ratio, can be linked to the variation of the deuteron binding energy with the quark mass [15]. Our results are summarized in Fig. 2 where the curves correspond to the values of K_D and K_P calculated here (solid line), as well as the values used by Berengut *et al.* [27]. In that reference the range of values for $\frac{\delta m_q}{m_q}$ are chosen such that the Lithium problem can be solved. However, even within that range our results show that such variations in the quark mass (at least for the $p(n, \gamma)d$ reaction) does not yield a significant change in the ^7Li abundance. The reason is that the contribution from the deuteron tends to cancel the one from the 1S_0 state [9]. It is different from [15] because their sign for the 1S_0 is opposite to ours. Therefore their contributions are enhanced. We finally note that it will be necessary to include the variation of the strange quark mass in the calculations, especially with the new findings for the sigma commutators.

ACKNOWLEDGEMENTS

This work was supported by the Australian Research Council through the ARC Centre for Excellence in Particle Physics at the Terascale and through the Australian Laureate Fellowship (FL0992247), as well as by the University of Adelaide.




Figure 2: (Color online) Abundance of ${}^7\text{Li}$ with respect to changes in the quark mass in $p(n, \gamma)d$ calculated in the same way as [27] (dashed-red line) and using our results for K_D and K_P (continuous-blue line).

References

- [1] V. A. Dzuba, V. V. Flambaum and J. K. Webb, Phys. Rev. Lett. **82**, 888 (1999).
- [2] J. K. Webb *et al.*, Phys. Rev. Lett. **82**, 884 (1999).
- [3] M. T. Murphy, J. K. Webb and V. V. Flambaum, Mon. Not. Roy. Astron. Soc. **345**, 609 (2003).
- [4] H. B. Sandvik, J. D. Barrow and J. Magueijo, Phys. Rev. Lett. **88**, 031302 (2002).
- [5] R. Srikanth, H. Chand, P. Petitjean and B. Aracil, Phys. Rev. Lett. **92**, 121302 (2004).
- [6] W. J. Marciano, Phys. Rev. Lett. **52**, 489 (1984). X. Calmet, H. Fritzsch, Eur. Phys. J. **C24**, 639 (2002). P. Langacker, G. Segré, M. J. Strassler, Phys. Lett. **B528**, 121-128 (2002).
- [7] J. D. Bekenstein, Phys. Rev. D **25**, 1527 (1982).
- [8] K. A. Olive *et al.*, Phys. Rev. D **66**, 045022 (2002); B. A. Campbell and K. A. Olive, Phys. Lett. B **345**, 429 (1995).
- [9] M. E. Carrillo-Serrano, I. C. Clöet, K. Tsushima, A. W. Thomas and I. R. Afnan, Phys. Rev. C **87**, 015801 (2013).
- [10] S. R. Beane and M. J. Savage, Nucl. Phys. A **713**, 148 (2003).
- [11] E. Epelbaum, U. -G. Meissner and W. Gloeckle, Nucl. Phys. A **714**, 535 (2003).
- [12] S. R. Beane and M. J. Savage, Nucl. Phys. A **717**, 91 (2003).
- [13] J. Soto and J. Tarrus, Phys. Rev. C **85**, 044001 (2012).
- [14] V. V. Flambaum, R. B. Wiringa, Phys. Rev. **C76**, 054002 (2007).
- [15] V. F. Dmitriev, V. V. Flambaum and J. K. Webb, Phys. Rev. D **69**, 063506 (2004).
- [16] Y. Nambu and G. Jona-Lasinio, Phys. Rev. **122**, 345 (1961).
- [17] W. Bentz and A. W. Thomas, Nucl. Phys. A **696**, 138 (2001).
- [18] I. Caprini, G. Colangelo, H. Leutwyler, Phys. Lett. **96**, 132001 (2006).
- [19] B. Ananthanarayan, G. Colangelo, J. Gasser, H. Leutwyler, Phys. Rep. **353**, 207-297 (2001). S. M. Roy, Phys. Lett. **36B**, 353 (1971).
- [20] Particle Data Group, Phys. Lett. **D66**, 1 (2002).
- [21] P. Gell-Mann, R. J. Oakes, B. Renner, Phys. Rev. **175**, 2195 (1968).
- [22] W. Armour, C. R. Allton, D. B. Leinweber, A. W. Thomas, R. D. Young, J. Phys. G. **32**, 971 (2006); D. B. Leinweber, A. W. Thomas, K. Tsushima, S. V. Wright, Phys. Rev. **D64**, 094502 (2001); C. R. Allton, W. Armour, A. W. Thomas, R. D. Young, Phys. Lett. **B628**, 125-130 (2005).
- [23] P. E. Shanahan, A. W. Thomas and R. D. Young, arXiv:1205.5365 [nucl-th].
- [24] P. A. M. Guichon, Phys. Lett. **B200**, 235 (1988); P. A. M. Guichon, K. Saito, E. N. Rodionov, A. W. Thomas, Nucl. Phys. **A601**, 349 (1996).
- [25] K. Saito, K. Tsushima, A. W. Thomas, Nucl. Phys. **A609**, 339 (1996).
- [26] Roland Bryan, and Bruce L. Scott, Phys. Rev. **177**, 1435 (1969).
- [27] J. C. Berengut, V. V. Flambaum and V. F. Dmitriev, Phys. Lett. B **683**, 114 (2010).