

# PUBLISHED VERSION

Alicia M. Montgomerie, Catherine R. Chittleborough, Anne W. Taylor  
**Physical inactivity and incidence of obesity among South Australian adults**  
PLoS One, 2014; 9(11):e112693-1-e112693-7

© 2014 Montgomerie et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Originally published at:

<http://doi.org/10.1371/journal.pone.0112693>

## PERMISSIONS

<http://creativecommons.org/licenses/by/4.0/>



This is a human-readable summary of (and not a substitute for) the [license](#).

[Disclaimer](#)



### You are free to:

**Share** — copy and redistribute the material in any medium or format

**Adapt** — remix, transform, and build upon the material

for any purpose, even commercially.

The licensor cannot revoke these freedoms as long as you follow the license terms.

### Under the following terms:



**Attribution** — You must give **appropriate credit**, provide a link to the license, and **indicate if changes were made**. You may do so in any reasonable manner, but not in any way that suggests the licensor endorses you or your use.

**No additional restrictions** — You may not apply legal terms or **technological measures** that legally restrict others from doing anything the license permits.

<http://hdl.handle.net/2440/87765>



# Physical Inactivity and Incidence of Obesity among South Australian Adults

Alicia M. Montgomerie<sup>1\*</sup>, Catherine R. Chittleborough<sup>2</sup>, Anne W. Taylor<sup>1</sup>

**1** Discipline of Medicine, School of Medicine, The University of Adelaide, Adelaide, Australia, **2** Discipline of Public Health, School of Population Health, The University of Adelaide, Adelaide, Australia

## Abstract

The aim of this paper is to examine the association of physical inactivity with incidence of obesity in the South Australian adult population. Two representative data sources were used – the South Australian Monitoring and Surveillance System (SAMSS), a monthly surveillance system, and the North West Adelaide Health Study (NWAHS), a biomedical cohort study. There were 75.3% (n = 12873) SAMSS participants and 72.8% (n = 1521) of NWAHS participants that were not obese at baseline. The cumulative incidence of obesity for SAMSS participants from the previous year to the current year was 2.7%. The cumulative incidence of obesity for NWAHS participants between baseline and stage 3 was 14.4%. Physical inactivity was associated with incident obesity (RR 1.48, 95% CI 1.14–1.90 [SAMSS] and RR 1.41, 95% CI 1.03–1.93 [NWAHS]). This association remained, but was attenuated after adjustment for chronic conditions, risk factors and socio-demographic factors. However, physical activity should be continued to be encouraged in the population for its known additional health benefits.

**Citation:** Montgomerie AM, Chittleborough CR, Taylor AW (2014) Physical Inactivity and Incidence of Obesity among South Australian Adults. PLoS ONE 9(11): e112693. doi:10.1371/journal.pone.0112693

**Editor:** Julian W. Tang, Alberta Provincial Laboratory for Public Health/University of Alberta, Canada

**Received:** July 8, 2014; **Accepted:** October 12, 2014; **Published:** November 10, 2014

**Copyright:** © 2014 Montgomerie et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Data Availability:** The authors confirm that, for approved reasons, some access restrictions apply to the data underlying the findings. Data from the South Australian Monitoring and Surveillance System (SAMSS) and North West Adelaide Healthy Study (NWAHS) were accessed from a third party. SAMSS is owned by the Government of South Australia, SA Health. To gain access to the data for this manuscript ethics approval was sought and granted. NWAHS data are available from Janet Grant on janet.grant@adelaide.edu.au and SAMSS data are available from Eleonora DalGrande on eleonora.dalgrande@adelaide.edu.au.

**Funding:** The authors have no support or funding to report.

**Competing Interests:** The authors have declared that no competing interests exist.

\* Email: alicia.montgomerie@adelaide.edu.au

## Introduction

Obesity is a major public health concern with the prevalence rates doubling and even tripling in many countries in the last three decades [1]. The prevalence of obesity in Australian adults was 28.3% in 2011–2012 [2]. Obesity has been linked to an increased risk of premature death and conditions that have an adverse effect on quality of life [3]. These conditions include diabetes, cardiovascular diseases, musculoskeletal disorders, stroke, some cancers, a number of respiratory and gastrointestinal diseases, osteoarthritis, and some mental health conditions [4–6]. Health risks associated with obesity impose a considerable economic burden on individuals, families and communities, with the total annual direct cost of overweight and obesity in Australia estimated to be \$58.2 billion in 2008 [7]. In recognition of this, the Australian Government endorsed obesity as a national health priority area in 2008 [8].

There is evidence that the increase in obesity prevalence is associated with a decrease in physical activity [9–11]. The overall prevalence of Australian adults reporting low or sedentary levels of physical activity was 66.9% in 2011–12 [12]. In 2009 it was reported that 27% of physically inactive Australian adults were obese, compared to only 16% of people undertaking a high level of physical activity [13]. The Australian Diabetes, Obesity and Lifestyle Study (AusDiab) reported a positive association between obesity and lower physical activity [14]. Previous cross-sectional studies have also reported an inverse relationship between physical

activity and body mass index (BMI) [15–20]. A recent study among US adults reported that obesity incidence was significantly higher among adults who did not participate in leisure time activity [21]. However, these studies are cross-sectional and causality cannot be determined from the observed association. From these studies, it is not clear whether physical inactivity causes obesity or whether physical inactivity is a consequence of obesity. A longitudinal study of adults in Copenhagen reported that there was no evidence that physical inactivity promotes the development of obesity [22]. Evidence of the association of physical inactivity and the development of obesity from other prospective observational population studies that measure physical activity at baseline, are few and give inconsistent results [23–26].

There are currently no nationally representative longitudinal studies on obesity and its association with physical inactivity, and more specifically there are no studies on the incidence of obesity and physical inactivity within the South Australian population. Prevalence data are important for conveying the magnitude of the obesity problem, but incidence conveys information about the rate of its development. By examining the association of physical inactivity with incident cases of obesity, we can be clearer about the causal nature of physical inactivity in the development of obesity.

The aim of this paper is to examine the association of physical inactivity with incidence of obesity in the South Australian adult population. Providing evidence for the link between physical inactivity and incident obesity may strengthen the argument for

improving physical activity among Australians, in an effort to prevent people becoming obese. We used two data sources to examine this association: one cross-sectional which relies on retrospective, self-reported data to establish obesity incidence, and the other a longitudinal cohort study following the same people prospectively over time.

## Methods

### Samples

This paper used two existing representative South Australian data sources of adults aged 18 years and over. The first data source is the South Australian Monitoring and Surveillance System (SAMSS); a monthly chronic disease and risk factor telephone survey of randomly selected people established in July 2002 [27]. All households in South Australia with a telephone number listed in the Electronic White Pages are eligible for selection in the sample. Within each household, the person who had a birthday last is selected for interview. There is no replacement for non-contactable persons. Data are collected by a contracted agency using Computer Assisted Telephone Interviewing (CATI) and interviews are conducted in English. Detailed SAMSS methodology has been published elsewhere [28]. We used a weighted SAMSS sample from data collected in the period January 2008 to December 2011 from participants aged 18 years and over ( $n = 22,535$ ). The response rate of SAMSS for this period was between 60% and 65% each month.

The second data source is the North West Adelaide Health Study (NWAHS), a representative population cohort study of 4056 randomly selected adults aged 18 years and over recruited from the northern and western regions of Adelaide. This region represents approximately half of the metropolitan area and one-third of the population in South Australia [29]. Detailed methodology of the study has been previously reported [30]. All households in the north-western area of Adelaide with a telephone connected and the telephone number listed in the Electronic White Pages were eligible for selection. Within each household, the person who had their birthday last and was 18 years or older, was selected for interview and invited to attend the clinic. Baseline assessment (Stage 1) of the cohort occurred between December 1999 and July 2003 with 4056 adults participating in the clinical assessment, resulting in an overall response rate of 49.4%. The first follow-up (Stage 2) of the cohort occurred between May 2004 and February 2006 with survey data collected for 88% ( $n = 3574$ ) and clinic data for 79% ( $n = 3206$ ). The second follow-up (Stage 3) of the cohort occurred between June 2008 and August 2010, with survey data collected for 67% ( $n = 2710$ ) and clinic data for 61% ( $n = 2487$ ) of eligible participants. The same survey methodologies were used at all stages of the study. Self-reported demographics and health-related information was collected in the CATI survey and a self-completed questionnaire. Biomedical data were obtained at the clinic appointment.

### Obesity

SAMSS participants were asked, 'What is your height without shoes?', 'What is your weight? (Undressed in the morning)' and 'How much did you weigh a year ago?' Participants in the NWAHS cohort had their height measured without shoes using a wall mounted stadiometer in centimetres to the nearest 0.5 centimetre and weight measured on calibrated scales in kilograms to the nearest 0.1 kilogram. BMI was calculated as weight in kilograms divided by the square of height in metres. Current BMI for SAMSS participants was calculated from self-reported current weight and height, and BMI in the previous year was calculated

from self-reported weight in the previous year and current self-reported height. BMI was categorised according to the World Health Organization criteria [5]. Participants with a BMI greater than or equal to  $30 \text{ kg/m}^2$  were classified as obese [5].

### Physical inactivity

SAMSS participants were asked to recall physical activity behaviours in the past week on the time they spent undertaking walking, moderate and vigorous activity for sport, recreation or fitness from items from the validated Active Australia Questionnaire [AAQ] [31–34]. Moderate exercise was defined as activity undertaken for fitness, recreation or sport that causes a moderate increase in heart rate or breathing, while vigorous exercise was defined as causing a large increase in a person's heart rate or breathing. The National Physical Activity Guidelines for Australians recommends accumulating 30 minutes of moderate or greater intensity physical activity on most days of the week, translated as 150 minutes per week [35]. Sufficient Physical Activity (SPA) can thus be defined as the completion of 150 minutes of walking, moderate and vigorous activity (with vigorous multiplied by two to account for its greater intensity) in the past week [31]. Participants not achieving SPA were classified as physically inactive. In the NWAHS at baseline, physical activity was assessed from the self-completed questionnaire using items from the Australian Bureau of Statistics (ABS) National Health Survey that required participants to recall physical activity behaviours in the past two weeks [36]. Questions focused on the typical frequency, intensity and duration of walking for sport, recreation or fitness, moderate and vigorous exercise, where moderate exercise was any exercise which caused a moderate increase in heart rate or breathing and vigorous exercise was any exercise which caused a large increase in heart rate or breathing. From these components, an exercise score was calculated by frequency of exercise sessions multiplied by the average time per session multiplied by intensity, where walking has an intensity score of 3.5, with 5.0 for moderate exercise and 7.5 for vigorous exercise. Scores were classified as sedentary (below 100), low (between 100 and below 1600), moderate (scores from 1600 to 3200, or greater than 3200 and less than two hours of vigorous exercise) and high (over 3200 and more than two hours of vigorous exercise), as determined by the ABS [33]. Participants with a score of sedentary or low were classified as physically inactive.

### Covariates

In both SAMSS and NWAHS, the socio-demographic characteristics included were age (in years), sex, country of birth (born in Australia; not born in Australia), highest educational attainment (Bachelor degree or higher; trade, apprenticeship, certificate or diploma; secondary education or less), marital status (married or living with partner; separated, divorced, widowed, never married), gross annual household income (more than \$60,000; less than or equal to \$60,000), work status (employed; not employed), and socioeconomic status as measured by the Socio Economic Indexes for Areas (SEIFA) Index of Relative Disadvantage [37]. SEIFA was dichotomised into most advantaged (middle, high or highest quintile) and most disadvantaged (lowest or low quintile). For NWAHS analyses, region (western; northern) was also included.

For NWAHS participants, diabetes was defined as having fasting plasma glucose of at least 7.0 mmol/L, or reporting having been told by a doctor that they have diabetes in the self-completed questionnaire [38]. Asthma was based on three pre- and three post-salbutamol measurements, and defined as at least a 12% increase in FEV1 (forced expiratory volume in one second) from

pre-Ventolin to post-Ventolin if the absolute difference in FEV1 was greater than 200 ml, or an absolute change in FEV1 of greater than or equal to 400 ml [39], or if participants reported having been told by a doctor that they have asthma. COPD was defined as a post bronchodilator FEV1:FVC ratio of less than 70% [40]. For SAMSS participants self-reported doctor diagnosed health conditions included diabetes, arthritis, osteoporosis, asthma (diagnosed with asthma by a doctor and experienced symptoms and/or treatment in the last 12 months [41]), and chronic obstructive pulmonary disease (COPD, chronic bronchitis or emphysema). For both NWAHS and SAMSS participants cardiovascular disease (CVD) was defined as being told by a doctor that they had ever had a heart attack, a stroke or angina and current mental health condition was defined as a doctor diagnosis of anxiety, depression, a stress related problem or any other mental health problem in the last twelve months.

Risk factors included smoking (current smokers; non- or ex-smokers), alcohol consumption, high cholesterol, high blood pressure and self-reported general health status (NWAHS and SAMSS), as well as fruit, vegetable and fast food consumption and psychological distress (SAMSS only). Alcohol consumption was classified according to the National Health and Medical Research Council guidelines for the risks associated with consuming alcohol, where lifetime risk of harm from alcohol-related disease or injury was those participants who drank two or more standard drinks in one occasion and for alcohol related injury risk, those participants who drank four or more standard drinks in one occasion [42]. High cholesterol for NWAHS participants was determined from a fasting blood sample taken at the study clinic and defined as total blood cholesterol greater than or equal to 5.5 mmol/L [43]. SAMSS participants were classified as having high cholesterol if they reported having been told by a doctor that they had high cholesterol. High blood pressure for NWAHS was defined as systolic blood pressure greater than or equal to 140 mmHg and/or diastolic blood pressure greater than or equal to 90 mmHg [44]. SAMSS participants were classified as having high blood pressure if they reported having been told by a doctor that they had high blood pressure and/or if they were on antihypertensive treatment [45]. Self-reported general health status [46] asked participants to rate their general health as excellent, very good, good, fair or poor; dichotomised into excellent, very good or good, vs. fair or poor. Insufficient daily vegetable or fruit consumption was defined as having consumed less than five serves of vegetables, or less than two serves of fruit (SAMSS). High fast food consumption was classified as more than one serve of fast food per week (SAMSS). SAMSS participants were classified as having high or very high levels of psychological distress if they had a score of 22 or above on the Kessler 10 Psychological Distress Scale [47].

### Data analysis

SAMSS data were weighted by age, sex, area and probability of selection in the household to estimated resident population data so that the results were representative of the South Australian population. We excluded participants with missing data for height ( $n = 380$ ), current ( $n = 856$ ) or previous weight ( $n = 419$ ), those who reported a height  $\geq 215$  cm ( $n = 1$ ) or  $< 100$  cm ( $n = 2$ ), those who had a previous high BMI ( $> 75$  kg/m<sup>2</sup>,  $n = 1$ ), those who lost  $\geq 90$  kg ( $n = 5$ ) and those missing data for at least one covariate ( $n = 2998$ ). This yielded a final study size of  $n = 17,085$  (Figure 1).

NWAHS data were weighted to the Australian Bureau of Statistics' Census [48] and Estimated Residential Population for South Australia [49], by region, age group, sex, and probability of selection in the household, to provide population representative estimates. We excluded participants who were missing data on

weight ( $n = 2$ ), those that did not complete stage 3 ( $n = 1684$ , of which  $n = 208$  had died) and those missing data for at least one covariate ( $n = 282$ ). This yielded a final study size of  $n = 2088$  (Figure 1).

Data were analysed using Statistical Package for the Social Sciences (SPSS) version 19 for Windows (Chicago, IL) and STATA version 10.0 (College Station, TX). Cumulative incidence of obesity in SAMSS was calculated by dividing the number of participants who were currently obese (BMI  $\geq 30$  kg/m<sup>2</sup>) by the number of participants who were not obese (BMI  $< 30$  kg/m<sup>2</sup>) one year ago. Cumulative incidence of obesity in NWAHS was calculated by dividing the number of participants who became obese at Stage 3 by the number of participants who were not obese at baseline (Stage 1). Incidence of obesity was also reported per 1000 person years, taking into account the different follow-up time periods in SAMSS (12 months) and NWAHS (mean follow-up time 2898.9 days, SD [Standard Deviation] 402.29).


Generalised linear regression models using the log link were used to estimate the relative risk (RR) for the association of physical inactivity and incident obesity. Model 1 was adjusted for age and sex. Model 2a was adjusted for age, sex and chronic conditions (diabetes, asthma, COPD, cardiovascular disease, mental health) and model 2b was adjusted for age, sex and chronic conditions as in model 2a and also adding arthritis and osteoporosis from SAMSS. Model 3a was adjusted for age, sex, chronic conditions and risk factors (smoking, alcohol, SF-1, blood pressure and cholesterol) and model 3b was adjusted for age, sex, chronic conditions and risk factors as in model 3a and also adding vegetable consumption, fruit consumption, fast food consumption and psychological distress from SAMSS. Model 4a was adjusted for age, sex, chronic conditions, risk factors and demographics (SEIFA, country of birth, education, marital status, income and work status) and model 4b was adjusted for age, sex, chronic conditions, risk factors and demographics as in model 4a and region from NWAHS.

Ethics approval was granted by the Government of South Australia SA Health Human Research Ethics Committee (HREC/13/SAH/44).

### Results

The baseline characteristics of NWAHS and SAMSS study participants who were obese or not obese are described in Table 1. In the baseline SAMSS sample one year before the interview, 24.7% ( $n = 4212$ ) were classified as obese (BMI  $\geq 30$  kg/m<sup>2</sup>), 35.6% ( $n = 6079$ ) were classified as overweight (BMI 25.0–29.9 kg/m<sup>2</sup>), 37.4% ( $n = 6397$ ) were classified as normal weight (BMI 18.50–24.99 kg/m<sup>2</sup>) and 2.3% ( $n = 397$ ) were classified as underweight (BMI  $< 18.50$  kg/m<sup>2</sup>). At baseline, 27.2% ( $n = 567$ ) of NWAHS participants were classified as obese (BMI  $\geq 30$  kg/m<sup>2</sup>), 38.3% ( $n = 799$ ) were classified as overweight (BMI 25.0–29.9 kg/m<sup>2</sup>), 33.8% ( $n = 706$ ) were classified as normal weight (BMI 18.50–24.99 kg/m<sup>2</sup>) and 0.8% ( $n = 16$ ) were classified as underweight (BMI  $< 18.50$  kg/m<sup>2</sup>).

There were 75.3% ( $n = 12873$ ) of SAMSS participants who were not obese at baseline and 72.8% ( $n = 1521$ ) of NWAHS participants who were not obese at baseline. The cumulative incidence of obesity for SAMSS participants from the previous year to the current year was 2.7% (Table 2). The cumulative incidence of obesity for NWAHS participants between baseline and Stage 3 was 14.4% (Table 2). Incidence of obesity was higher among those who were physically inactive (SAMSS 3.3%; NWAHS 16.3%) compared to those who were physically active (SAMSS 2.3%; NWAHS 11.5%) (Table 2). Weight gain among



**Figure 1. South Australian Monitoring and Surveillance System (SAMSS) and North West Adelaide Health Study (NWAHS) study sample.**

doi:10.1371/journal.pone.0112693.g001

**Table 1. Socio-demographic characteristics of participants who were obese and not obese, South Australian Monitoring and Surveillance System (SAMSS) and North West Adelaide Health Study (NWAHS).**

	SAMSS			NWAHS		
	% or mean ± SD			% or mean ± SD		
	Overall (n = 17085)	Not obese one year ago (n = 12873)	Obese one year ago (n = 4212)	Overall (n = 2088)	Not obese at baseline (n = 1521)	Obese at baseline (n = 567)
Age (years)	48.4±17.06	47.9±17.71	49.6±14.81	45.5±15.78	44.6±16.22	47.8±14.28
Male (%)	50.6	51.1	49.1	49.2	50.6	45.6
Lowest and low quintile (IRSD) (%)	36.1	33.4	44.4	54.4	51.9	61.3
Born in Australia (%)	79.2	78.7	80.8	71.2	73.1	70.5
Secondary education or less (%)	47.7	46.5	51.4	42.1	41.1	44.9
Married (%)	71.3	70.4	73.9	69.6	68.4	72.8
Household income ≤ \$60,000 (%)	43.4	41.7	48.4	71.0	69.0	76.2
Employed (%)	66.1	66.5	64.7	62.6	64.8	56.8
Western region (%) (NWAHS only)				44.8	48.1	36.2

SD: Standard Deviation, IRSD: Index of Relative Disadvantage

Note: Socio-demographic characteristics are current for SAMSS participants and baseline for NWAHS participants. Obesity status is reported one year ago for SAMSS participants and measured at baseline for NWAHS participants.

doi:10.1371/journal.pone.0112693.t001

**Table 2.** Obesity incidence overall, and by physical activity, South Australian Monitoring and Surveillance System (SAMSS) and North West Adelaide Health Study (NWAHS).

	n	events	Cumulative incidence (95% CI)	Incidence rate per 1000 person-years
SAMSS Obesity	12873	353	2.7 (2.5–3.0)	27.4
Physically active	7209	163	2.3 (1.9–2.6)	22.7
Physically inactive	5664	189	3.3 (2.9–3.8)	33.5
NWAHS Obesity	1521	219	14.4 (12.7–16.3)	18.1
Physically active	607	70	11.5 (9.2–14.3)	15.0
Physically inactive	914	149	16.3 (14.0–18.8)	20.2

CI: Confidence Interval.

doi:10.1371/journal.pone.0112693.t002

SAMSS participants who became obese ( $n = 353$ ) was greater (mean 9.2 kg, SD 6.99) than among the participants who did not become obese ( $n = 12,520$ , mean 0.3 kg, SD 3.72). Weight gain among NWAHS participants who became obese ( $n = 219$ ) was greater (11.7 kg, SD 8.55) than among the participants who did not become obese ( $n = 1302$ , 1.2 kg, SD 5.98).

Physical inactivity was associated with incident obesity (RR 1.48, 95% CI 1.14–1.90 [SAMSS] and RR 1.41, 95% CI 1.03–1.93 [NWAHS], Table 3). This association remained, but was attenuated, after adjustment for chronic conditions, risk factors and socio-demographic factors.

## Discussion

In these two representative population studies of South Australia adults, we found that physical inactivity is associated with incident obesity but this association is attenuated when adjusted for confounders. Physical inactivity was associated with incident obesity using both data sources. The association was attenuated when the models were adjusted for risk factors and socio-demographic factors. The relative risks for the association of

physical inactivity and incident obesity are similar in each model for both studies. Smaller P-values observed with the SAMSS data could be explained by the larger sample size. By including a retrospective question of weight in SAMSS, a surveillance system, we are able to continually monitor and identify those at high risk of becoming obese to target prevention efforts.

The attenuation in the model including fruit, vegetable and fast food consumption using SAMSS data is consistent with recent studies reporting that weight gain is explained more by increased energy intake than physical inactivity [50,51]. Other studies have also demonstrated associations between physical inactivity and obesity but do not provide evidence that physical inactivity promotes the development of obesity [22,52–54]. More recently, results from the Whitehall II study reported that there was no association between sitting time and incident obesity [55].

The difference in incidence rate per 1000 person years between SAMSS and NWAHS may be explained by the fact that NWAHS participants and their nominated general practitioner received a result letter after each clinic assessment. The letter highlighted any results outside the healthy range, and indicated that the desirable

**Table 3.** Association of physical inactivity with incident obesity, South Australian Monitoring and Surveillance System (SAMSS) and North West Adelaide Health Study (NWAHS).

	SAMSS (n = 12873)		NWAHS (n = 1521)	
	RR (95% CI)	P-value	RR (95% CI)	P-value
Basic model (unadjusted)	1.48 (1.14–1.90)	0.003	1.41 (1.03–1.93)	0.031
Model 1 <sup>a</sup>	1.51 (1.17–1.95)	0.002	1.39 (1.02–1.92)	0.039
Model 2a <sup>b</sup>	1.43 (1.10–1.85)	0.007	1.42 (1.03–1.95)	0.030
Model 2b <sup>c</sup>	1.36 (1.05–1.77)	0.022	-	-
Model 3a <sup>d</sup>	1.30 (1.00–1.69)	0.050	1.32 (0.96–1.81)	0.093
Model 3b <sup>e</sup>	1.24 (0.94–1.64)	0.125	-	-
Model 4a <sup>f</sup>	1.26 (0.97–1.64)	0.083	1.27 (0.15–1.76)	0.920
Model 4b <sup>g</sup>	-	-	1.26 (0.91–1.73)	0.164

RR: relative risk. CI: Confidence Interval.

<sup>a</sup>Adjusted for age and sex.<sup>b</sup>Adjusted by age, sex and chronic conditions (diabetes, asthma, COPD, cardiovascular disease and mental health).<sup>c</sup>Adjusted by age, sex and chronic conditions (diabetes, asthma, COPD, cardiovascular disease, mental health, arthritis [SAMSS only] and osteoporosis [SAMSS only]).<sup>d</sup>Adjusted for age, sex, chronic conditions and risk factors (smoking, alcohol, SF-1, blood pressure and cholesterol).<sup>e</sup>Adjusted for age, sex, chronic conditions and risk factors (smoking, alcohol, SF-1, blood pressure, cholesterol, vegetable consumption [SAMSS only], fruit consumption [SAMSS only], fast food consumption [SAMSS only] and psychological distress [SAMSS only]).<sup>f</sup>Adjusted for age, sex, chronic conditions, risk factors and sociodemographics (SEIFA, country of birth, education, marital status, income, work status).<sup>g</sup>Adjusted for age, sex, chronic conditions, risk factors and sociodemographics (SEIFA, country of birth, education, marital status, income, work status and region [NWAHS only]).

doi:10.1371/journal.pone.0112693.t003

level for BMI should be between 18.50 and 24.99. This feedback on their health may have made participants more aware of their weight and prompted them to make changes in order to prevent them becoming obese. In the NWAHS there were more obese participants at baseline than SAMSS participants which may also explain the difference in the incidence rate. The cumulative incidence for SAMSS was consistent with a similar study in the US (4.0%) using the same retrospective weight question [21]. The cumulative incidence for NWAHS was consistent with another prospective cohort study (15.3%) with that had a median follow-up of 6.4 years [56].

A strength of this study is that it utilised two South Australian data sources of adults aged 18 years and over to examine the association of physical inactivity and incident obesity. Both studies have a large random sample, and participants are representative of the whole State (SAMSS) or a large region of South Australia. In addition, the NWAHS has measured height and weight and other biomedical measures at more than one time point.

A limitation of this study is that BMI for SAMSS participants was based on self-reported weight and height, and previous studies have shown that height is generally over-estimated and weight is under-estimated [57,58]. SAMSS participants were also asked to recall weight in the past year, however self-reported past weight has been shown to be a reasonably reliable measure of past body weight [59–62]. The samples were restricted to participants with complete data on height, weight and covariates. There was also loss to follow-up for NWAHS participants, although a proportion ( $n = 208$ ) of the attrition in this sample is mortality related. This attrition could cause selection bias if the relationship between physical inactivity and obesity is different between participants and non-participants [63] but we have no reason to suggest that this may be the case. We also made the assumption that current health behaviours and socio-demographic information of SAMSS participants accurately reflected their health behaviours and socio-demographic information in the previous year. The recall periods for physical activity behaviours differed between the two studies, with NWAHS participants asked about physical activity undertaken in the last two weeks, whereas SAMSS participants were asked about physical activity undertaken in the past week. Self-

reported measures of physical activity are limited due to recall error, and social desirability; however the items have been shown to have acceptable reliability and validity for population level physical activity [32–34]. Physical activity in these two studies only includes exercise for sport, recreation or fitness, and therefore is not necessarily indicative of total physical activity; for example it excludes physical activity at work. The studies also do not take into account the time participants spent being sedentary. A recent study has found that time spent sitting for prolonged periods may not only be detrimental to people's health but may also counteract the benefits of regular moderate to vigorous physical activity [64]. The studies also lack data on participants' complete dietary intake, so only limited dietary factors could be included in our models.

## Conclusion

In conclusion, we found that physical inactivity is associated with incident obesity but this association is attenuated when adjusted for confounders. Further study is needed to take into account sedentary time, physical activity at work and dietary behaviours and incident obesity. Physical activity should, however, continue to be encouraged in the population for its known additional health benefits [65].

## Acknowledgments

SAMSS is owned by SA Health, South Australia, Australia. All collected source data are maintained and managed by Population Research and Outcomes Studies, University of Adelaide. The opinions expressed in this work are those of the authors and may not represent the position or policy of SA Health. We are most grateful to the clinic and recruiting staff for their enormous contribution to the success of the study, and for the generosity of the NWAHS participants in the giving of their time and effort.

## Author Contributions

Conceived and designed the experiments: AM CC AT. Analyzed the data: AT CC. Contributed reagents/materials/analysis tools: AM CC AT. Wrote the paper: AM AT CC.

## References

- Walls HL, Backholer K, Proietto J, McNeil JJ (2012) Obesity and trends in life expectancy. *Journal of Obesity* 2012.
- Australian Bureau of Statistics (2012) National Health Survey: First Results, 2011–12. Overweight and Obesity. Canberra.
- Laforce G, De Looper M, Balestat G (2009) Health at a glance 2009: OECD indicators. OECD Publishing, Paris.
- Guh D, Zhang W, Bansback N, Amarsi Z, Birmingham CL, et al. (2009) The incidence of co-morbidities related to obesity and overweight: a systematic review and meta-analysis. *BMC Public Health* 9: 88.
- World Health Organization (2000) Obesity: preventing and managing the global epidemic. World Health Organization.
- World Health Organization (2003) Diet, nutrition and the prevention of chronic diseases. Geneva; 2003. WHO technical report series 916.
- Economics A, Australia D (2008) The growing cost of obesity in 2008: three years on: Access Economics.
- Australian Institute of Health and Welfare (2012) National Health Priority Area. Canberra: Australian Institute of Health and Welfare.
- Dipietro L (1999) Physical activity in the prevention of obesity: current evidence and research issues. *Medicine and Science in Sports and Exercise* 31: S542–S546.
- Hill JO, Wyatt HR (2005) Role of physical activity in preventing and treating obesity. *Journal of Applied Physiology* 99: 765–770.
- Wareham NJ, van Sluijs EMF, Ekelund U (2005) Physical activity and obesity prevention: a review of the current evidence. *Proceedings of the Nutrition Society* 64: 229–248.
- Australian Bureau of Statistics (2012) National Health survey: Profiles of Health, Australia 2011–2013. Canberra: Australian Bureau of Statistics.
- Australian Bureau of Statistics (2009) National Health survey: Australian Social Trends, December 2009 – Smoking, risky drink and obesity. Canberra: Australian Bureau of Statistics.
- Cameron AJ, Welborn TA, Zimmet PZ, Dunstan DW, Owen N, et al. (2003) Overweight and obesity in Australia: the 1999–2000 Australian diabetes, obesity and lifestyle study (AusDiab). *Medical Journal of Australia* 178: 427–432.
- Ball K, Owen N, Salmon J, Bauman A, Gore C (2001) Associations of physical activity with body weight and fat in men and women. *International Journal of Obesity* 25: 914–919.
- Bernstein MS, Costanza MC, Morabia A (2004) Association of physical activity intensity levels with overweight and obesity in a population-based sample of adults. *Preventive Medicine* 38: 94–104.
- Chan CB, Spangler E, Valcour J, Tudor-Locke C (2003) Cross-sectional Relationship of Pedometer-Determined Ambulatory Activity to Indicators of Health. *Obesity Research* 11: 1563–1570.
- Cooper AR, Page A, Fox KR, Misson J (2000) Physical activity patterns in normal, overweight and obese individuals using minute-by-minute accelerometry. *European Journal of Clinical Nutrition* 54: 887.
- Ewing R, Schmid T, Killingsworth R, Zlot A, Raudenbush S (2003) Relationship between urban sprawl and physical activity, obesity, and morbidity. *American Journal of Health Promotion* 18: 47–57.
- Thompson DL, Rakow J, Perdue SM (2004) Relationship between accumulated walking and body composition in middle-aged women. *Medicine and Science in Sports and Exercise* 36: 911–914.
- Pan L, Freedman DS, Gillespie C, Park S, Sherry B (2011) Incidences of obesity and extreme obesity among US adults: findings from the 2009 Behavioral Risk Factor Surveillance System. *Population Health Metrics* 9: 56.

22. Petersen L, Schnohr P, Sørensen TIA (2003) Longitudinal study of the long-term relation between physical activity and obesity in adults. *International Journal of Obesity* 28: 105–112.
23. Erlichman J, Kerbey AL, James WPT (2002) Physical activity and its impact on health outcomes. Paper 2: Prevention of unhealthy weight gain and obesity by physical activity: an analysis of the evidence. *Obesity Reviews* 3: 273–287.
24. Fogelholm M, Kukkonen-Harjula K (2000) Does physical activity prevent weight gain—a systematic review. *Obesity Reviews* 1: 95–111.
25. Lee IM, Djoussé L, Sesso HD, Wang L, Buring JE (2010) Physical activity and weight gain prevention. *JAMA: the journal of the American Medical Association* 303: 1173–1179.
26. Saris WHM, Blair SN, Van Baak MA, Eaton SB, Davies PSW, et al. (2003) How much physical activity is enough to prevent unhealthy weight gain? Outcome of the IASO 1st Stock Conference and consensus statement. *Obesity Reviews* 4: 101–114.
27. Unit PRaOS (2002) South Australian Monitoring and Surveillance System (SAMSS). Brief Report. No. 2002-20. Adelaide: Population Research and Outcomes Studies Unit.
28. Taylor A, Dal Grande E (2008) Chronic disease and risk factor surveillance using the SA Monitoring and Surveillance System (SAMSS)—history, results and future challenges. *Public Health Bulletin SA* 5: 17–21.
29. Grant JF, Chittleborough CR, Taylor AW, Dal Grande E, Wilson DH, et al. (2006) The North West Adelaide Health Study: detailed methods and baseline segmentation of a cohort for selected chronic diseases. *Epidemiologic Perspectives & Innovations* 3: 4.
30. Grant JF, Taylor AW, Ruffin RE, Wilson DH, Phillips PJ, et al. (2009) Cohort Profile: The North West Adelaide Health Study (NWAHS). *International Journal of Epidemiology* 38: 1479–1486.
31. Australian Institute of Health and Welfare (2003) *The Active Australia Survey: A Guide and Manual for Implementation, Analysis and Reporting*. Australian Institute of Health and Welfare.
32. Heesch KC, Hill RL, Van Uffelen JGZ, Brown WJ (2011) Are Active Australia physical activity questions valid for older adults? *Journal of Science and Medicine in Sport* 14: 233–237.
33. Brown WJ, Burton NW, Marshall AL, Miller YD (2008) Reliability and validity of a modified self-administered version of the Active Australia physical activity survey in a sample of mid-age women. *Australian and New Zealand Journal of Public Health* 32: 535–541.
34. Brown WJ, Trost SG, Bauman A, Mummery K, Owen N (2004) Test-retest reliability of four physical activity measures used in population surveys. *Journal of Science and Medicine in Sport* 7: 205–215.
35. Commonwealth of Australia (1999) *National Physical Activity Guidelines for Adults*. Canberra: Department of Health and Aged Care.
36. McLennan W, Podger AS (1998) *National Nutrition Survey Users' Guide, 1995*. Australian Bureau of Statistics [and] Commonwealth Department of Health and Family Services.
37. Pink B (2006) *Socio-Economic Indexes for Areas (SEIFA)—Technical Paper*. Commonwealth of Australia.
38. World Health Organization (1997) *preventing and managing the global epidemic [report of a WHO Consultation on Obesity]*. Geneva: World Health Organization.
39. Adams RJ, Wilson DH, Appleton S, Taylor A, Dal Grande E, et al. (2003) Underdiagnosed asthma in South Australia. *Thorax* 58: 846–850.
40. Pauwels R (2001) Global initiative for chronic obstructive lung diseases (GOLD): time to act. *European Respiratory Journal* 18: 901–902.
41. Ampon R, Correll P, Marks G, Poulos L, Waters AM (2007) *Asthma in Australia: findings from the 2004–05 National Health Survey*. Canberra: Australian Institute of Health and Welfare.
42. National Health and Medical Research Council (NHMRC) (2009) *Australian Guidelines to Reduce Health Risks from Drinking Alcohol*. Canberra.
43. Australian Institute of Health and Welfare, National Heart Foundation of Australia, National Stroke Foundation of Australia (2001) *Heart, Stroke and Vascular Disease Australian Facts 2001*. Canberra.
44. Bennett SA, Magnus P (1994) Trends in cardiovascular risk factors in Australia. Results from the National Heart Foundation's Risk Factor Prevalence Study, 1980–1989. *The Medical Journal of Australia* 161: 519–527.
45. Australian Institute of Health and Welfare (2004) *Heart, stroke and vascular diseases, Australian facts 2004*. Canberra.
46. Ware Jr JE, Sherbourne CD (1992) The MOS 36-item short-form health survey (SF-36): I. Conceptual framework and item selection. *Medical Care*: 473–483.
47. Kessler RC, Andrews G, Colpe LJ, Hiripi E, Mroczek DK, et al. (2002) Short screening scales to monitor population prevalences and trends in non-specific psychological distress. *Psychological Medicine* 32: 959–976.
48. Australian Bureau of Statistics (2001) *Census of Population and Housing Selected Social and Housing Characteristics for Statistical Local Areas South Australia, 2001*.
49. Australian Bureau of Statistics (2002) *Estimated Resident Population by Age and Sex 1999*. Canberra.
50. Scarborough P, Burg MR, Foster C, Swinburn B, Sacks G, et al. (2011) Increased energy intake entirely accounts for increase in body weight in women but not in men in the UK between 1986 and 2000. *British Journal of Nutrition* 105: 1399.
51. Swinburn B, Sacks G, Ravussin E (2009) Increased food energy supply is more than sufficient to explain the US epidemic of obesity. *The American journal of clinical nutrition* 90: 1453–1456.
52. Brien SE, Katzmarzyk PT, Craig CL, Gauvin L (2007) Physical activity, cardiorespiratory fitness and body mass index as predictors of substantial weight gain and obesity: the Canadian physical activity longitudinal study. *Canadian journal of public health* 98: 121–124.
53. Heitmann BL, Kaprio J, Harris JR, Rissanen A, Korkeila M, et al. (1997) Are genetic determinants of weight gain modified by leisure-time physical activity? A prospective study of Finnish twins. *The American journal of clinical nutrition* 66: 672–678.
54. Raguso CA, Kyle U, Kossovsky MP, Roynette C, Paoloni-Giacobino A, et al. (2006) A 3-year longitudinal study on body composition changes in the elderly: role of physical exercise. *Clinical Nutrition* 25: 573–580.
55. Pulsford RM, Stamatakis E, Britton AR, Brunner EJ, Hillsdon MM (2013) *Sitting Behavior and Obesity: Evidence from the Whitehall II Study*. *American Journal of Preventive Medicine* 44: 132–138.
56. Núñez-Córdoba JM, Bes-Rastrollo M, Pollack KM, Seguí-Gómez M, Beunza JJ, et al. (2013) *Annual Motor Vehicle Travel Distance and Incident Obesity: A Prospective Cohort Study*. *American Journal of Preventive Medicine* 44: 254–259.
57. Taylor AW, Grande ED, Gill TK, Chittleborough CR, Wilson DH, et al. (2006) How valid are self-reported height and weight? A comparison between CATI self-report and clinic measurements using a large cohort study. *Australian and New Zealand Journal of Public Health* 30: 238–246.
58. Australian Bureau of Statistics (1995) *How Australians measure up*. Canberra: ABS.
59. Casey VA, Dwyer JT, Berkey CS, Coleman KA, Gardner J, et al. (1991) Long-term memory of body weight and past weight satisfaction: a longitudinal follow-up study. *The American journal of clinical nutrition* 53: 1493–1498.
60. Klipstein-Grobusch K, Kroke A, Boeing H (1998) Reproducibility of self-reported past body weight. *European Journal of Clinical Nutrition* 52: 525.
61. Kovalchik S (2009) Validity of adult lifetime self-reported body weight. *Public Health Nutrition* 12: 1072.
62. Perry GS, Byers TE, Mokdad AH, Serdula MK, Williamson DF (1995) The validity of self-reports of past body weights by US adults. *Epidemiology* 6: 61–66.
63. Carter KN, Imlach-Gunasekara F, McKenzie SK, Blakely T (2012) Differential loss of participants does not necessarily cause selection bias. *Australian and New Zealand Journal of Public Health* 36: 218–222.
64. Ponsonby A-L, Sun C, Ukoumunne OC, Pezic A, Venn A, et al. (2011) Objectively Measured Physical Activity and the Subsequent Risk of Incident Dysglycemia The Australian Diabetes, Obesity and Lifestyle Study (AusDiab). *Diabetes Care* 34: 1497–1502.
65. Warburton DER, Nicol CW, Bredin SSD (2006) Health benefits of physical activity: the evidence. *Canadian Medical Association Journal* 174: 801–809.