

PUBLISHED VERSION

Alexandre Emidio Ribeiro Silva, Ana Maria Baptista Menezes, Maria Cecília Formoso Assunção, Helen Gonçalves, Flávio Fernando Demarco, Fabiana Vargas-Ferreira, Marco Aurélio Peres

Validation of self-reported information on dental caries in a birth cohort at 18 years of age

PLoS One, 2014; 9(9):e106382-1-e106382-8

© 2014 Silva et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Originally published at:

<http://doi.org/10.1371/journal.pone.0106382>

PERMISSIONS

<http://creativecommons.org/licenses/by/4.0/>



Attribution 4.0 International (CC BY 4.0)

This is a human-readable summary of (and not a substitute for) the [license](#).

[Disclaimer](#)



You are free to:

Share — copy and redistribute the material in any medium or format

Adapt — remix, transform, and build upon the material

for any purpose, even commercially.

The licensor cannot revoke these freedoms as long as you follow the license terms.

Under the following terms:



Attribution — You must give [appropriate credit](#), provide a link to the license, and [indicate if changes were made](#). You may do so in any reasonable manner, but not in any way that suggests the licensor endorses you or your use.

No additional restrictions — You may not apply legal terms or [technological measures](#) that legally restrict others from doing anything the license permits.



Validation of Self-Reported Information on Dental Caries in a Birth Cohort at 18 Years of Age

Alexandre Emídio Ribeiro Silva^{1*}, Ana Maria Baptista Menezes¹, Maria Cecília Formoso Assunção¹, Helen Gonçalves¹, Flávio Fernando Demarco¹, Fabiana Vargas-Ferreira¹, Marco Aurélio Peres²

1 Postgraduate Programme in Epidemiology, Federal University of Pelotas, Pelotas, Rio Grande do Sul, Brazil, **2** Australian Research Centre for Population Oral Health, School of Dentistry, University of Adelaide, Adelaide, Austrália

Abstract

Objective: Estimate the prevalence of dental caries based on clinical examinations and self-reports and compare differences in the prevalence and effect measures between the two methods among 18-year-olds belonging to a 1993 birth cohort in the city of Pelotas, Brazil.

Method: Data on self-reported caries, socio-demographic aspects and oral health behaviour were collected using a questionnaire administered to adolescents aged 18 years ($n=4041$). Clinical caries was evaluated ($n=1014$) by a dentist who had undergone training and calibration exercises. Prevalence rates of clinical and self-reported caries, sensitivity, specificity, positive and negative predictive values, absolute and relative bias, and inflation factors were calculated. Prevalence ratios of dental caries were estimated for each risk factor.

Results: The prevalence of clinical and self-reported caries ($DMFT > 1$) was 66.5% (95%CI: 63.6%–69.3%) and 60.3% (95%CI: 58.8%–61.8%), respectively. Self-reports underestimated the prevalence of dental caries by 9.3% in comparison to clinical evaluations. The analysis of the validity of self-reports regarding the DMFT index indicated high sensitivity (81.8%; 95%CI: 78.7%–84.7%) and specificity (78.1%; 95%CI: 73.3%–82.4%) in relation to the gold standard (clinical evaluation). Both the clinical and self-reported evaluations were associated with gender, schooling and self-rated oral health. Clinical dental caries was associated with visits to the dentist in the previous year. Self-reported dental caries was associated with daily tooth brushing frequency.

Conclusions: Based on the present findings, self-reported information on dental caries using the DMFT index requires further studies prior to its use in the analysis of risk factors, but is valid for population-based health surveys with the aim of planning and monitoring oral health actions directed at adolescents.

Citation: Silva AER, Menezes AMB, Assunção MCF, Gonçalves H, Demarco FF, et al. (2014) Validation of Self-Reported Information on Dental Caries in a Birth Cohort at 18 Years of Age. PLoS ONE 9(9): e106382. doi:10.1371/journal.pone.0106382

Editor: Peter M. Milgrom, University of Washington, United States of America

Received April 18, 2014; **Accepted** July 29, 2014; **Published** September 9, 2014

Copyright: © 2014 Silva et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability: The authors confirm that all data underlying the findings are fully available without restriction. All relevant data are within the paper and its Supporting Information files.

Funding: The cohort study was funded by Wellcome Trust. The initial phases of the cohort were funded by the European Union and the Brazilian agencies National Programme for Centres of Excellence (PRONEX), National Research Council (CNPq) and the Ministry of Health. The providers of funding played no role in the study design, data collection and analysis, decision to publish or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* Email: emidio3@ig.com.br

Introduction

Self-reported health status is considered a valid, acceptable method for the assessment of the prevalence of diseases in the general population, such as hypertension and diabetes [1], as well as risk factors, such as poor diet and lack of physical activity [2]. In the field of dentistry, self-reported information is an economically feasible option for measuring oral health conditions in population-based multidisciplinary surveys and diminishes the need for time-consuming clinical exams [3]. Moreover, self-reports have the potential to be a useful method for monitoring oral health conditions and trends over time, which is important to the planning and evaluation of public health policies. For instance, the Centers for Disease Control and Prevention in the United States of

America use health information acquired from telephone surveys [4].

Health inquiries involving self-reported information indicate that characteristics related to socio-demographic aspects [5] and past disease experience [6] have a direct influence on the knowledge of individuals regarding their health, producing greater or lesser agreement between self-reports and clinically determined data. In the field of oral health, studies indicate an underestimation of periodontal disease in self-reported information. [2,7,8]. A systematic review on this issue demonstrates that the use of self-reported data is inadequate for information on gingivitis, but other self-reported measures of periodontal disease have proven to be valid [1], such as the occurrence of periodontal pockets. However, standardisation is required regarding the different methods used for the acquisition of self-reported information.

The literature demonstrates greater validity in self-reported information related to the use of dentures and the number of teeth in the oral cavity [2,4,5,9–11]. However, some studies indicate a slight underestimation [2] or overestimation [2,7] regarding the number of teeth. For measures related to tooth decay, an underestimate of the number of teeth with carious tissue has been found in self-reports [2] and reports of missing teeth should be used with the proper corrections [10]. However, with regard to the presence/absence of dental caries, studies involving adults indicate a lack of agreement with clinical measures, mainly because individuals are often unable to recognise caries [2,6].

Few studies have been conducted on the validity of self-reported information regarding dental caries in 18-year-olds in countries of low to middle income. Therefore, the aims of the present study were to estimate the prevalence of dental caries based on clinical examinations and self-reports and compare differences in the prevalence and effect measures between the two methods among 18-year-olds belonging to a 1993 birth cohort in the city of Pelotas, Brazil.

Materials and Methods

Description of 1993 Pelotas birth cohort

This was the second birth cohort in the city of Pelotas (southern Brazil) and was separated from the first cohort study by 11 years. The aim was to allow the comparison of mother/child characteristics of the population and changes in the main health indicators as well as to evaluate the influence of factors related to birth and early childhood on health throughout the life cycle. Moreover, it is one of the few population-based studies to investigate oral health.

All live births recorded at hospitals in the city of Pelotas in 1993 from mothers residing in urban areas were included in this cohort. Among the 5265 women who had children in the period, 5249 (99.7%) agreed to participate in the study. Subsamples were revisited when the members of the cohort were 1, 3, and 6 months of age as well as 1, 3, 6, and 9 years of age. In 2004, when the members were 11 years of age, the entire cohort was evaluated; the same occurred in 2008 (15 years of age) and 2011 (18 years of age). The last visit (18 years of age) occurred between September 2011 and April 2012 and all members of the cohort were asked to appear at the Federal University of Pelotas for a clinical examination and the administration of questionnaires. Descriptions of the methods employed during past visits to the members of the cohort are found in previous studies [12,13].

Oral health subsamples of 1993 Pelotas birth cohort

The oral health follow up evaluations were performed with subsamples at six, 12 and 18 years. The subsample at six years was obtained from the follow up of the cohort performed when the children were one year of age, in which all underweight children were included. The oral health subsample at six years involved 359 members of the cohort, including 28.7% of underweight children. This information was weighted to represent the true proportion of live births of underweight children. Another home follow up was performed at 12 years, when 339 members of those visited at six years were located. The description of the oral health follow-up methods at six and 12 years is found in a previous publication [14].

The third follow up occurred at 18 years in a clinical setting during morning and afternoon shifts from Monday to Saturday from September 2011 to March 2012. The oral health subsample at 18 years was formed by adolescents who appeared for evaluations during the four predefined appointments (two in the morning and two in the afternoon) throughout the week, which were scheduled in a randomised fashion. Besides the adolescents

who appeared for the predefined appointments, attempts were made to examine all 359 member members of the cohort that had been evaluated in the previous oral health subsamples. The oral health subsample of the final follow up evaluation consisted of 1014 adolescents, 307 of whom had been part of the subsample at six years of age and 301 of whom had been part of the subsample at 12 years of age.

Oral health exams at 18 years of age

At the 2012 follow up, the oral health exam was performed by a single examiner who had undergone training and calibration exercises using the diagnostic criteria of the World Health Organization [15]. Two calibration exercises were conducted (August and December 2011) with the participation of the researcher of the study and three dentists. The aim of the two exercises was to maintain the reproducibility of the exams, as the fieldwork occurred over approximately eight months. Weighted Kappa coefficients were calculated. The lowest values during the two calibration exercises were 0.78 and 0.83 for intra-examiner and intra-examiner agreement, respectively.

Outcome – Dental Caries

1. Clinical dental caries. A World Health Organization (WHO) periodontal probe and mouth mirror were used for the clinical detection of dental caries based on the WHO criteria [15]. The distal, vestibular, mesial, and lingual/palatine surfaces of the anterior teeth were examined. These same surfaces plus the occlusal surface were examined on the posterior teeth. The overall decayed, missing, and filled teeth (DMFT) index and each of its components were calculated for each participant. The prevalence of dental caries was determined based on the proportion of adolescents with DMFT ≥ 1 .

2. Self-reported dental caries. For the self-reported data on dental caries, a 24-item questionnaire (Figure 1) was administered by trained interviewers to 4041 adolescents of the overall cohort. This questionnaire was first tested on individuals with a similar age to those in the cohort. The aim of obtaining self-reported information was to identify the individuals' knowledge regarding the number of decayed, missing and filled teeth. Based on the findings, the DMFT index was calculated for each participant. As with the clinical exam, the prevalence of dental caries was determined based on the proportion of adolescents with DMFT ≥ 1 .

Exploratory variables

A questionnaire addressing socioeconomic and demographic variables, visits to the dentist, tooth brushing habits, and self-rated oral health was administered to all adolescents in the cohort by the same interviewers who administered the questionnaire on self-reported dental caries. The demographic and socioeconomic variables analysed were gender, adolescent's schooling in number of completed years of study (categorised as ≤ 4 , 5 to 8, 9 to 11, and ≥ 12), and household income in the previous month (sum of the monthly income of all members of the household in Brazilian currency [Real – R\$], converted to a figure based on the minimum Brazilian wage and categorised in tertiles).

Variables related to oral health (visits to the dentist, daily tooth brushing frequency, and self-rated oral health) were obtained from the following questions: 1) Have you visited the dentist since <month> of last year (yes or no); 2) How many times a day do you brush your teeth? (categorised as <2 times a day and ≥ 2 times

THE FOLLOWING QUESTIONS ARE ABOUT THE NUMBER OF TEETH YOU HAVE IN THE UPPER PART AND LOWER PART OF YOUR MOUTH.

1. We have a maximum of 16 natural teeth in the upper part of the mouth, counting both wisdom teeth. How many natural teeth do you have in the upper part of your mouth?

16 15 14 13 12 11 10 9 8 7 6 5 4 3 2 1 0

IF LESS THAN 16 TEETH:

2. Were any teeth pulled because of cavities?

(0) No → GO TO QUESTION 4 (1) Yes

IF YES: 3. How many teeth in the upper part of your mouth were pulled because of cavities? ____ teeth [00=IGN]

4. Do you have any natural tooth with an untreated cavity or hole in the upper part of your mouth?

(0) No → GO TO QUESTION 6 (1) Yes

IF YES: 5. How many natural teeth with an untreated cavity or hole in the upper part of your mouth do you have? ____ teeth [00=IGN]

6. Do you have any natural tooth with a restoration or filling in the upper part of your mouth?

(0) No → GO TO QUESTION 8 (1) Yes

IF YES: 7. How many natural teeth with a restoration or filling in the upper part of your mouth do you have? ____ teeth [00=IGN]

8. Do you have bridges or partial denture in the upper part of your mouth?

(0) No → GO TO QUESTION 10 (1) Yes

IF YES: 9. How many natural teeth have been replaced with a bridge or partial denture in the upper part of your mouth? ____ teeth [00=IGN]

10. Do you have any fixed bridge in the upper part of your mouth?

(0) No → GO TO QUESTION 12 (1) Yes

IF YES: 11. How many natural teeth have been replaced with a fixed bridge in the upper part of your mouth?

____ teeth. [00=IGN]

NOW WE ARE GOING TO TALK ABOUT THE LOWER PART OF YOUR MOUTH

12. We have a maximum of 16 natural teeth in the lower part of the mouth, counting both wisdom teeth. How many natural teeth do you have in the lower part of your mouth?

16 15 14 13 12 11 10 9 8 7 6 5 4 3 2 1 0

IF LESS THAN 16 TEETH:

13. Were any teeth pulled due to cavities?

(0) No → GO TO QUESTION 15 (1) Yes

IF YES: 14. How many teeth in the lower part of your mouth were pulled because of cavities? ____ teeth [00=IGN]

15. Do you have any natural tooth with an untreated cavity or hole in the lower part of your mouth?

(0) No → GO TO QUESTION 17 (1) Yes

IF YES: 16. How many natural teeth with an untreated cavity or hole in the lower part of your mouth do you have? ____ teeth [00=IGN]

17. Do you have any natural tooth with a restoration or filling in the lower part of your mouth?

(0) No → GO TO QUESTION 19 (1) Yes

IF YES: 18. How many natural teeth with a restoration or filling in the lower part of your mouth do you have? ____ teeth [00=IGN]

19. Do you have bridges or partial denture in the lower part of your mouth?

(0) No → GO TO QUESTION 21 (1) Yes

IF YES: 20. How many natural teeth have been replaced with a bridge or partial denture in the lower part of your mouth? ____ teeth [00=IGN]

21. Do you have any fixed bridge in the lower part of your mouth?

(0) No → GO TO QUESTION 23 (1) Yes

IF YES: 22. How many natural teeth have been replaced with a fixed bridge in the lower part of your mouth?

____ teeth. [00=IGN]

IF YOUTH HAS NO TEETH IN UPPER ARCH: 23. Do you wear a complete denture on the upper part of your mouth?

(0) No (1) Yes

IF YOUTH HAS NO TEETH IN LOWER ARCH: 23. Do you wear a complete denture on the lower part of your mouth?

(0) No (1) Yes

Figure 1. Self-reported survey of dental caries. Pelotas, Rio Grande do Sul, Brazil, 2012.
doi:10.1371/journal.pone.0106382.g001

a day); and 3) How would you rate the health of your teeth today? (very good, good, fair, poor, and very poor [subsequently categorised as very good/good, fair, and poor/very poor]).

Statistical analysis

The data were analysed with the aid of the Stata 12.0 program. The chi-square test was used to compare the participants in the subsample at 18 years of age with all members of the original cohort. Descriptive analysis was then performed of the DMFT index as well as the individual decayed, missing, and filled components, with the calculation of mean values and 95% confidence intervals (CI). The following were also calculated: intraclass correlation coefficient (ICC) for the DMFT index as well as the individual decayed, missing, and filled components; prevalence of clinically determined dental caries (gold standard); prevalence of self-reported dental caries; estimates of sensitivity, specificity, and positive and negative predictive values; absolute bias (prevalence of self-reported caries minus gold standard prevalence); relative bias (percentage of underestimation of true prevalence = absolute bias/gold standard prevalence × 100); and inflation factor (gold standard prevalence/self-reported prevalence) [16]. Estimates of sensitivity, specificity, and positive and negative predictive values were stratified by gender and schooling (12 or more years of study *versus* 4 years or less). Prevalence ratios for clinically determined and self-reported dental caries were calculated for risk factors using Poisson regression models with robust variance [17].

Ethical aspects

This study received approval from the Human Research Ethics Committee of the Federal University of Pelotas (Brazil) under process number 67/11. All participants received clarifications regarding the objectives and procedures and agreed to participate by signing a statement of informed consent.

Results

A total of 1014 members of the cohort were evaluated with regard to oral health at 18 years of age, 526 (51.9%) of whom were male and 386 (41.7%) had mothers with five to eight years of schooling. Regarding household income, the first, second, and third tertiles corresponded to less than two times the Brazilian minimum wage, between two and four times the minimum wage, and more than four times the minimum wage, respectively. Table 1 displays the characteristics of the original cohort ($n=5248$) and oral health subsample ($n=1014$). Statistically significant differences were found regarding gender, mother's schooling, and household income, with an increase in the proportion of male subjects, mean mother's schooling, and household income in the subsample.

Both the clinical and self-reported DMFT indices had a median value of 1 decayed, missing or filled tooth and an interquartile interval (Q3–Q1) of 3 DMFT. Regarding the mean values, the clinical DMFT was 2.06, and the self-reported DMFT was 1.75. All mean values of the DMFT components demonstrated the underestimation of self-reported information in comparison to the clinical evaluation. In the analysis of the ICC, which allowed measuring the concordance or reliability of the tools used to compare the clinical and self-reported DMFT and its components, the lowest degree of reliability was found for the decayed component ($ICC=0.43$) and the highest was found for the missing component ($ICC=0.61$) (Table 2).

Table 3 displays the prevalence rates of dental caries and estimates of sensitivity, specificity, positive and negative predictive

values, absolute bias, relative bias, and inflation factor. Self-reports underestimated the prevalence of dental caries by 9.3% in comparison to the clinical evaluation. The results indicate high sensitivity (81.8%) and specificity (78.1%). The positive predictive value of 88% indicates that, among those identified with dental caries, the majority actually had the condition. The negative predictive value indicates that, among those identified as not having dental caries, only 68.6% were confirmed as actually not having the condition. The inflation factor for self-reported dental caries was 1.10.

Table 4 displays the estimates of sensitivity, specificity, and positive and negative predictive values stratified by gender, greater (12 or more years of study) or less schooling (4 years or less), lower (1st tertile) or higher (3rd tertile) household income, and visit to the dentist in the previous year among the adolescents. Sensitivity was higher for female subjects (87.3%), adolescents with a greater level of schooling (82.6%), those with a lower household income (84.5%) and those who visited the dentist in the previous year (81.9%). Specificity was higher for male subjects (84.1%), adolescents with a greater level of schooling (92.6%), those with a higher household income (83.5%) and those who did not visit the dentist in the previous year (78.2%). The confirmation of the diagnosis of dental caries (measured by the positive predictive value) was also higher among male subjects (89.5%), adolescents with a greater level of schooling (90.5%), those with a higher household income (89.3%) and those who visited the dentist in the previous year (89.8%). The confirmation of the absence of the condition (measured by the negative predictive value) was greater among female subjects (71.6%), adolescents with a greater level of schooling (86.2%), those with a lower household income (69.7%) and those who did not visit the dentist in the previous year (72.5%).

Table 5 displays the prevalence ratios according to the independent variables evaluated for the binary DMFT (0/ ≥ 1) based on the clinical and self-reported assessments. Higher prevalence ratios were found for the majority of variables when considering clinically detected dental caries. Adolescent's schooling and household income were inversely associated with clinically detected dental caries. Self-rated oral health was positively associated with both the clinical and self-reported measures of dental caries. Clinically determined dental caries was positively associated with visits to the dentist in the previous year. The self-reported measure of dental caries was inversely associated with daily tooth brushing frequency.

Discussion

The findings of the present study indicate adequate validity in self-reported data regarding dental caries in adolescents in relation to the clinical measure using the DMFT index. Self-reported measures exhibited high sensitivity and specificity.

To adjust the prevalence of self-reported dental caries (which was underestimated by 9.3% in comparison to the clinical evaluation in the present study), the literature proposes the use of correction factors, such as the inflation factor [16], for health surveys that assess self-reported information regarding periodontal disease [18].

Moderate agreement was found between self-reported and clinically determined tooth decay in the present study. Studies on the validation of self-reported oral health information have indicated low agreement regarding the comparison of the decayed component [2]. This may be related to the fact that laypersons are unable to recognise dental caries [2,6] or only perceive the

Table 1. Characteristics of original 1993 birth cohort and oral health subsample of cohort members at 18 years age, Pelotas, Rio Grande do Sul, Brazil, 2012.

	Original cohort		Subsample at 18 years of ages		p-value *
	N	% (95%CI)	N	% (95%CI)	
Gender					
Male	2603	49.6 (48.3; 50.9)	526	52.2 (49.8; 55.3)	<0.001
Female	2645	50.4 (49.1; 51.8)	487	47.8 (44.6; 50.9)	
Mother's schooling (years)					
≤4	1468	28.0 (26.8; 29.2)	228	24.6 (21.8; 27.4)	
5–8	2424	46.24 (44.9; 47.6)	386	41.8 (38.5; 44.9)	<0.001
9–11	923	17.60 (16.6; 18.6)	261	28.4 (25.5; 31.4)	
≥12	427	8.1 (7.4; 8.9)	48	5.3 (3.8; 6.8)	
Household income (based on minimum wage)					
1 st tertile	2226	43.3 (41.9; 44.7)	339	32.7(29.9; 35.7)	
2 nd tertile	1445	28.1 (26.9; 29.4)	335	33.2(30.2; 36.2)	<0.001
3 rd tertile	1466	28.5 (27.3; 29.8)	336	34.0(31.0; 37.0)	

*chi-square test.

doi:10.1371/journal.pone.0106382.t001

presence of the condition when it affects their social relations or when they experience pain [19].

Stronger agreement is described regarding reports of the number of missing and filled teeth [20]. The literature states that the number of self-reported missing teeth in adults and elderly individuals differs from the number determined clinically [10] because individuals have difficulties remembering treatment received years ago. Moreover, the loss of the first permanent molar may have occurred early in life, which further hinders the recollection of such an event. The stronger agreement in the present study regarding the number of missing teeth may be explained by the fact that adolescents are more likely to remember adverse oral conditions in their life. Moreover, missing teeth is a rare occurrence in this group. Indeed, the most recent national oral health surveys in Brazil [21,22] report that adolescents between 15 and 19 years of age have an increasingly fewer number of decayed teeth, which is the main cause of tooth loss in this age group. This decrease in the prevalence of caries is directly related to an improved socioeconomic status, which allows the population access to fluoridated toothpaste, a fluoridated water supply and oral healthcare services [23].

The findings for the filled teeth component are in agreement with those reported in a previous study [2]. The greater agreement between the self-reported and clinical assessments regarding

restored teeth is likely related to the aforementioned reduction in the prevalence of dental caries as well as the change in the oral healthcare model adopted in Brazil, which was previously directed more toward surgical and restorative procedures and currently involves a health vigilance model [24]. This aspect may have contributed to the fact that the adolescents analysed had an average of less than one tooth with carious tissue.

The similar prevalence rates between clinically determined and self-reported dental caries measured by the DMFT index in the present study were likely due to the fact that the clinical exam for dental caries based on the WHO criteria for epidemiological surveys identifies more advanced stages of tooth decay. Another aspect that diminished the possibility of error in self-reported information on DMFT was the fact that all participants were informed regarding the maximum number of teeth in the upper and lower arches, which differs from methods reported in previous studies [2,5,6].

The high sensitivity and specificity in the present investigation differ from findings reported in a study carried out in the United States involving individuals in different age groups (19 to 78 years), but the prevalence rate of dental caries was similar (63.8%) [2]. It should be stressed that the study cited only compared the decayed component, whereas the present investigation involved the entire DMFT index. Nonetheless, the difference between studies is likely

Table 2. Description and correlation of clinically determined and self-reported Decayed, Missing and Filled Teeth (DMFT) index of members of 1993 birth cohort at 18 years of age, Pelotas, Rio Grande do Sul, Brazil, 2012.

	Clinical Mean (95% CI)	Self-reported mean (95% CI)	Intraclass correlation coefficient	p-value
DMFT	2.06 (1.92; 2.21)	1.75 (1.68; 1.82)	0.50	<0.001
Decayed component	0.79 (0.71; 0.88)	0.63 (0.59; 0.67)	0.43	<0.001
Missing component	0.33 (0.29; 0.38)	0.28 (0.25; 0.30)	0.61	<0.001
Filled component	0.94 (0.84; 1.04)	0.84 (0.79; 0.89)	0.58	<0.001

doi:10.1371/journal.pone.0106382.t002

Table 3. Prevalence of clinically determined and self-reported dental caries (DMFT ≥ 1) with estimates of sensitivity, specificity, positive predictive value, negative predictive value, absolute bias, relative bias and inflation factor for members of 1993 birth cohort at 18 years of age, Pelotas, Rio Grande do Sul, Brazil, 2012.

Prevalence of dental caries						
Clinical % (95% CI)		Self-reported % (95% CI)			p-value¹	
66.5(63.6; 69.3)		60.3(58.8; 61.8)			0,001	
Dental caries determined by clinical (gold standard) and self-reported methods						
Sensitivity % (95% CI)	Specificity % (95% CI)	PPV² % (95% CI)	NPV³ % (95% CI)	Absolute bias⁴	Relative bias⁵	Inflation factor⁶
81.8(78.7; 84.7)	78.1(73.3; 82.4)	88(85.2; 90.5)	68.6(63.7; 73.2)	-6.2	9.3	1.1

¹chi-squared test.

²positive predictive value.

³negative predictive value.

⁴absolute bias = tested prevalence – gold standard prevalence.

⁵relative bias = underestimated true prevalence = absolute bias/gold standard prevalence X 100.

⁶inflation factor = gold standard prevalence/tested prevalence.

doi:10.1371/journal.pone.0106382.t003

due to the age groups analysed, which may have exerted an influence on the quality of the self-reported information, as discussed previously. Sensitivity is dependent on the prevalence of a disease [20]. Moreover, socioeconomic issues and health-related behaviour exert an influence on the quality of self-reported information regarding the health of a population and consequently affect the degree of sensitivity. The positive predictive value is affected by these same factors. In the analysis of adolescent's schooling, higher sensitivity and positive predictive values were found among those with a greater level of schooling and those who visited the dentist in the previous year. Thus, different results can be found in different socioeconomic contexts, indicating that the findings of the present study should be analysed with caution, considering the socioeconomic profile of Brazil.

In the analysis of the unadjusted prevalence ratios for socio-demographic aspects and oral health habits, some variables were only associated with the clinical determination of caries and others were only associated with self-reported caries. This indicates that

self-reported information should be used with caution in epidemiological studies with the aim of establishing significant associations between risk factors and dental caries. None of the studies consulted in the literature performed such comparisons, with the exception of studies addressing self-reported information on periodontal disease [18].

The advantage of using self-reported information for knowledge on the situation of dental caries in adolescents resides in the fact that the data collection questionnaire can be administered by any trained interviewer and does not require a dental professional. For oral health professionals, especially those linked to public healthcare services, such information is of extreme importance to the planning and monitoring of oral health policies and actions.

The quality of the information obtained in the present study should be stressed, as the data came from a sample from a cohort study, with due methodological care taken to confer a high degree of reliability to the information obtained.

Table 4. Estimates of sensitivity, specificity, positive predictive value and negative predictive value for clinically determined and self-reported dental caries (DMFT ≥ 1) according to gender, adolescent's schooling, household income and visit to dentist in previous year in 1993 birth cohort at 18 years of age, Pelotas, Rio Grande do Sul, Brazil, 2012.

	Sensitivity % (95% CI)	Specificity % (95% CI)	PPV¹ % (95% CI)	NPV² % (95% CI)	Accuracy
Female	87.3 (83.2; 90.7)	70.7(62.7; 77.8)	86.8 (82.7; 90.2)	71.6 (63.6; 78.7)	82.1
Male	76.4 (71.5; 80.9)	84.1 (78.1; 89.0)	89.5 (85.4; 92.8)	66.8 (60.4; 72.8)	78.2
Lesser schooling of adolescent (≤ 4 years of study)	75.0 (56.8; 88.5)	65.0 (40.8; 84.6)	77.4 (58.9; 90.4)	61.9 (38.4; 81.9)	71.1
Greater schooling of adolescent (≥ 12 years of study)	82.6 (61.2; 95.0)	92.6 (75.7; 99.1)	90.5 (69.6; 98.8)	86.2 (68.2; 96.1)	88.0
Lower household income (1st tertile)	84.5 (79.8; 88.5)	75.4 (67.2; 82.4)	87.9 (83.4; 91.5)	69.7 (61.5; 77.0)	83.9
Higher household income (3rd tertile)	78.0 (71.3; 83.8)	83.5 (74.9; 90.1)	89.3 (83.4; 93.6)	68.3 (59.4; 76.3)	80.0
Visited dentist in previous year	81.9 (77.6; 85.6)	78.0 (70.7; 84.2)	89.8 (86.1; 92.8)	64.6 (57.4; 71.3)	80.7
Did not visit dentist in previous year	80.8 (75.9; 86.0)	78.2 (71.4; 84.0)	85.9 (81.3; 89.8)	72.5 (65.7; 78.7)	80.4

¹positive predictive value.

²negative predictive value.

doi:10.1371/journal.pone.0106382.t004

Table 5. Prevalence ratios (95% CI) for exploratory variables of clinically determined and self-reported dental caries (DMFT ≥1) in 1993 cohort at 18 years of age, Pelotas, Rio Grande do Sul, Brazil, 2012.

	Clinical DMFT	Self-reported DMFT
Gender	p = 0.003	p < 0.001
Male	1.0	1.0
Female	1.2 (1.1; 1.4)	1.25 (1.2; 1.4)
Adolescent's schooling (years of study)	p < 0.001	p < 0.001
≤4	2.1 (1.2; 3.6)	1.0 (0.9; 1.3)
5–8	2.5 (1.6; 4.0)	1.3 (1.1; 1.6)
9–11	1.8 (1.2; 3.6)	1.3 (1.0; 1.7)
≥12	1.0	1.0
Household income (based on minimum wage)	p = 0.05	p = 0.27
1 st tertile	1.2 (1.0; 1.5)	1.2 (1.0; 1.4)
2 nd tertile	1.1 (0.9; 1.3)	1.1 (0.9; 1.3)
3 rd tertile	1.0	1.0
Visit to dentist in previous year	p = 0.001	p = 0.098
No	1.0	1.0
Yes	1.3 (1.2; 1.5)	1.1 (1.0; 1.3)
Daily tooth brushing frequency	p = 0.81	p = 0.007
<2 times	1.1 (0.7; 1.6)	1.3 (1.1; 1.6)
≥2 times	1.0	1.0
Self-rated oral health	p < 0.001	p < 0.001
Very good/good	1.0	1.0
Fair	1.4 (1.2; 1.6)	1.7 (1.4; 1.9)
Poor/very poor	2.1 (1.8; 2.6)	(2.6; 3.8)

doi:10.1371/journal.pone.0106382.t005

Based on the present findings, self-reported information on dental caries using the DMFT index requires further studies prior to its use in the analysis of risk factors, but is valid for population-based health surveys with the aim of planning and monitoring oral health actions directed at adolescents without the need to submit the population to clinical exams for the diagnosis of dental caries.

Author Contributions

Conceived and designed the experiments: AERS AMBM MAP MCFA HG FFD. Performed the experiments: AERS AMBM MAP MCFA HG FFD. Analyzed the data: AERS AMBM MAP FVF. Contributed reagents/materials/analysis tools: AERS MCFA HG FFD. Contributed to the writing of the manuscript: AERS MCFA HG FFD FVF.

References

- Blicher B, Joshipura K, Eke P (2005) Validation of self-reported periodontal disease: A Systematic Review. *J Dent Res* 84: 881–890.
- Pitiphat W, Garcia RI, Douglass CW, Joshipura KJ (2002) Validation of self-reported oral health measures. *J Public Health Dent* 62: 122–128.
- Joshipura KJ, Pitiphat W, Douglass CW (2002) Validation of Self-reported periodontal measures among health professionals. *J Public Health Dent* 62: 115–121.
- Taylor GW, Borgnakke WS (2007) Self-Reported periodontal disease: validation in an epidemiological survey. *Periodontol* 78: 1407–1420.
- Axellsson G, Helgadottir S (1995) Comparison of oral health data from self-administered questionnaire and clinical examination. *Community Dent Oral Epidemiol* 23: 365–8.
- Pinelli C, Loffredo LCM (2007) Reproducibility and validity of self-perceived oral health conditions. *Clin Oral Invest* 11: 431–437.
- Heloe LA (1972) Comparison of dental health data obtained from questionnaires, interviews and clinical examination. *Scand J Dent Res* 80: 495–499.
- Reisine ST, Bailit HL (1980) Clinical oral health status and adult perceptions of oral health. *Soc Sci Med Med Psychol Med Sociol* 14A: 597–605.
- Kononen M, Lipasti J, Murtomaa H (1986) Comparison of dental information obtained from self-examination and clinical examination. *Community Dent Oral Epidemiol* 14: 258–260.
- Palmqvist S, Soderfeldt B, Arnbjerg D (1991) Self-assessment of dental conditions: validity of a questionnaire. *Community Dent Oral Epidemiol* 19: 249–51.
- Gilbert GH, Duncan RP, Kulley AM (1997) Validity of self-reported tooth counts during a telephone screening interview. *J Public Health Dent* 57: 176–180.
- Araújo CL, Menezes AMB, Vieira MFA, Neutzling MB, Gonçalves H, et al. (2010) The 11-year follow-up of the 1993 Pelotas (Brazil) birth cohort study: methods. *Cad Saúde Pública* 26: 1875–186.
- Victora CG, Halal PC, Araújo CL, Menezes AM, Wells JC, et al. (2008) Cohort profile: the 1993 Pelotas (Brazil) birth cohort study. *Int J Epidemiol*. 37: 704–9.
- Peres MA, Barros AJ, Peres KG, Araújo CL, Menezes AMB, et al. (2010) Oral health follow-up studies in the 1993 Pelotas (Brazil) birth cohort study: methodology and principal results. *Cad Saúde Pública* 26: 1990–1999.
- World Health Organization (1997) *Oral Health Surveys: Basic Methods*. Geneva: World Health Organization. 66p.
- Albandar JM (2011) Underestimation of periodontitis in NHANES surveys. *J Periodontol* 82: 337–341.
- Barros AJ, Hirakata VN (2003) Alternatives for logistic regression in cross-sectional studies: An empirical comparison of models that directly estimate the prevalence ratio. *BMC Med Res Methodol* 3: 21.
- Peres MA, Peres KG, Cascaes AM, Correia MB, Demarco FF, et al. (2012) Validity of Partial Protocols to Assess the Prevalence of Periodontal Outcomes and Associated Sociodemographic and Behavior Factors in Adolescents and Young Adults. *J Periodontol* 83: 369–378.
- Gooch BF, Dolan TA, Bourque LB (1989) Correlates of self-reported dental health status upon enrollment in the Rand Health Insurance Experiment. *J Dent Educ*. 53: 629–637.
- Szklarz M, Javier Nieto F (2004) *Epidemiology: Beyond the Basics*. Sudbury, MA: Jones and Bartlett Publishers. 309 p.
- Brasil. Ministério da Saúde. Departamento de Atenção Básica: Condições de Saúde Bucal da População Brasileira 2002–2003. Disponível: <http://dtr2001>.

- sauda.gov.br/editora/produtos/livros/pdf/05_0053_M.pdf. Acessado 25 de março de. 2012.
22. Brasil. Ministério da Saúde. Departamento de Atenção Básica: SB Brasil 2010-Pesquisa Nacional de Saúde Bucal. Disponível: http://189.28.128.100/dab/docs/geral/projeto_sb2010_relatorio_final.pdf. Acessado 25 de março de 2012.
23. Costa SM, Martins CC, Bonfim MLC, Zina LG, Paiva SM, et al. (2012) A Systematic Review of Socioeconomic Indicators and Dental Caries in Adults. *Int J Environ Res Public Health* 9: 3540–74.
24. Moreira RSM, Nico LS, Tomita NE, Ruiz T (2005) Oral health of Brazilian elderly: a systematic review of epidemiologic status and dental care access. *Cad Saúde Pública* 21: 1665–1675.