

PUBLISHED VERSION

Melanie Manetsch, Md. Mostafizur Rahman, Brijeshkumar S. Patel, Emma E. Ramsay, Nowshin N. Rumzhum, Hatem Alkhouri, Qi Ge, Alaina J. Ammit
Long-acting B2-agonists increase fluticasone propionate-induced mitogen-activated protein kinase phosphatase 1 (MKP-1) in airway smooth muscle cells
PLoS One, 2013; 8(3):e59635-1-e59635-9

© 2013 Manetsch et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Originally published at:

<http://doi.org/10.1371/journal.pone.0059635>

PERMISSIONS

<http://creativecommons.org/licenses/by/4.0/>



Attribution 4.0 International (CC BY 4.0)

This is a human-readable summary of (and not a substitute for) the [license](#).

[Disclaimer](#)



You are free to:

Share — copy and redistribute the material in any medium or format

Adapt — remix, transform, and build upon the material

for any purpose, even commercially.

The licensor cannot revoke these freedoms as long as you follow the license terms.

Under the following terms:



Attribution — You must give [appropriate credit](#), provide a link to the license, and [indicate if changes were made](#). You may do so in any reasonable manner, but not in any way that suggests the licensor endorses you or your use.

No additional restrictions — You may not apply legal terms or [technological measures](#) that legally restrict others from doing anything the license permits.

<http://hdl.handle.net/2440/98105>

Long-Acting β_2 -Agonists Increase Fluticasone Propionate-Induced Mitogen-Activated Protein Kinase Phosphatase 1 (MKP-1) in Airway Smooth Muscle Cells

Melanie Manetsch¹, Md. Mostafizur Rahman¹, Brijeshkumar S. Patel¹, Emma E. Ramsay¹,
Nowshin N. Rumzhum¹, Hatem Alkhouri¹, Qi Ge², Alaina J. Ammit^{1*}

¹ Respiratory Research Group, Faculty of Pharmacy, University of Sydney, Sydney, Australia, ² Woolcock Institute of Medical Research, University of Sydney, Sydney, Australia

Abstract

Mitogen-activated protein kinase phosphatase 1 (MKP-1) represses MAPK-driven signalling and plays an important anti-inflammatory role in asthma and airway remodelling. Although MKP-1 is corticosteroid-responsive and increased by cAMP-mediated signalling, the upregulation of this critical anti-inflammatory protein by long-acting β_2 -agonists and clinically-used corticosteroids has been incompletely examined to date. To address this, we investigated MKP-1 gene expression and protein upregulation induced by two long-acting β_2 -agonists (salmeterol and formoterol), alone or in combination with the corticosteroid fluticasone propionate (abbreviated as fluticasone) in primary human airway smooth muscle (ASM) cells *in vitro*. β_2 -agonists increased MKP-1 protein in a rapid but transient manner, while fluticasone induced sustained upregulation. Together, long-acting β_2 -agonists increased fluticasone-induced MKP-1 and modulated ASM synthetic function (measured by interleukin 6 (IL-6) and interleukin 8 (IL-8) secretion). As IL-6 expression (like MKP-1) is cAMP/adenylate cyclase-mediated, the long-acting β_2 -agonist formoterol increased IL-6 mRNA expression and secretion. Nevertheless, when added in combination with fluticasone, β_2 -agonists significantly repressed IL-6 secretion induced by tumour necrosis factor α (TNF α). Conversely, as IL-8 is not cAMP-responsive, β_2 -agonists significantly inhibited TNF α -induced IL-8 in combination with fluticasone, where fluticasone alone was without repressive effect. In summary, long-acting β_2 -agonists increase fluticasone-induced MKP-1 in ASM cells and repress synthetic function of this immunomodulatory airway cell type.

Citation: Manetsch M, Rahman MM, Patel BS, Ramsay EE, Rumzhum NN, et al. (2013) Long-Acting β_2 -Agonists Increase Fluticasone Propionate-Induced Mitogen-Activated Protein Kinase Phosphatase 1 (MKP-1) in Airway Smooth Muscle Cells. PLoS ONE 8(3): e59635. doi:10.1371/journal.pone.0059635

Editor: Kazuhiro Ito, National Heart and Lung Institute, United Kingdom

Received: August 31, 2012; **Accepted:** February 16, 2013; **Published:** March 22, 2013

Copyright: © 2013 Manetsch et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This research was funded by a project grant (570856) from the National Health and Medical Research Council of Australia (www.nhmrc.gov.au). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: alaina.ammit@sydney.edu.au

Introduction

Asthma is a chronic airway disease characterized by excessive inflammation. Corticosteroids are widely-used to repress inflammation in asthma and control of asthma symptoms can be improved by combination therapy with long-acting β_2 -agonists [1,2,3]. There are a number of mechanisms by which this enhanced anti-inflammatory effect might be achieved at the cellular level: these include control of histone acetylation, increased function of glucocorticoid receptor (GR) or the β_2 -adrenergic receptor, and upregulation of anti-inflammatory proteins (reviewed in [4]). Mitogen-activated protein kinase phosphatase 1 (MKP-1) is one such anti-inflammatory protein whose upregulation is increasingly recognized as an important molecular mechanism responsible for the improved anti-inflammatory action of corticosteroids and long-acting β_2 -agonists in combination.

MKP-1 is an immediate-early gene [5] that is corticosteroid-responsive [6] and contains a cAMP-responsive element (CRE) in its 5'-promoter [7,8]. We and others have demonstrated that the upregulation of MKP-1 by corticosteroids is responsible, in part,

for their anti-inflammatory actions in clinically-relevant airway cell types such as airway smooth muscle (ASM) and bronchial epithelial cells [9,10,11,12,13,14]. Moreover, MKP-1 mRNA expression and protein upregulation can be enhanced by long-acting β_2 -agonists [12,15]. The increased expression of MKP-1 may explain the beneficial effects of β_2 -agonists/corticosteroid combination therapies in the repression of inflammatory gene expression in asthma [12,16]. However the upregulation of this critical anti-inflammatory protein by long-acting β_2 -agonists and clinically-used corticosteroids has been incompletely examined to date.

To address this gap in knowledge, the aim of this study is to examine MKP-1 mRNA expression and protein upregulation induced by two long-acting β_2 -agonists (salmeterol and formoterol), alone or in combination with the corticosteroid fluticasone propionate (abbreviated as fluticasone). We will assess the anti-inflammatory function of MKP-1 induced by these combination therapies on the synthetic function of primary human ASM cell *in vitro* by measuring repression of interleukin 6 (IL-6) and interleukin 8 (IL-8); two cytokines known to be upregulated in a MAPK-mediated manner [10,17,18].




Figure 1. Salmeterol increases fluticasone-induced MKP-1 protein upregulation. (A) Growth-arrested ASM cells were treated with vehicle, salmeterol (100 nM), fluticasone (1 nM), salmeterol (100 nM)+fluticasone (1 nM), for 0, 0.5, 1, 2, 4, 8, and 24 h. MKP-1 protein (molecular weight:~39 kDa) was quantified by Western blotting, using α -tubulin as the loading control (molecular weight: 55~kDa), where (A) illustrates representative Western blots and (B) demonstrates densitometric analysis where results are expressed as fold increase over 0 h (mean+SEM values from n=6–8 primary ASM cell cultures). Statistical analysis was performed using two-way ANOVA then Bonferroni's post-test (where * indicates a significant effect of salmeterol or fluticasone on MKP-1 protein upregulation, compared to vehicle-treated cells, and § indicates a significant effect of salmeterol on fluticasone-induced MKP-1 ($P<0.05$)). doi:10.1371/journal.pone.0059635.g001

Materials and Methods

Cell Culture

Human bronchi were obtained from patients undergoing surgical resection for carcinoma or lung transplant donors in accordance with procedures approved by the Sydney South West Area Health Service and the Human Research Ethics Committee of the University of Sydney. ASM cells were dissected, purified and cultured as previously described by Johnson *et al.* [19]. A minimum of three different ASM primary cell cultures established from individual patients were used for each experiment.

Unless otherwise specified, all chemicals used in this study were purchased from Sigma-Aldrich (St. Louis, MO).

Western Blotting

MKP-1 was quantified by Western blotting using a rabbit polyclonal antibody against MKP-1 (C-19: Santa Cruz Biotechnology, Santa Cruz, CA), compared to α -tubulin used as the loading control (mouse monoclonal IgG₁, DM1A: Santa Cruz Biotechnology, Santa Cruz, CA). Primary antibodies were detected with goat anti-mouse or anti-rabbit HRP-conjugated




Figure 2. Formoterol increases fluticasone-induced MKP-1 protein upregulation. (A) Growth-arrested ASM cells were treated with vehicle, formoterol (10 nM), fluticasone (1 nM), formoterol (10 nM)+fluticasone (1 nM), for 0, 0.5, 1, 2, 4, 8, and 24 h. MKP-1 protein (molecular weight:~39 kDa) was quantified by Western blotting, using α -tubulin as the loading control (molecular weight:~55 kDa), where (A) illustrates representative Western blots and (B) demonstrates densitometric analysis where results are expressed as fold increase over 0 h and results for fluticasone (from Figure 1B) shown graphically for comparative purposes (mean+SEM values from $n=3-8$ primary ASM cell cultures). Statistical analysis was performed using two-way ANOVA then Bonferroni's post-test (where * indicates a significant effect of formoterol or fluticasone on MKP-1 protein upregulation, compared to vehicle-treated cells, and § indicates a significant effect of formoterol on fluticasone-induced MKP-1 ($P<0.05$)). doi:10.1371/journal.pone.0059635.g002

secondary antibodies (Cell Signaling Technology) and visualized by enhanced chemiluminescence (PerkinElmer, Wellesley, MA).

Real-time RT-PCR

Total RNA was extracted using the RNeasy Mini Kit (Qiagen Australia, Doncaster, VIC, Australia) and reverse transcription performed by using the RevertAid First strand cDNA. MKP-1, IL-6 and IL-8 mRNA levels were measured using real-time RT-PCR on an ABI Prism 7500 (Applied Biosystems, Foster City, CA) with the MKP-1 (DUSP1: Hs00610256_g1), IL-6 (Hs00174131_m1) or IL-8 (Hs00174103_m1) TaqMan[®] Gene Expression Assays and the eukaryotic 18S rRNA endogenous control probe (Applied

Biosystems) subjected to the following cycle parameters: 50°C for 2 min, 1 cycle; 95°C for 10 min, 1 cycle; 95°C for 15 s, 60°C for 1 min, 40 cycles and mRNA expression (fold increase) quantified by delta delta Ct calculations.

ELISAs

Cell supernatants were collected and stored at -20°C for later analysis by ELISA. IL-6 and IL-8 ELISAs were performed according to the manufacturer's instructions (BD Biosciences Pharmingen, San Diego, CA).




Figure 3. Long-acting β_2 -agonists increase fluticasone-induced MKP-1 mRNA expression. Growth-arrested ASM cells were treated for 1 h with vehicle, fluticasone (1 nM), salmeterol (100 nM), formoterol (10 nM), salmeterol (100 nM)+fluticasone (1 nM), formoterol (10 nM)+fluticasone (1 nM). MKP-1 mRNA expression was quantified by real-time RT-PCR and results expressed as fold increase compared to vehicle-treated cells (mean \pm SEM values from $n=9$ primary ASM cell cultures). Statistical analysis was performed using repeated measures ANOVA with Bonferroni's multiple comparison test (where * denotes a significant increase in MKP-1 mRNA expression and § denotes a significant effect of the β_2 -agonists on fluticasone-induced MKP-1 mRNA expression ($P<0.05$)).

doi:10.1371/journal.pone.0059635.g003

MKP-1 siRNA Knock-down

Protein levels of MKP-1 were reduced by RNA interference and the effect on cytokine secretion examined. ASM cells were transiently transfected using nucleofection with 1 μ g MKP-1-specific SMART pool siRNA, consisting of a pool of four individual siRNA from Dharmacon (Dharmacon, Lafayette, CO) or a scrambled siRNA control, using methods established in our previous publication [10]. Briefly, ASM cells were transfected with the Nucleofector (Amaxa, Koln, Germany), using the basic kit for primary smooth muscle cells with the manufacturer's optimized protocol of P-024. ASM cells were plated for 16 h after transfection, before being growth-arrested for a further 24 h. Cells were pretreated for 1 h with formoterol+fluticasone (along with relevant controls) before stimulation for 24 h with TNF α (10 ng/ml). Supernatants were then removed for IL-6 and IL-8 protein secretion by ELISA and lysates utilized for MKP-1 Western blotting (compared to α -tubulin as a loading control).

Statistical Analysis

Statistical analysis was performed using either Student's unpaired t tests, repeated measured ANOVA or two-way ANOVA followed by Bonferroni's multiple comparison tests. P values <0.05 were sufficient to reject the null hypothesis for all analyses.

Results

Long-acting β_2 -agonists Increase Fluticasone-induced MKP-1 Protein Upregulation.

Although MKP-1 is known to be corticosteroid-responsive [10,20] and increased by cAMP-mediated signalling [20] in ASM cells, the upregulation of this critical anti-inflammatory protein by long-acting β_2 -agonists and clinically-used corticosteroids has been incompletely examined to date. To address this, we examine here for the first time, the effect of treating ASM cells with fluticasone, alone or in combination with the long-acting β_2 -agonists - salmeterol (100 nM: Figure 1) or formoterol (10 nM: Figure 2). These concentrations of β_2 -agonists were chosen for the current

study as they had been used previously in *in vitro* investigations of MKP-1 gene expression by Kaur *et al.* [12].

We first examined the temporal kinetics of MKP-1 protein upregulation in ASM cells treated for up to 24 h with salmeterol or fluticasone, alone or in combination (compared to vehicle-treated controls). As shown in Figure 1A, salmeterol alone induced rapid, but transient upregulation of MKP-1 with a peak at 1 h. In contrast, fluticasone alone induced a sustained increase in MKP-1 protein upregulation; in accordance with our earlier report [10]. This was confirmed by densitometry (as shown in Figure 1B) where salmeterol induced significantly greater levels of MKP-1 protein (4.8 ± 1.0 -fold) than cells treated with vehicle alone (1.7 ± 0.2 -fold) at 1 h ($P<0.05$). While MKP-1 upregulation induced by salmeterol was transient (returning back to basal levels at 4 h), fluticasone significantly increased MKP-1 protein levels by 4.4 ± 0.9 -fold at 1 h and this increase was sustained for up to 24 h (3.9 ± 0.8 -fold)($P<0.05$). When we then treat cells with salmeterol in combination with fluticasone (Figure 1A), the temporal pattern of MKP-1 protein upregulation reflects the combined contribution of the β_2 -agonist and corticosteroid. This is best demonstrated by densitometric analysis (Figure 1B), where both drugs in combination are shown to act in an additive manner and salmeterol added in combination with fluticasone induces significantly greater levels of MKP-1 protein (8.2 ± 1.4 -fold), than fluticasone alone at 1 h (4.4 ± 0.9 -fold) ($P<0.05$).

We then compared the effect of formoterol on fluticasone-induced MKP-1 (Figure 2). In corroboration of our earlier report [20], formoterol alone induces a transient increase in MKP-1 upregulation with a peak at 1–2 h (Figure 2A). Importantly, Western blotting (Figure 2A) revealed that when formoterol and fluticasone are added in combination, the peak of MKP-1 upregulation that appears at 1–2 h after the long-acting β_2 -agonist and enhances the sustained pattern of MKP-1 upregulation observed with fluticasone treatment alone. This is supported by densitometric analysis where formoterol significantly increased fluticasone-induced MKP-1 at 1 h (11.3 ± 2.1 -fold) and 2 h (10.1 ± 1.4 -fold), compared to fluticasone alone at the same time points (4.4 ± 0.9 - and 3.9 ± 0.7 -fold, respectively) ($P<0.05$).




Figure 4. Effect of long-acting β_2 -agonists on IL-6 and IL-8 mRNA expression and secretion. (A, B, C, D) Growth-arrested ASM cells were treated with vehicle, salmeterol (100 nM) or formoterol (10 nM). (A) IL-6 and (B) IL-8 mRNA expression was quantified by real-time RT-PCR at 1 h and results expressed as fold increase compared to vehicle-treated cells (mean+SEM values from $n=9$ primary ASM cell cultures). (C) IL-6 and (D) IL-8 secretion was measured by ELISA at 24 h and results are expressed as fold increase compared to vehicle-treated cells (where IL-6 and IL-8 protein levels in cells treated with vehicle were 295.0 ± 34.8 and 26.3 ± 4.6 pg/ml, respectively (mean+SEM values from $n=6$ primary ASM cell cultures)). Statistical analysis was performed using repeated measures ANOVA with Bonferroni's multiple comparison test (where * denotes a significant increase in mRNA expression or secretion ($P < 0.05$)). (E, F) Growth-arrested ASM cells were treated with vehicle, dibutyryl cAMP (1 mM), forskolin (10 μ M) and (E) IL-6 and (F) IL-8 mRNA expression was quantified by real-time RT-PCR at 1 h. Results are expressed as fold increase compared to vehicle-treated cells (mean+SEM values from $n=3$ primary ASM cell cultures). Statistical analysis was performed using the Student's unpaired t test where * denotes a significant increase in mRNA expression or secretion ($P < 0.05$). doi:10.1371/journal.pone.0059635.g004

Long-acting β_2 -agonists Increase Fluticasone-induced MKP-1 mRNA Expression

MKP-1 is an immediate-early gene and we previously reported that the corticosteroid dexamethasone and the long-acting β_2 -agonist formoterol rapidly increased MKP-1 gene expression by 1 h [20]. In the current study we extend these observations and perform a side-by-side comparison of MKP-1 mRNA upregulation in response to the long-acting β_2 -agonists salmeterol and formoterol, and the corticosteroid fluticasone, alone and in combination. While MKP-1 gene expression induced by flutica-

none (5.7 ± 1.2 -fold, compared to vehicle-treated cells) trended towards significance, salmeterol and formoterol added alone significantly increased MKP-1 gene expression by 10.2 ± 2.8 -fold and 16.4 ± 3.2 -fold, respectively ($P < 0.05$). Importantly, when long-acting β_2 -agonists were added in combination with fluticasone they significantly augmented steroid-induced MKP-1 mRNA expression ($P < 0.05$). As shown in Figure 3, salmeterol increased MKP-1 expression to 16.4 ± 3.2 -fold and formoterol to 21.3 ± 3.5 -fold, compared to fluticasone alone (5.7 ± 1.2 -fold, respectively) ($P < 0.05$).




Figure 5. The effect of long-acting β_2 -agonists and fluticasone, alone or in combination, on repression of TNF α -induced IL-6 and IL-8 secretion. Growth-arrested ASM cells were pretreated for 1 h with vehicle, salmeterol (100 nM), or formoterol (10 nM), alone or in combination with fluticasone (1 nM), before stimulation for 24 h with TNF α (10 ng/ml). (A) IL-6 and (B) IL-8 secretion was measured by ELISA and results expressed as (A) % TNF α -induced IL-6 secretion or (B) % TNF α -induced IL-8 secretion (where TNF α -induced IL-6 and IL-8 protein secretion was 5153.9 ± 618.9 and 5913.1 ± 480.0 pg/ml, respectively (mean \pm SEM values from $n=6$ primary ASM cell cultures)). Statistical analysis was performed using repeated measures ANOVA with Bonferroni's multiple comparison test where * denotes a significant increase in IL-6 secretion and § denotes a significant repression of TNF α -induced cytokine secretion ($P < 0.05$). doi:10.1371/journal.pone.0059635.g005

Effect of Long-acting β_2 -agonists on IL-6 and IL-8 mRNA Expression and Secretion

As the final aim of the current study was to demonstrate the anti-inflammatory function of MKP-1 on ASM synthetic function by measuring repression of IL-6 and IL-8 secretion induced by TNF α (see next section), it was first important to examine the effects of β_2 -agonists alone on secretion of these cytokines by ASM cells. Interestingly, IL-6 (like MKP-1) is cAMP-responsive. In this way, we show that salmeterol and formoterol significantly enhance IL-6 mRNA expression at 1 h (Figure 4A) and that formoterol increases IL-6 secretion from ASM cells at 24 h (Figure 4C) ($P < 0.05$). These protein secretion results are in confirmation of our earlier study [21] and in accord with the relative ability of these β_2 -agonists to elevate cAMP in ASM cells [21]. The involvement of cAMP/adenylate cyclase-mediated pathway in IL-6 mRNA expression is further substantiated by treatment of ASM cells with the cell-permeable cAMP analogue, dibutyryl cAMP, and the adenylylate cyclase activator, forskolin. As shown in Figure 4E, dibutyryl cAMP and forskolin increased IL-6 mRNA expression by 2.4 ± 0.4 -fold and 18.8 ± 1.7 -fold, respectively, compared to vehicle control ($P < 0.05$). IL-8, in contrast, is not regulated by cAMP and thus long-acting β_2 -agonists do not

increase IL-8 mRNA expression (Figure 4B), nor protein secretion (Figure 4D) in ASM cells. Moreover, dibutyryl cAMP and forskolin are without effect on IL-8 mRNA expression (Figure 4F).

The Effect of Long-acting β_2 -agonists and Fluticasone, Alone or in Combination, on Repression of TNF α -induced IL-6 and IL-8 Secretion

Finally, the effect of MKP-1 upregulation induced by β_2 -agonists, alone or in combination with the corticosteroid fluticasone, was demonstrated by pretreating ASM cells for 1 h before stimulation with TNF α . As shown in Figure 5A, and explained by the cAMP-responsiveness of IL-6 (described earlier), formoterol significantly enhanced TNF α -induced IL-6 secretion. Despite this, when β_2 -agonists were added in combination with fluticasone they still significantly repressed TNF α -induced IL-6 ($P < 0.05$). IL-8, on the other hand, is not cAMP-responsive; thus, β_2 -agonists did not significantly increase TNF α -induced IL-8 (Figure 5B). Interestingly, fluticasone alone did not repress TNF α -induced IL-8, while salmeterol, or formoterol, in combination with fluticasone significantly inhibited IL-6 secretion (Figure 5B). The combination of the long-acting β_2 -agonists with fluticasone, has a statistically greater effect than fluticasone alone on TNF α -induced IL-8 secretion ($P < 0.05$). Taken together, our results indicate that long-acting β_2 -agonists increase fluticasone-induced MKP-1 in an additive manner and in this way MKP-1 acts to repress cytokine secretion from ASM cells.

In preliminary studies, we then knocked-down MKP-1 with siRNA in an attempt to reverse the repression of cytokine secretion by formoterol and fluticasone, in combination. Although MKP-1 siRNA successfully reduced protein levels of MKP-1 in ASM cells (Figure 6A) we were unable to reverse the inhibition of TNF α -induced IL-6 (Figure 6B) or IL-8 (Figure 6C) secretion by formoterol and fluticasone. Investigations are now ongoing to examine the hypothesis that in the absence of MKP-1, the p38 MAPK-dependent RNA destabilizing factor – tristetraprolin (TTP) – may serve in a compensatory manner to repress inflammation. However further studies will be required to fully understand how these critical anti-inflammatory molecules are intrinsically intertwined and act together as a fail-safe mechanism to regulate inflammation in health and disease.

Discussion

This study demonstrates that long-acting β_2 -agonists increase fluticasone-induced MKP-1 mRNA expression and protein upregulation in ASM cells and together these asthma drugs have significant effects on cytokine secretion from ASM, an important immunomodulatory cell type in asthma.

Long-acting β_2 -agonists mediate their actions via interaction with the β_2 -adrenergic receptor, activation of adenylylate cyclase and production of cAMP. Gene transcription is then induced in genes that contain cAMP-responsive elements (CRE) in their 5'-promoter. MKP-1 has a CRE-containing promoter [7,8] and we recently reported that the long-acting β_2 -agonist formoterol induced MKP-1 mRNA and protein expression in a cAMP-mediated manner via the β_2 -adrenergic receptor-protein kinase A pathway [15]. MKP-1 gene expression and protein upregulation was rapid and peaked by 1 h [15]. We confirm these studies here and extend them to show that another commonly-used long-acting β_2 -agonist, salmeterol, also rapidly upregulates MKP-1 in a similar manner. Taken together with the demonstration that MKP-1 can be upregulated by cAMP analogues [20] these results present a unifying explanation that agonists that act to increase cAMP will increase MKP-1. Indeed, this has proven to be the case




Figure 6. Effect of MKP-1 siRNA. ASM cells were transiently transfected using nucleofection with scrambled control or MKP-1 siRNA, growth-arrested, then pretreated for 1 h with formoterol (10 nM)+fluticasone (1 nM) (along with relevant controls) before stimulation for 24 h with TNF α (10 ng/ml). Supernatants were then removed for IL-6 and IL-8 protein secretion by ELISA and lysates utilized for MKP-1 Western blotting (compared to α -tubulin as a loading control). (A) illustrates a representative Western blot, while (B, C) demonstrates results expressed as % TNF α -induced (B) IL-6 or (C) IL-8 secretion in cells transfected with scrambled control siRNA (mean+SEM values from n=6 primary ASM cell cultures). doi:10.1371/journal.pone.0059635.g006

for sphingosine 1-phosphate (S1P), a bioactive sphingolipid known to be elevated in asthma [22]. S1P acts via G_s-coupled receptors to increase cAMP [22] and we recently demonstrated that S1P increased MKP-1 in a cAMP-dependent manner to serve an important role as a negative feedback controller and restrain the extent and duration of pro-inflammatory cellular signalling [20].

Importantly, when long-acting β_2 -agonists are added in combination with the corticosteroid fluticasone they significantly and rapidly upregulated MKP-1 protein and mRNA expression. The temporal pattern of MKP-1 protein upregulation reflects the additive contribution of each asthma drug. That is, β_2 -agonists rapidly, but transiently, upregulate MKP-1 at 1–2 h, while corticosteroids induced a sustained increase in MKP-1 protein. Thus, when added together they upregulate MKP-1 early (by 1 h) but in a sustained manner. This additive upregulation was supported by mRNA levels. These results suggest a molecular basis for the added anti-inflammatory effects of both asthma drugs in combination.

To support the above assertion that increased expression of MKP-1 may explain the beneficial effects of β_2 -agonists/corticosteroid combination therapies in the repression of inflammatory gene expression in asthma [16], we performed *in vitro* studies measuring secretion of IL-6 and IL-8 from ASM cells after stimulation with TNF α . It is important to note that IL-6 is a cAMP-responsive gene. The 5'-promoter for IL-6 contains a CRE [21] and can be induced by cAMP analogues, as well as short-acting and long-acting β_2 -agonists [21,23]. Herein we confirmed the cAMP-responsiveness of IL-6 by demonstrating significant IL-

6 mRNA expression induced by the cell-permeable analogue, dibutyryl cAMP, as well as the adenylate cyclase activator, forskolin. In contrast, cAMP is only a weak activator of IL-8 in ASM cells [24] and, as highlighted by Holden *et al.* in lung epithelial cells [25], the promoter for IL-8 does not contain CRE consensus sequences [26] and that transcriptional regulation may involve cross-talk between transcription factors. Our current study demonstrates the lack of effect of cAMP-elevating agents on IL-8 expression in ASM cells.

Thus, due to the cAMP-dependence of IL-6 expression, the long-acting β_2 -agonist formoterol significantly enhanced TNF α -induced IL-6 secretion. This corroborated our earlier report [21] and underscored the importance of the study by Holden *et al.* [25] where the authors stated that enhancement of IL-6 (and IL-8) release by β_2 -agonists may contribute to the deleterious effects of β_2 -agonist used monotherapy. Notably, when added in combination with fluticasone, β_2 -agonists significantly repressed IL-6 secretion, but not to a greater extent than fluticasone alone. These results are in agreement with Holden *et al.* where increased inflammatory cytokine released was reversed by corticosteroid action [25].

The repressive effect on TNF α -induced IL-8 were more clear-cut, as long-acting β_2 -agonists added on their own did not increase IL-8 expression. Importantly, we observed that fluticasone alone was without repressive effect on TNF α -induced IL-8 secretion, but that β_2 -agonists added in combination significantly repressed TNF α -induced IL-8. This enhanced repressive effect of β_2 -agonists and corticosteroids correlates with the increased MKP-1

expression by both asthma drugs in combination. Why the combined effects of LABA/fluticasone on MKP-1 result in an additive suppressive effect on IL-8, but not on IL-6, cannot be completely delineated at this stage, but may reflect the relative MAPK-dependent contribution to IL-6 and IL-8 secretion. However, our *in vitro* results support those recently published *in vivo* in mild asthmatics [27], where formoterol in combination with corticosteroids improved clinical symptoms (FEV₁) and reduced IL-8 in a manner linked to MKP-1 upregulation. Interestingly, it has been previously shown by Knobloch *et al.* [28] that the corticosteroid dexamethasone suppresses IL-8 in ASM cells of non-smokers and smokers, but not in those of COPD; whether MKP-1 is involved warrants further investigation.

Our preliminary studies have raised the hypothesis that in the absence of MKP-1 (knocked-down by siRNA), the p38 MAPK-dependent TTP may be upregulated and serve as a compensatory anti-inflammatory factor. TTP is an immediate-early gene [29] that is rapidly and robustly expressed via p38 MAPK signalling [30,31,32,33]. Moreover, TTP's function as an mRNA destabilising factor is dictated by phosphorylation on two key serine residues (S52 and S178). Further studies investigating the regulation of the RNA destabilising factor TTP by MKP-1, and *visa versa*, in ASM cells are warranted.

References

- Pauwels RA, Lofdahl CG, Postma DS, Tattersfield AE, O'Byrne P, et al. (1997) Effect of inhaled formoterol and budesonide on exacerbations of asthma. Formoterol and Corticosteroids Establishing Therapy (FACET) International Study Group. *New England Journal of Medicine* 337: 1405–1411.
- O'Byrne PM, Naya IP, Kallen A, Postma DS, Barnes PJ (2008) Increasing doses of inhaled corticosteroids compared to adding long-acting inhaled beta2-agonists in achieving asthma control. *Chest* 134: 1192–1199.
- Chung KF, Caramori G, Adcock IM (2009) Inhaled corticosteroids as combination therapy with beta-adrenergic agonists in airways disease: present and future. *European Journal of Clinical Pharmacology* 65: 853–871.
- Barnes PJ (2011) Biochemical basis of asthma therapy. *Journal of Biological Chemistry* 286: 32899–32905.
- Sun H, Charles CH, Lau LF, Tonks NK (1993) MKP-1 (3CH134), an immediate early gene product, is a dual specificity phosphatase that dephosphorylates MAP kinase *in vivo*. *Cell* 75: 487–493.
- Clark AR (2003) MAP kinase phosphatase 1: a novel mediator of biological effects of glucocorticoids? *Journal of Endocrinology* 178: 5–12.
- Kwak SP, Hakes DJ, Martell KJ, Dixon JE (1994) Isolation and characterization of a human dual specificity protein-tyrosine phosphatase gene. *Journal of Biological Chemistry* 269: 3596–3604.
- Sommer A, Burkhardt H, Keyse SM, Luscher B (2000) Synergistic activation of the mkp-1 gene by protein kinase A signaling and USF, but not c-Myc. *FEBS Lett* 474: 146–150.
- Issa R, Xie S, Khorasani N, Sukkar M, Adcock IM, et al. (2007) Corticosteroid inhibition of growth-related oncogene protein-alpha via mitogen-activated kinase phosphatase-1 in airway smooth muscle cells. *Journal of Immunology* 178: 7366–7375.
- Quante T, Ng YC, Ramsay EE, Henness S, Allen JC, et al. (2008) Corticosteroids reduce IL-6 in ASM cells via up-regulation of MKP-1. *Am J Respir Cell Mol Biol* 39: 208–217.
- Kang BN, Jude JA, Panettieri RA Jr, Walseth TF, Kannan MS (2008) Glucocorticoid regulation of CD38 expression in human airway smooth muscle cells: role of dual specificity phosphatase 1. *Am J Physiol Lung Cell Mol Physiol* 295: L186–193.
- Kaur M, Chivers JE, Gienbycz MA, Newton R (2008) Long-acting beta2-adrenoceptor agonists synergistically enhance glucocorticoid-dependent transcription in human airway epithelial and smooth muscle cells. *Mol Pharmacol* 73: 203–214.
- King EM, Holden NS, Gong W, Rider CF, Newton R (2009) Inhibition of NF-kappaB-dependent transcription by MKP-1: transcriptional repression by glucocorticoids occurring via p38 MAPK. *Journal of Biological Chemistry* 284: 26803–26815.
- Newton R, King EM, Gong W, Rider CF, Staples KJ, et al. (2010) Glucocorticoids inhibit IL-1beta-induced GM-CSF expression at multiple levels: roles for the ERK pathway and repression by MKP-1. *Biochemical Journal* 427: 113–124.
- Manetsch M, Ramsay E, King E, Seidel P, Che W, et al. (2012) Corticosteroids and beta(2) -agonists upregulate mitogen-activated protein kinase phosphatase 1: *in vitro* mechanisms. *British Journal of Pharmacology* 166: 2049–2059.
- Gienbycz MA, Kaur M, Leigh R, Newton R (2008) A Holy Grail of asthma management: toward understanding how long-acting beta(2)-adrenoceptor agonists enhance the clinical efficacy of inhaled corticosteroids. *British Journal of Pharmacology* 153: 1090–1104.
- Amrani Y, Ammit AJ, Panettieri RA Jr (2001) Tumor Necrosis Factor Receptor (TNFR) 1, but Not TNFR2, Mediates Tumor Necrosis Factor-alpha-Induced Interleukin-6 and RANTES in Human Airway Smooth Muscle Cells: Role of p38 and p42/44 Mitogen-Activated Protein Kinases. *Molecular Pharmacology* 60: 646–655.
- Henness S, van Thoor E, Ge Q, Armour CL, Hughes JM, et al. (2006) IL-17A acts via p38 MAPK to increase stability of TNF-alpha-induced IL-8 mRNA in human ASM. *American Journal of Physiology* 290: L1283–L1290.
- Johnson PR, McKay KO, Armour CL, Black JL (1995) The maintenance of functional activity in human isolated bronchus after cryopreservation. *Pulmonary Pharmacology* 8: 43–47.
- Che W, Manetsch M, Quante T, Rahman MM, Patel BS, et al. (2012) Sphingosine 1-phosphate induces MKP-1 expression via p38 MAPK- and CREB-mediated pathways in airway smooth muscle cells. *Biochimica et Biophysica Acta* 1823: 1658–1665.
- Ammit AJ, Lazaar AL, Irani C, O'Neill GM, Gordon ND, et al. (2002) TNF-alpha-induced secretion of RANTES and IL-6 from human airway smooth muscle cells: Modulation by glucocorticoids and beta-agonists. *American Journal of Respiratory Cell and Molecular Biology* 26: 465–474.
- Ammit AJ, Hastie AT, Edsall LC, Hoffman RK, Amrani Y, et al. (2001) Sphingosine 1-phosphate modulates human airway smooth muscle cell functions that promote inflammation and airway remodeling in asthma. *FASEB Journal* 15: 1212–1214.
- Ammit AJ, Hoffman RK, Amrani Y, Lazaar AL, Hay DWP, et al. (2000) TNF-alpha-induced secretion of RANTES and IL-6 from human airway smooth muscle cells: Modulation by cAMP. *American Journal of Respiratory Cell and Molecular Biology* 23: 794–802.
- Pang L, Knox AJ (2000) Synergistic inhibition by beta-agonists and corticosteroids on tumor necrosis factor-alpha-induced interleukin-8 release from cultured human airway smooth-muscle cells. *American Journal of Respiratory Cell and Molecular Biology* 23: 79–85.
- Holden NS, Rider CF, Bell MJ, Velayudhan J, King EM, et al. (2010) Enhancement of inflammatory mediator release by beta(2)-adrenoceptor agonists in airway epithelial cells is reversed by glucocorticoid action. *British Journal of Pharmacology* 160: 410–420.
- Roebuck KA (1999) Regulation of interleukin-8 gene expression. *Journal of Interferon and Cytokine Research* 19: 429–438.
- Essilfie-Quaye S, Ito K, Ito M, Kharitonov SA, Barnes PJ (2011) Comparison of Symbicort(R) versus Pulmicort(R) on steroid pharmacodynamic markers in asthma patients. *Respiratory Medicine* 105: 1784–1789.
- Knobloch J, Sibbing B, Jungck D, Lin Y, Urban K, et al. (2010) Resveratrol impairs the release of steroid-resistant inflammatory cytokines from human airway smooth muscle cells in chronic obstructive pulmonary disease. *Journal of Pharmacology and Experimental Therapeutics* 335: 788–798.

29. Carballo E, Lai WS, Blakeshear PJ (1998) Feedback inhibition of macrophage tumor necrosis factor- α production by tristetraprolin. *Science* 281: 1001–1005.
30. Mahtani KR, Brook M, Dean JL, Sully G, Saklatvala J, et al. (2001) Mitogen-activated protein kinase p38 controls the expression and posttranslational modification of tristetraprolin, a regulator of tumor necrosis factor α mRNA stability. *Molecular and Cellular Biology* 21: 6461–6469.
31. Tchen CR, Brook M, Saklatvala J, Clark AR (2004) The stability of tristetraprolin mRNA is regulated by mitogen-activated protein kinase p38 and by tristetraprolin itself. *Journal of Biological Chemistry* 279: 32393–32400.
32. Brook M, Tchen CR, Santalucia T, McIlrath J, Arthur JS, et al. (2006) Posttranslational regulation of tristetraprolin subcellular localization and protein stability by p38 mitogen-activated protein kinase and extracellular signal-regulated kinase pathways. *Molecular and Cellular Biology* 26: 2408–2418.
33. Hiiti E, Iakovleva T, Brook M, Deppenmeier S, Gruber AD, et al. (2006) Mitogen-activated protein kinase-activated protein kinase 2 regulates tumor necrosis factor mRNA stability and translation mainly by altering tristetraprolin expression, stability, and binding to adenine/uridine-rich element. *Molecular and Cellular Biology* 26: 2399–2407.